ON PAIRED ROOT SYSTEMS OF COXETER GROUPS

FU, XIANG

School of Mathematics and Statistics University of Sydney, NSW 2006, Australia

xifu9119@mail.usyd.edu.au xiangf@maths.usyd.edu.au

Preliminary version, March 18, 2021

ABSTRACT. This paper examines a systematic method to construct a pair of (inter-related) root systems for arbitrary Coxeter groups from a class of non-standard geometric representations. This method can be employed to construct generalizations of root systems for a large family of linear groups generated by involutions. We then give a characterization of Coxeter groups, among these groups, in terms of such paired root systems. Furthermore, we use this method to construct and study the paired root systems for reflection subgroups of Coxeter groups.

1. Introduction

A Coxeter group W is an abstract group generated by a set of involutions R, called its Coxeter generators, subject only to certain braid relations. Despite the simplicity of this definition, there is a rich theory for Coxeter groups with non-trivial applications in a multitude of areas of mathematics and physics. When studying Coxeter groups, one of the most powerful tools we have at our disposal is the notion of root systems. In classical literature ([2, Ch.V, §4] or [21, §5.3–5.4], for example), the root system of a Coxeter group W is a geometric construction arising from the Tits representation of W. The Tits representation of W is an embedding of W into the orthogonal group of a certain bilinear form on a suitably chosen vector space V subject to the requirement that the W-conjugates of elements of R are mapped to reflections with respect to certain hyperplanes in V. In the case that W is finite, V is Euclidean (of dimension equal to the cardinality of R), and the root system of W simply consists of representative normal vectors for these hyperplanes. Those elements of the root system corresponding to the

²⁰¹⁰ Mathematics Subject Classification. 20F55 (20F10, 20F65).

Key words and phrases. Coxeter groups, reflection groups, Kac-Moody Lie algebras, root systems.

elements of R are known as *simple roots*, and in most classical literature ([2, Ch.V, §4] and [21, §5.3], for example), the simple roots are linearly independent.

Similar constructions of root systems can be extended to infinite Coxeter groups and Kac-Moody Lie algebras. However, the actual constructions of root systems differ depending whether the root systems are associated to Kac-Moody Lie algebras or infinite Coxeter groups. As discussed in the introduction of [17], while all definitions of root systems are related to a given bilinear form, the actual bilinear forms considered in the case of Kac-Moody Lie algebras are different from the ones in Coxeter groups. Furthermore, it is well known ([9] and [10, Ch.3) that within an arbitrary Coxeter group W, all of its reflection subgroups are themselves Coxeter groups, but in the literature ([10] or [11], for example), the construction of the root systems corresponding to such reflections subgroups as subsets of the root system of W requires special care. In particular, since a proper reflection subgroup may have strictly more Coxeter generators than the over-group (as seen in [17, Example 5.1]), the equivalent of the simple roots in these root systems need not be linearly independent, making the overall theory of root systems and root bases somewhat less uniform. Consequently, it seems profitable to develop a universal method for constructing root systems that is applicable to arbitrary Coxeter groups and their reflection subgroups, as well as to objects like Kac-Moody Lie algebras (in fact, to all groups with a so-called root group datum, as surveyed in [6]).

In [25], [7], [8] and [6], a number of more general notions of root systems have been studied. In these approaches, a pair of root systems are constructed in two vector spaces which are essentially algebraic duals of each other (apart from [25], the two vector spaces involved are explicitly required to be algebraic duals of each other, whereas in [25] the two vector spaces are linked by a non-degenerate bilinear pairing satisfying certain integrality conditions).

Recently, an approach taken in [12] and [14] generalizes those of [25], [7], [8] and [6]. In this approach, for an arbitrary Coxeter group, a pair of root systems are constructed in two vector spaces linked only by a bilinear pairing which does not require the non-degeneracy and integrality conditions of [25]. This particular approach allows more abstract geometry in the two root systems to take place (for example, the two representation spaces need not be algebraic duals of each other), whilst providing a unified theory of root systems, especially with respect to reflection subgroups (this last point is to be established in Section 3 of this present paper).

In this paper, we present a few results demonstrating the "universalness" of the notion of root systems in [12] and [14]. In fact, this new approach applies to a large family of linear groups generated by

involutions, and one of the key results of this paper (Theorem 2.8) shows that these groups are Coxeter groups only if the corresponding root systems decompose as disjoint unions of those roots generalizing the classical concept of positive roots and those roots generalizing the classical concept of negative roots. In fact, this result provides an alternative characterization for Coxeter groups, since it is well known that for any Coxeter group we may construct a root system that decomposes in the same way. This alternative characterization is implicitly suggested in the work of Matthew Dyer ([10]), and we are very grateful to Prof. Dyer for a large number of helpful suggestions leading to the development of this generalized notion of root systems.

The main body of this paper is organized into 2 sections, namely, Section 2 and Section 3. In Section 2 we develop a notion of root systems applicable to a large family of groups that are generated only by involutions, and we investigate when these root systems may decompose into disjoint unions of the so-called positive roots and the so-called negative roots, and we prove that such groups are Coxeter groups only if such decompositions take place (Theorem 2.3 and Theorem 2.8). In Section 3 we prove that the notion of root systems in [12] and [14] applies to all the reflection subgroups of any Coxeter group. In particular, we give a geometric characterization of the roots that correspond to the Coxeter generators of reflection subgroups (Proposition 3.15 and Proposition 3.22), and we show that these characterizations are precisely those allowing these roots to be the simple roots for the root systems of such reflection subgroups within the root systems of the respective over-groups.

Notation. If A is a subset of a real vector space then we define the positive linear cone of A, denoted PLC(A), to be the set

$$\left\{\sum_{a\in A} c_a a \mid c_a \geq 0 \text{ for all } a\in A, \text{ and } c_{a'}>0 \text{ for some } a'\in A \right\}.$$

Furthermore, we define $-A := \{ -v \mid v \in A \}$. Also, if B is a subset of a group G then $\langle B \rangle$ denotes the subgroup of G generated by B.

2. Decomposition of Root Systems and Coxeter Datum

Let V_1 and V_2 be vector spaces over the real field \mathbb{R} equipped with a bilinear pairing $\langle , \rangle \colon V_1 \times V_2 \to \mathbb{R}$. Let S be an indexing set, and suppose that $\Pi_1 := \{ \alpha_s \mid s \in S \} \subseteq V_1 \text{ and } \Pi_2 := \{ \beta_s \mid s \in S \} \subseteq V_2 \text{ are both in bijective correspondence with } S$. Further, suppose that Π_1 and Π_2 satisfy the following conditions:

- (D1) $\langle \alpha_s, \beta_s \rangle = 1$ for all $s \in S$;
- (D2) (i) $0 \notin PLC(\Pi_1)$ and $0 \notin PLC(\Pi_2)$.
 - (ii) $\alpha_s \notin PLC(\Pi_1 \setminus \{\alpha_s\})$ and $\beta_s \notin PLC(\Pi_2 \setminus \{\beta_s\})$ for each $s \in S$.

Observe that condition (D2) (i) implies that $\alpha_s \notin PLC(-\Pi_1 \setminus \{-\alpha_s\})$ and $\beta_s \notin PLC(-\Pi_2 \setminus \{-\beta_s\})$ for each $s \in S$. We remark that there do exist examples for which Π_1 (resp. Π_2) is linearly dependent, in which case necessarily some α_s (resp. β_s) will be expressible as a linear combination of $\Pi_1 \setminus \{\alpha_s\}$ (resp. $\Pi_2 \setminus \{\beta_s\}$) with coefficients of mixed signs.

Definition 2.1. For $s \in S$, define $\rho_1(s) \in GL(V_1)$ and $\rho_2(s) \in GL(V_2)$ by the rules

$$\rho_1(s)(x) := x - 2\langle x, \beta_s \rangle \alpha_s$$

for all $x \in V_1$, and

$$\rho_2(s)(y) := y - 2\langle \alpha_s, y \rangle \beta_s$$

for all $y \in V_2$. Further, we define, for each $i \in \{1, 2\}$:

$$R_{i} := \{ \rho_{i}(s) \mid s \in S \};$$

$$W_{i} := \langle R_{i} \rangle;$$

$$\Phi_{i} := W_{i}\Pi_{i};$$

$$\Phi_{i}^{+} := \Phi_{i} \cap PLC(\Pi_{i});$$

and

$$\Phi_i^- := -\Phi_i^+$$

For each $i \in \{1, 2\}$, and for each $s \in S$, we call $\rho_i(s)$ the reflections corresponding to s in V_i . We call Φ_i the root system for the Weyl group W_i realized in V_i , and we call Π_i the set of simple roots in Φ_i . Furthermore, we call Φ_i^+ the set of positive roots in Φ_i and Φ_i^- the set of negative roots in Φ_i .

Remark 2.2. For each $i \in \{1, 2\}$ and each $s \in S$ note that $\rho_i(s)$ is an involution with a -1-eigenvector of multiplicity 1. Furthermore, it is a consequence of condition (D2) that $\Phi_i^+ \cap \Phi_i^- = \emptyset$. Use \forall to denote disjoint unions, we have:

Theorem 2.3. Given conditions (D1) and (D2), the following are equivalent:

- $\begin{array}{ll} \text{(i)} \ \Phi_1 = \Phi_1^+ \uplus \Phi_1^-. \\ \text{(ii)} \ \Phi_2 = \Phi_2^+ \uplus \Phi_2^-. \end{array}$
- (iii) For all $s, t \in S$ the following three conditions are satisfied:
 - (D3) $\langle \alpha_s, \beta_t \rangle \leq 0$ and $\langle \alpha_t, \beta_s \rangle \leq 0$ whenever $s \neq t$.
 - (D4) $\langle \alpha_s, \beta_t \rangle = 0$ if and only if $\langle \alpha_t, \beta_s \rangle = 0$.
 - (D5) Either $\langle \alpha_s, \beta_t \rangle \langle \alpha_t, \beta_s \rangle = \cos^2 \frac{\pi}{m_{st}}$ for some integer $m_{st} \geq 2$, or else $\langle \alpha_s, \beta_t \rangle \langle \alpha_t, \beta_s \rangle > 1$.

It is a consequence of this theorem that if any of the equivalent conditions in it is satisfied then W_1 and W_2 are isomorphic Coxeter groups. To prove this theorem we shall need a few technical results first. These are essentially taken from [10], and for completeness, the relevant proofs are included here.

Let \mathscr{A} be a commutative \mathbb{R} -algebra, let $q^{1/2}$ and X be units of \mathscr{A} , and let $\gamma \in \mathbb{R}$. Define A, B to be 2×2 matrices over \mathscr{A} given by

$$A = \begin{pmatrix} -1 & 2\gamma q^{1/2}X \\ 0 & q \end{pmatrix} \qquad B = \begin{pmatrix} q & 0 \\ 2\gamma q^{1/2}X^{-1} & -1 \end{pmatrix}.$$

It is easily proved by induction on $n \in \mathbb{N}$ that

$$B(AB)^{n} = \begin{pmatrix} q^{n+1}p_{2n+1} & -q^{n+\frac{1}{2}}p_{2n}X\\ q^{n+\frac{1}{2}}p_{2n+2}X^{-1} & -q^{n}p_{2n+1} \end{pmatrix}$$
(2.1)

$$A(BA)^{n} = \begin{pmatrix} -q^{n}p_{2n+1} & q^{n+\frac{1}{2}}p_{2n+2}X\\ -q^{n+\frac{1}{2}}p_{2n}X^{-1} & q^{n+1}p_{2n+1} \end{pmatrix}$$
(2.2)

$$(BA)^{n} = \begin{pmatrix} -q^{n} p_{2n-1} & q^{n+\frac{1}{2}} p_{2n} X \\ -q^{n-\frac{1}{2}} p_{2n} X^{-1} & q^{n} p_{2n+1} \end{pmatrix}$$
 (2.3)

and

$$(AB)^n = \begin{pmatrix} q^n p_{2n+1} & -q^{n-\frac{1}{2}} p_{2n} X \\ q^{n+\frac{1}{2}} p_{2n} X^{-1} & -q^n p_{2n-1} \end{pmatrix}$$
 (2.4)

where $p_n \in \mathbb{R}$ $(n \in \{-1\} \cup \mathbb{N})$ are defined recursively by

$$p_{-1} = -1,$$
 $p_0 = 0,$ $p_{n+1} = 2\gamma p_n - p_{n-1} \quad (n \in \mathbb{N}).$ (2.5)

The solutions of the recurrence equation (2.5) is

$$p_{n} = \begin{cases} n & \gamma = 1\\ (-1)^{n+1}n & \gamma = -1\\ \frac{\sinh n\theta}{\sinh \theta} & \text{(where } \theta = \cosh^{1}\gamma) & |\gamma| > 1\\ \frac{\sin n\theta}{\sin \theta} & \text{(where } \theta = \cos^{-1}\gamma) & |\gamma| < 1. \end{cases}$$
 (2.6)

Proposition 2.4. ([10, Lemma 2.2]) Keep all the above notation.

- Conditions (1) and (2) below are equivalent:

 - (1) $p_n p_{n+1} \ge 0$ for all $n \in \mathbb{N}$; (2) $\gamma \in \{\cos \frac{\pi}{m} \mid m \in \mathbb{N}, m \ge 2\} \cup [1, \infty)$.
- (ii) If $\gamma = \cos \frac{k\pi}{m}$ for some $k, m \in \mathbb{N}$ with 0 < k < m then the matrices A and B satisfies the equation

$$ABA \cdot \cdot \cdot = BAB \cdot \cdot \cdot$$

where there are m factors on each side.

(iii) If q = 1 then the matrix AB has order m if $\gamma = \cos \frac{k\pi}{m}$ for some $k, m \in \mathbb{N}$ with 0 < k < m and gcd(m, k) = 1, and the matrix AB has infinite order otherwise.

Proof. (i): First assume that (1) holds. Observe that (2.5) yields that $p_1 = 1$ and $p_2 = 2\gamma$, hence $\gamma \geq 0$. Since (2) obviously holds if $\gamma \geq 1$, we may assume that $0 \leq \gamma < 1$. Choose θ so that $0 < \theta \leq \frac{\pi}{2}$ and $\cos \theta = \gamma$, and let m be the largest integer such that

$$0 < \theta < 2\theta < \dots < m\theta < \pi$$
.

Note that $m \geq 2$. Now if $m\theta \neq \pi$ then $\pi < (m+1)\theta < 2\pi$, and in view of (2.6) we have $p_m = \frac{\sin m\theta}{\sin \theta} > 0$, whereas $p_{m+1} = \frac{\sin(m+1)\theta}{\sin \theta} < 0$, contradicting (1). Hence $m\theta = \pi$ and $\gamma = \cos \frac{\pi}{m}$ for some integer $m \geq 2$, whence (2) holds as desired. Conversely, if (2) holds then it follows immediately from (2.6) that (1) holds.

(ii): If m = 2r is even then our task is to prove that $(AB)^r = (BA)^r$. It follows from (2.6) that $p_n = \frac{\sin(nk\pi/2r)}{\sin(k\pi/2r)}$, and hence, $p_{2r+1} = (-1)^k$ and $p_{2r-1} = (-1)^{k+1}$, while $p_{2r} = 0$. Then it follows from (2.3) and (2.4) that $(AB)^r = (BA)^r$.

If m = 2r+1 is odd then our task is to prove that $B(AB)^r = A(BA)^r$. In this case we find from (2.6) that $p_{2r+1} = 0$, while $p_{2r+2} = (-1)^k$ and $p_{2r} = (-1)^{k+1}$, and then the required result follows immediately from (2.2) and (2.1).

(iii): If $\gamma = \cos \frac{k\pi}{m}$ then it follows immediately from (ii) above that $(AB)^m = 1$, because $A^2 = B^2 = 1$ when q = 1. If 0 < n < m and $\gcd(k,m) = 1$, then (2.6) yields that $p_n = \frac{\sin(nk\pi/m)}{\sin(k\pi/m)} \neq 0$, and it then follows from (2.4) that $(AB)^n \neq 1$, proving that AB has order m. On the other hand, if γ is of any other form then it follows from (2.6) that $p_n \neq 0$ for all integer n > 0. Then it is clear from (2.4) that $(AB)^n \neq 1$ for all such n, proving that AB has infinite order.

Now we are ready to prove Theorem 2.3.

Proof of Theorem 2.3. We give a proof that (i) is equivalent to (iii). An entirely similar argument shows that (ii) and (iii) are also equivalent.

First we show that (iii) implies (i). Given conditions (D3), (D4) and (D5) of the present paper we observe that $\mathscr{C} := (S, V_1, V_2, \Pi_1, \Pi_2, \langle, \rangle)$ forms a Coxeter datum in the sense of [14], and hence (i) follows immediately from Lemma 3.2 of [14].

Conversely, suppose that $\Phi_1 = \Phi_1^+ \uplus \Phi_1^-$. We first prove that (D3) holds. Let $s, t \in S$ be distinct. By definition we have

$$\rho_1(t)\alpha_s = \alpha_s - 2\langle \alpha_s, \beta_t \rangle \alpha_t. \tag{2.7}$$

The condition $\Phi_1 = \Phi_1^+ \uplus \Phi_1^-$ implies that either

$$\rho_1(t)\alpha_s = \sum_{r \in S} c_r \alpha_r, \text{ where all } c_r \ge 0,$$
(2.8)

or else

$$\rho_1(t)\alpha_s = \sum_{r \in S} -c_r \alpha_r, \text{ where all } c_r \ge 0.$$
(2.9)

The following argument involving inspecting the coefficients rules out the possibility of (2.9). Indeed, in view of (2.7) we would have from (2.9) that

$$(1+c_s) + \sum_{r \in S \setminus \{s,t\}} c_r \alpha_r = (2\langle \alpha_s, \beta_t \rangle - c_t) \alpha_t.$$

Now if $2\langle \alpha_s, \beta_t \rangle - c_t > 0$ then we have a contradiction to (D2), since then $\alpha_t \in \text{PLC}(\Pi_1 \setminus \{\alpha_t\})$; whereas if $2\langle \alpha_s, \beta_t \rangle - c_t \leq 0$ then we again have a contradiction to (D2), since then $0 \in \text{PLC}(\Pi_1)$. Thus (2.8) must be the case, and in view of (2.7) we have

$$(1 - c_s)\alpha_s = (2\langle \alpha_s, \beta_t \rangle + c_t)\alpha_t + \sum_{r \in S \setminus \{s, t\}} c_r \alpha_r.$$

Suppose for a contradiction that $\langle \alpha_s, \beta_t \rangle > 0$. Then $2\langle \alpha_s, \beta_t \rangle + c_t > 0$. Now if $1 - c_s > 0$ then we have a contradiction to condition (D2), since $\alpha_s \in \text{PLC}(\Pi_1 \setminus \{\alpha_s\})$; whereas if $1 - c_s \leq 0$ then we again have a contradiction to (D2), since $0 \in \text{PLC}(\Pi_1)$. Thus it follows from these contradictions that $\langle \alpha_s, \beta_t \rangle \leq 0$, and interchange the roles of s and t, we see that $\langle \alpha_t, \beta_s \rangle \leq 0$, whence (D3) holds.

Next, suppose that further $\langle \alpha_s, \beta_t \rangle = 0$, and we prove that (D4) holds. Observe that

$$\rho_1(t)\rho_1(s)\alpha_t = \rho_1(t)(\rho_1(s)\alpha_t) = -\alpha_t - 2\langle \alpha_t, \beta_s \rangle \alpha_s + 4\langle \alpha_t, \beta_s \rangle \langle \alpha_s, \beta_t \rangle \alpha_t$$
$$= -\alpha_t - 2\langle \alpha_t, \beta_s \rangle \alpha_s.$$

Again the assumption that $\Phi_1 = \Phi_1^+ \uplus \Phi_1^-$ implies that either

$$-\alpha_t - 2\langle \alpha_t, \beta_s \rangle \alpha_s = \sum_{r \in S} c_r \alpha_r, \text{ where all } c_r \ge 0,$$
 (2.10)

or else

$$-\alpha_t - 2\langle \alpha_t, \beta_s \rangle \alpha_s = \sum_{r \in S} -c_r \alpha_r, \text{ where all } c_r \ge 0.$$
 (2.11)

A similar argument involving inspecting the coefficients together with (D2) make it possible to conclude that only (2.11) is possible. Hence

$$(-2\langle \alpha_t, \beta_s \rangle + c_s)\alpha_s + \sum_{r \in S \setminus \{s, t\}} c_r \alpha_s = (1 - c_t)\alpha_t.$$
 (2.12)

Observe that (D3) just prove above yields that $-\langle \alpha_t, \beta_s \rangle \geq 0$. Now if $1 - c_t < 0$ then we will have a contradiction to (D2), since then $0 \in \text{PLC}(\Pi_1)$; whereas if $1 - c_t > 0$ then we again have a contradiction to (D2), since then $\alpha_t \in \text{PLC}(\Pi_1 \setminus \{\alpha_s\})$. Thus $c_t = 1$, and then (2.12) implies, in view of (D2) and (D3), that $\langle \alpha_t, \beta_s \rangle = 0 = c_s$ (and $c_r = 0$ for all $r \in S \setminus \{s, t\}$). Interchange the roles of s and t we deduce that $\langle \alpha_t, \beta_s \rangle = 0$ implies that $\langle \alpha_s, \beta_t \rangle = 0$, whence (D4) holds.

To prove that (D5) holds, we may assume that $\langle \alpha_s, \beta_t \rangle \langle \alpha_t, \beta_s \rangle \neq 0$, for otherwise $\langle \alpha_s, \beta_t \rangle \langle \alpha_t, \beta_s \rangle = \cos^2 \frac{\pi}{2}$, trivially satisfying (D5). We let

 \mathcal{A} , γ , q, X, p_n , A and B be as defined before Proposition 2.4. If we set

$$\mathcal{A} = \mathbb{R};$$

$$q = 1;$$

$$\gamma = \sqrt{\langle \alpha_s, \beta_t \rangle \langle \alpha_t, \beta_s \rangle};$$

and

$$X = \frac{-\langle \alpha_t, \beta_s \rangle}{\sqrt{\langle \alpha_s, \beta_t \rangle \langle \alpha_t, \beta_s \rangle}},$$

then it is readily checked that A and B are the matrices representing the actions of $\rho_1(s)$ and $\rho_1(t)$ respectively, on the $\langle \{ \rho_1(s), \rho_1(t) \} \rangle$ -invariant subspace $\mathbb{R}\alpha_s + \mathbb{R}\alpha_t$. It follows from (2.1) to (2.4) and a similar argument involving inspecting the coefficients as used above that the requirement

$$\langle \{ \rho_1(s), \rho_1(t) \} \rangle \alpha_s \cup \langle \{ \rho_1(s), \rho_1(t) \} \rangle \alpha_t \subseteq \Phi_1^+ \uplus \Phi_1^-$$

is equivalent to $p_n p_{n+1} \ge 0$ for all $n \in \mathbb{N}$. By Proposition 2.4, this later condition is, in turn, equivalent to

$$\langle \alpha_s, \beta_t \rangle \langle \alpha_t, \beta_s \rangle \in \{ \cos^2 \frac{\pi}{m} \mid m \in \mathbb{N}_{\geq 2} \} \cup [1, \infty),$$

whence (D5) holds, finally establishing that (i) implies (iii).

Notation 2.5. For $w_i \in W_i$ (for each $i \in \{1, 2\}$), let $\operatorname{ord}(w_i)$ denote the order of w_i in W_i . For those $s, t \in S$ with $\langle \alpha_s, \beta_t \rangle \langle \alpha_t, \beta_s \rangle \geq 1$, extend the definition of m_{st} (given in Theorem 2.3) by setting $m_{st} = \infty$.

Proposition 2.6. Suppose that one of the (equivalent) statements of Theorem 2.3 is satisfied. Then $\operatorname{ord}(\rho_i(s)\rho_i(t)) = m_{st}$.

Proof. If one of the (equivalent) statements of Theorem 2.3 is satisfied, then $\mathscr{C} := (S, V_1, V_2, \Pi_1, \Pi_2, \langle, \rangle)$ forms a Coxeter datum in the sense of [14], and thus the required result follows immediately from Proposition 2.8 of [14].

We point out that a Coxeter datum in the sense of [14] automatically satisfies the conditions (D1) to (D5) of the present paper. Indeed, the only possible difference of these two formulations is that in (D2) of the present paper we require a seemingly extra condition that $\alpha_s \notin \text{PLC}(\Pi_1 \setminus \{\alpha_s\})$ and $\beta_s \notin \text{PLC}(\Pi_2 \setminus \{\beta_s\})$ for each $s \in S$. However, it can be checked that this condition is an immediate consequence of (C1), (C2) and (C5) of a Coxeter datum in the sense of [14] (in fact, this is just [14, Lemma 2.5]). Thus we have:

Proposition 2.7. The following are equivalent:

(i) $\mathscr{C} := (S, V_1, V_2, \Pi_1, \Pi_2, \langle , \rangle)$ satisfies one of the (equivalent) statements of Theorem 2.3;

(ii) $\mathscr{C} := (S, V_1, V_2, \Pi_1, \Pi_2, \langle , \rangle)$ is a Coxeter datum in the sense of [14].

Next we have a result which enables us to give a characterization of Coxeter groups, among a large family of linear groups that are generated by involutions, in terms of their root systems:

Theorem 2.8. Let S, Π_1 and Π_2 be the same as at the beginning of this section, and let R_1 , W_1 , Φ_1 , R_2 , W_2 and Φ_2 be as in Definition 2.1. Let (W, R) be a Coxeter system in the sense of [2] or [21], with W being an abstract group generated by a set of involutions $R := \{r_s \mid s \in S\}$ subject only to the condition that for $s, t \in S$ the order of $r_s r_t$ is either equal to m if $\langle \alpha_s, \beta_t \rangle \langle \alpha_t, \beta_s \rangle = \cos^2(\pi/m)$, or else equal to infinity. Then $\Phi_1 = \Phi_1^+ \uplus \Phi_1^-$, or equivalently, $\Phi_2 = \Phi_2^+ \uplus \Phi_2^-$ only if there exist isomorphisms $f_1 \colon W \to W_1$ and $f_2 \colon W \to W_2$ such that $f_1(r_s) = \rho_1(s)$ and $f_2(r_s) = \rho_2(s)$ for all $s \in S$.

Proof. Follows immediately from Proposition 2.7 above and [14, Theorem 2.10].

Remark 2.9. Theorem 2.8 shows that if $\Phi_1 = \Phi_1^+ \uplus \Phi_1^-$, or equivalently, $\Phi_2 = \Phi_2^+ \uplus \Phi_2^-$ then (W_1, R_1) and (W_2, R_2) are Coxeter systems isomorphic to (W, R). It is well known in the literature that all Coxeter groups have root systems decomposable into a disjoint union of positive roots and negative roots ([1, Proposition 4.2.5] or [21, §5.4], for example). Furthermore, given an arbitrary Coxeter system (W', R'), it follows from [12] and [14] that we could associate a Coxeter datum $\mathscr{C}' := (S', V'_1, V'_2, \Pi'_1, \Pi'_2, \langle , \rangle')$ to (W', R'), such that the paired root systems Φ'_1 and Φ'_2 arising from this particular Coxeter datum admit decompositions $\Phi'_1 = \Phi'_1^+ \uplus \Phi'_1^-$ and $\Phi'_2 = \Phi'_2^+ \uplus \Phi'_2^-$. These facts combined with Theorem 2.8 yield that if a linear group is generated by involutions, then it is a Coxeter group if and only if it has a root system decomposable into a disjoint union of positive roots and negative roots.

Let W and R be as in Theorem 2.8, we call (W,R) the abstract Coxeter system corresponding to $\mathscr C$ with W being the corresponding abstract Coxeter group. We see immediately from the above theorem that f_1 and f_2 give rise to faithful W-actions on V_1 and V_2 in the natural way with $wx := (f_1(w))(x)$ and $wy := (f_2(w))(y)$ for all $w \in W$, $x \in V_1$ and $y \in V_2$.

To close this section we include the following useful result taken from [14]:

Lemma 2.10. (i) \langle , \rangle is W-invariant, that is, $\langle wx, wy \rangle = \langle x, y \rangle$ for all $w \in W$, $x \in V_1$ and $y \in V_2$.

(ii) There exists a W-equivariant bijection $\phi \colon \Phi_1 \to \Phi_2$ satisfying $\phi(\alpha_s) = \beta_s$ for all $s \in S$.

(iii) Let ϕ be as in (ii) above, and let $x, x' \in \Phi_1$. Then $\langle x, \phi(x') \rangle = 0$ if and only if $\langle x', \phi(x) \rangle = 0$.

Proof. (i): Lemma 2.13 of [14].

(ii): Proposition 3.18 of [14].

(iii): Corollary 3.25 of [14].
$$\square$$

For the rest of this paper, the notation ϕ will be fixed for the W-equivariant bijection in Lemma 2.10 (iii).

3. Reflection Subgroups and Canonical Generators in Coxeter Groups

Given a Coxeter group W and its Coxeter generators R, a subgroup W' of W is called a reflection subgroup if W' is generated by those elements of the form wrw^{-1} (where $w \in W$ and $r \in R$). It is well known that W' is a Coxeter group, and consequently the notion of a Coxeter datum as in the previous section applies to W'. In this section we study the paired root systems for W' as a subsets of the paired root systems for W. In the spirit of the previous section, our investigation of the paired root systems for W' is based on a Coxeter datum \mathscr{C}' closely related to the Coxeter datum for the over group W. In particular, we show that the Coxeter generators of W' are characterize by this Coxeter datum \mathscr{C}' . In addition to obtaining certain geometric insights of reflection subgroups of Coxeter groups, these investigations also establish the fact that the method of constructing paired root systems via Coxeter data applies to paired root systems for reflection subgroups of a Coxeter group, either on their own or as subsets of the paired root systems of the over group.

Suppose that $\mathscr{C} := (S, V_1, V_2, \Pi_1, \Pi_2, \langle , \rangle)$ satisfies conditions (D1) to (D5) of Section 2 inclusive (or in view of Proposition 2.7, we could equivalently suppose that \mathscr{C} is a Coxeter datum in the sense of [14]). Let (W, R) be the abstract Coxeter system associated to the Coxeter datum \mathscr{C} , and keep all the notation of the previous section.

Let $T := \bigcup_{w \in W} wRw^{-1}$, and call it the set of reflections in W. For $s \in S$ and $w \in W$, observe that for each $x \in V_1$ and $y \in V_2$, Lemma 2.10 yields that

$$wr_s w^{-1} x = w(w^{-1} x - 2\langle w^{-1} x, \beta_s \rangle \alpha_s) = x - 2\langle w^{-1} x, \beta_s \rangle w \alpha_s$$
$$= x - 2\langle x, \phi(w \alpha_s) \rangle w \alpha_s, \quad (3.1)$$

and

$$wr_s w^{-1} y = w(w^{-1}y - 2\langle \alpha_s, w^{-1}y \rangle \beta_s) = y - 2\langle w\alpha_s, y \rangle w\beta_s$$
$$= y - 2\langle \phi^{-1}(w\beta_s), y \rangle w\beta_s. \quad (3.2)$$

Now suppose that $\alpha \in \Phi_1$ and $\beta \in \Phi_2$ are arbitrary. Then $\alpha = w_1 \alpha_s$ and $\beta = w_2 \beta_t$ for some $w_1, w_2 \in W$ and $s, t \in S$. It follows from (3.1)

and (3.2) that we can unambiguously define $r_{\alpha}, r_{\beta} \in T$, the reflection corresponding to α and β respectively, by

$$r_{\alpha} = r_{w_1 \alpha_s} := w_1 r_s w_1^{-1}, \tag{3.3}$$

and

$$r_{\beta} = r_{w_2\beta_t} := w_2 r_t w_2^{-1}, \tag{3.4}$$

with

$$r_{\alpha}x = x - 2\langle x, \phi(\alpha) \rangle \alpha$$

for all $x \in V_1$ and

$$r_{\beta}y = y - 2\langle \phi^{-1}(\beta), y \rangle \beta$$

for all $y \in V_2$.

Definition 3.1. (i) A subgroup W' of W is called a *reflection subgroup* if $W' = \langle W' \cap T \rangle$.

- (ii) For each $i \in \{1, 2\}$, a subset Φ'_i of Φ_i is called a *root subsystem* if $r_x y \in \Phi'_i$ whenever $x, y \in \Phi'_i$.
- (iii) If W' is a reflection subgroup, set $\Phi_i(W') := \{ x \in \Phi_i \mid r_x \in W' \}$ for each $i \in 1, 2$.

Lemma 3.2. Let W' be a reflection subgroup of W. Then for each $i \in \{1, 2\}$

$$W'\Phi_i(W') = \Phi_i(W').$$

Proof. We prove that $W'\Phi_1(W') = \Phi_1(W')$ here, and we stress that the other half follows in the same way. Let $w \in W'$. By definition, we have $w = t_1 t_2 \cdots t_n$ where $t_1, t_2, \ldots, t_n \in W' \cap T$. Now let $x \in \Phi_1(W')$ be arbitrary. Then $r_x \in W'$, and hence $r_{t_n x} = t_n r_x t_n \in W'$, which in turn yields that $t_n x \in \Phi_1(W')$. Then it follows that $t_{n-1} t_n x \in \Phi_1(W')$ and so on. Thus $wx = t_1 \cdots t_n x \in \Phi_1(W')$. Since $x \in \Phi_1(W')$ is arbitrary, it follows that $w\Phi_1(W') \subseteq \Phi_1(W')$. Finally, replacing $w \in W'$ by w^{-1} we see that $\Phi_1(W') \subseteq w\Phi_1(W')$.

Remark 3.3. Let W' be a reflection subgroup. For each $i \in \{1, 2\}$, it follows from the above lemma that $\Phi_i(W')$ is a root subsystem of Φ_i , and we call it the root subsystem corresponding to W'. It is easily seen that there is a bijective correspondence between the set of reflection subgroups W' of W and the set of root subsystems Φ'_i of Φ_i : W' uniquely determines the corresponding root subsystem $\Phi_i(W')$; and Φ'_i uniquely determines the reflection subgroup $W' := \langle \{r_x \mid x \in \Phi'_i \} \rangle$.

In fact, for a reflection subgroup W', we shall see that $\Phi_1(W')$ and $\Phi_2(W')$ are the root systems for the Coxeter group W' arising from a suitably chosen Coxeter datum. In order to do this, we need a few preparatory results first.

Remark 3.4. For each $i \in \{1,2\}$, it has been observed in [14] that non-trivial scalar multiple of an element of Φ_i can still be an element

of Φ_i (see the example immediately after [14, Definition 3.1] and [14, Lemma 3.20]). Therefore, unlike in the classical setting of [21], we do not have a bijection from T to either Φ_1^+ or Φ_2^+ .

Definition 3.5. For each $i \in \{1, 2\}$, define an equivalence relation \sim_i on Φ_i as follows: if $z_1, z_2 \in \Phi_i$, then $z_1 \sim_i z_2$ if and only if z_1 and z_2 are (non-zero) scalar multiples of each other. For each $z \in \Phi_i$, write \widehat{z} for the equivalence class containing z and write $\widehat{\Phi}_i = \{\widehat{z} \mid z \in \Phi_i\}$.

Remark 3.6. Observe that W has a natural action on $\widehat{\Phi}_i$ (for each $i \in \{1,2\}$) given by $w\widehat{z} = \widehat{wz}$ for all $w \in W$ and $z \in \Phi_i$. Furthermore, given $z, z' \in \Phi_i$, the corresponding reflections r_z and $r_{z'}$ are equal if and only if $\widehat{z} = \widehat{z'}$.

Definition 3.7. For $i \in \{1, 2\}$, and for each $w \in W$, define

$$N_i(w) = \{ \widehat{z} \mid z \in \Phi_i^+ \text{ and } wz \in \Phi_i^- \}.$$

Note that for $w \in W$, the set $N_i(w)$ (i = 1, 2) can be alternatively characterized as $\{\hat{z} \mid z \in \Phi_i^- \text{ and } wz \in \Phi_i^+\}$. Hence $\hat{z} \in N_i(w)$ if and only if precisely one element of the set $\{z, wz\}$ is in Φ_i^+ .

Notation 3.8. Let $\ell: W \to \mathbb{N}$ denote the *length function* with respect to (W, R), that is, for $w \in W$,

$$\ell(w) = \min\{ n \in \mathbb{N} \mid w = r_1 r_2 \cdots r_n, \text{ where } r_1, r_2, \cdots, r_n \in R \}.$$

A mild generalization of the techniques used in ([21, §5.6]) then yields the following connection between the length function and the functions N_1 and N_2 :

Lemma 3.9. ([14, Lemma 3.8]) (i) $N_1(r_s) = \{\widehat{\alpha}_s\}$ and $N_2(r_s) = \{\widehat{\beta}_s\}$ for all $s \in S$.

- (ii) Let $w \in W$. Then $N_1(w)$ and $N_2(w)$ both have cardinality $\ell(w)$.
- (iii) Let $w_1, w_2 \in W$ and let \dotplus denote set symmetric difference. Then $N_i(w_1w_2) = w_2^{-1}N_i(w_1) \dotplus N_i(w_2)$ for each $i \in \{1, 2\}$.

The last lemma enables us to deduce the following generalization of [14, Lemma 3.2 (ii)]:

Proposition 3.10. For each $i \in \{1, 2\}$, let $w \in W$ and $x \in \Phi_i^+$. If $\ell(wr_x) > \ell(w)$ then $wx \in \Phi_i^+$, whereas if $\ell(wr_x) < \ell(w)$ then $wx \in \Phi_i^-$.

Proof. We prove the statement that $\ell(wr_x) > \ell(w)$ if and only if wx is positive in the case $x \in \Phi_1$, and again we stress that a similar argument also shows the desired result holds in Φ_2 .

Observe that the second statement follows from the first, applied to wr_x in place of w: indeed if $\ell(wr_x) < \ell(w)$ then $\ell((wr_x)r_x) > \ell(wr_x)$, forcing $(wr_x)x = w(r_xx) = -wx \in \Phi_+^+$, that is, $wx \in \Phi_-^-$.

Now we prove the first statement in Φ_1 . Proceed by induction on $\ell(w)$, the case $\ell(w) = 0$ being trivial. If $\ell(w) > 0$, then there exists

 $s \in S$ with $\ell(r_s w) = \ell(w) - 1$, and hence

$$\ell((r_s w)r_x) = \ell(r_s(wr_x)) \ge \ell(wr_x) - 1 > \ell(w) - 1 = \ell(r_s w).$$

Then the inductive hypothesis yields that $(r_s w)x \in \Phi_1^+$. Suppose for a contradiction that $wx \in \Phi_1^-$. Then $\widehat{wx} \in N_1(r_s)$ and Lemma 3.9 (i) yields that $wx = -\lambda \alpha_s$ for some $\lambda > 0$. But then $r_s wx = \lambda \alpha_s$, and hence $(r_s w)r_x(r_s w)^{-1} = r_s$ by calculations similar to (3.3) and (3.4). But this yields that $wr_x = r_s w$, contradicting $\ell(wr_x) > \ell(w) > \ell(r_s w)$, as desired.

Definition 3.11. For each $w \in W$, define

$$\overline{N}(w) := \{ t \in T \mid \ell(wt) < \ell(w) \}.$$

If $t \in T$ then $t = wr_s w^{-1}$ for some $w \in W$ and $s \in S$, and hence it follows from calculations like (3.3) and (3.4) that $t = r_{w\alpha_s} = r_{w\beta_s}$. This combined with Proposition 3.10 give us:

Proposition 3.12. Let $w \in W$. Then

$$\overline{N}(w) = \{ r_x \mid \widehat{x} \in N_i(w) \}$$

for each $i \in \{1, 2\}$.

Definition 3.13. Suppose that W' is a reflection subgroup. Then we define

$$S(W') := \{ t \in T \mid \overline{N}(t) \cap W' = \{t\} \}$$

and

$$\Delta_i(W') := \{ x \in \Phi_i^+ \mid r_x \in S(W') \}$$

for each $i \in \{1, 2\}$.

For a reflection subgroup W', the set S(W') is called the *canonical* generators of W' in [10], and it is well known that (W', S(W')) is a Coxeter system. Indeed, we have:

Lemma 3.14. [10] Let W' be a reflection subgroup of W.

- (i) If $t \in W' \cap T$, then there exist $m \in \mathbb{N}$ and $t_0, \dots, t_m \in S(W')$ such that $t = t_m \cdots t_1 t_0 t_1 \cdots t_m$.
- (ii) (W', S(W')) is a Coxeter system.

Proof. (i):
$$[10, Lemma (1.7) (ii)].$$
 (ii)]: $[10, Theorem (1.8) (i)].$

For a reflection subgroup W', we will show that $\Delta_1(W')$ and $\Delta_2(W')$ can be characterized in terms of a suitably defined Coxeter datum. Before we could prove this, we need a number of simple observations.

Observe that for a reflection subgroup W' we can equivalently define $\Delta_i(W')$ by requiring

$$\Delta_i(W') := \{ x \in \Phi_i^+ \mid N_i(r_x) \cap \widehat{\Phi_i(W')} = \{ \widehat{x} \} \}.$$
 (3.5)

Suppose that $\Delta'_1 \subseteq \Phi_1^+$ and $\Delta'_2 \subseteq \Phi_2^+$ are two sets of roots with $\phi(\Delta'_1) = \Delta'_2$ (where ϕ is as in Lemma 2.10 (iii)). Furthermore, suppose that Δ'_1 and Δ'_2 satisfy the following:

- (i') $\langle x, \phi(x') \rangle \leq 0$, for all distinct $x, x' \in \Delta'_1$;
- (ii') $\langle x, \phi(x') \rangle \langle x', \phi(x) \rangle \in \{ \cos^2(\pi/m) \mid m \in \mathbb{N}, m \geq 2 \} \cup [1, \infty), \text{ for all } x, x' \in \Delta'_1 \text{ with } r_x \neq r_{x'}.$

It follows from Lemma 2.10 that

$$\langle x, \phi(x) \rangle = 1$$
, for all $x \in \Delta'_1$. (3.6)

Since $\Delta'_1 \subseteq PLC(\Pi_1)$ and $\Delta'_2 \subseteq PLC(\Pi_2)$, it follows that $0 \notin PLC(\Delta'_1)$ and $0 \notin PLC(\Delta'_2)$. Also it can be readily checked from (i'), (ii') and (3.6) that $x \notin PLC(\Delta'_1 \setminus \{x\})$ and $\phi(x) \notin PLC(\Delta'_2 \setminus \{\phi(x)\})$ for all $x \in \Delta'_1$. Furthermore, Lemma 2.10 (iii) ensures that $\langle x, \phi(x') \rangle = 0$ whenever $\langle x', \phi(x) \rangle = 0$ for all $x, x' \in \Delta'_1 \subseteq \Phi_1$, Thus Δ'_1 and Δ'_2 satisfy conditions (D1) to (D5) of the present paper inclusive. If we let S' be an indexing set for both Δ'_1 and Δ'_2 then

$$\mathscr{C}' := (S', \operatorname{span}(\Delta_1'), \operatorname{span}(\Delta_2'), \Delta_1', \Delta_2', \langle , \rangle'),$$

(where \langle , \rangle' denotes the restriction of \langle , \rangle to span $(\Delta_1') \times$ span (Δ_2')) constitutes a Coxeter datum in the sense of [14]. Now if we let $R' := \{ r_x \mid x \in \Delta_1' \} (= \{ r_y \mid y \in \Delta_2' \})$, and set $W' = \langle R' \rangle$, then it is clear that W' is a reflection subgroup of W. Furthermore, it follows from Theorem 2.8 that (W', R') forms a Coxeter system. Then upon applying Lemma 3.9 and (3.5) to \mathscr{C}' and W' we may conclude that S(W') = R' and consequently $\widehat{\Delta_1(W')} = \widehat{\Delta_1'}$ and $\widehat{\Delta_2(W')} = \widehat{\Delta_2'}$. Summing up, we have:

Proposition 3.15. Suppose that $\Delta'_1 \subseteq \Phi_1^+$ and $\Delta'_2 \subseteq \Phi_2^+$ such that

- (A1) $\phi(\Delta_1') = \Delta_2';$
- (A2) $\langle x, \phi(x') \rangle \leq 0$, for all distinct $x, x' \in \Delta'_1$;
- (A3) $\langle x, \phi(x') \rangle \langle x', \phi(x) \rangle \in \{ \cos^2(\pi/m) \mid m \in \mathbb{N}, m \geq 2 \} \cup [1, \infty), \text{ for all } x, x' \in \Delta'_1 \text{ with } r_x \neq r_{x'}.$

Then
$$W' = \langle \{ r_x \mid x \in \Delta'_1 \} \rangle$$
 is a reflection subgroup of W with $\widehat{\Delta'_1} = \widehat{\Delta_1(W')}$ and $\widehat{\Delta'_2} = \widehat{\Delta_2(W')}$.

It turns out that the converse of Proposition 3.15 is also true, namely: if W' is a reflection subgroup of W, and if $x, x' \in \Delta_1(W')$ with $r_x \neq r_{x'}$, then conditions (A2) and (A3) of Proposition 3.15 must be satisfied. Since Lemma 2.10 (iii) ensures that $\langle x, \phi(x') \rangle = 0$ if and only if $\langle x'\phi(x) \rangle = 0$, it follows from this assertion and a quick argument similar to the one used immediately after (3.6) that representative elements from $\Delta_1(W')$ and $\Delta_2(W')$ can be used to form a Coxeter datum for W'. Hence this assertion and Proposition 3.15 together yield that for a reflection subgroup W', the corresponding $\Delta_i(W')$ (i = 1, 2) can be characterized by a suitable Coxeter datum. We devote the rest of this section to a proof of this assertion.

Lemma 3.16. Let W' be a reflection subgroup of W.

(i) For each $i \in \{1, 2\}$, let $x \in \Pi_i \setminus \Phi_i(W')$. Then $\Delta_i(r_x W' r_x) = r_x \Delta_i(W')$.

(ii) For each $i \in \{1, 2\}$, $\Phi_i(W') = W' \Delta_i(W')$.

Proof. (i): It is readily checked that $r\Phi_i(W') = \Phi_i(rW'r)$ for all $r \in T$. Since $x \in \Pi_i \setminus \Phi_i(W')$, it follows that $r_x \in R \setminus W'$. Let $y \in \Delta_i(W')$ be arbitrary. Then

$$N_{i}(r_{(rxy)}) \cap \widehat{\Phi_{i}(r_{x}W'r_{x})} = N_{i}(r_{x}r_{y}r_{x}) \cap \widehat{\Phi_{i}(r_{x}W'r_{x})}$$

$$(by (3.3) \text{ and } (3.4))$$

$$= (r_{x}N_{i}(r_{x}r_{y}) \dotplus N_{i}(r_{x})) \cap \widehat{\Phi_{i}(r_{x}W'r_{x})}$$

$$(by \text{ Lemma } 3.9 \text{ (iii)})$$

$$= (r_{x}r_{y}N_{i}(r_{x}) \dotplus r_{x}N_{i}(r_{y}) \dotplus N_{i}(r_{x}))$$

$$\cap \widehat{\Phi_{i}(r_{x}W'r_{x})}$$

$$(again by \text{ Lemma } 3.9 \text{ (iii)})$$

$$= r_{x}((r_{y}N_{i}(r_{x}) \dotplus N_{i}(r_{y}) \dotplus N_{i}(r_{x}))$$

$$\cap \widehat{\Phi_{i}(W')})$$

$$= r_{x}((r_{y}\{\widehat{x}\} \dotplus N_{i}(r_{y}) \dotplus \{\widehat{x}\}) \cap \widehat{\Phi_{i}(W')})$$

$$(by \text{ Lemma } 3.9 \text{ (i)})$$

$$= r_{x}(N_{i}(r_{y}) \cap \widehat{\Phi_{i}(W')})$$

$$(since \widehat{(x)}, r_{y}\widehat{x} \notin \widehat{\Phi_{i}(W')})$$

$$= \{\widehat{r_{x}y}\}$$

$$(since y \in \Delta_{i}(W')).$$

Hence $r_x y \in \Delta_i(r_x W' r_x)$. This proves that $r_x \Delta_i(W') \subseteq \Delta_i(r_x W' r_x)$. But $x \in \Pi_i \backslash r_x \Phi_i(W')$, so the above yields that $r_x \Delta_i(r_x W' r_x) \subseteq \Delta_i(W')$ proving the desired result.

(ii): Since $\Delta_i(W') \subseteq \Phi_i(W')$ for each $i \in \{1, 2\}$, it follows from Lemma 3.2 that $W'\Delta_i(W') \subseteq \Phi_i(W')$.

Conversely if $x \in \Phi_i(W')$ then $r_x \in W' \cap T$. By Lemma 3.14 (i), there are $x_0, x_1, \dots, x_m \in \Delta_i(W')$ $(m \in \mathbb{N})$ such that

$$r_x = r_{x_m} \cdots r_{x_1} r_{x_0} r_{x_1} \cdots r_{x_m}.$$

Calculations similar to those of (3.3) and (3.4) enable us to conclude that $\lambda x = r_{x_m} \cdots r_{x_1} x_0 \in W' \Phi_i(W')$ for some (nonzero) scalar λ . Now since $\frac{1}{\lambda} x_0 = (r_{x_m} \cdots r_{x_1})^{-1} x \in \Phi_i$, it follows that $\frac{1}{\lambda} x_0 \in \Delta_i(W')$ and hence $x = r_{x_m} \cdots r_{x_1}(\frac{1}{\lambda} x_0) \in W' \Delta_i(W')$ as required.

Definition 3.17. Let W' be a reflection subgroup of W, and let $\ell_{W'}$: $W' \to \mathbb{N}$ be the length function on (W', S(W')) defined by

$$\ell_{W'}(w) = \min\{ n \in \mathbb{N} \mid w = r_1 \cdots r_n, \text{ where } r_i \in S(W') \}.$$

If $w = r_1 \cdots r_n \in W'$ $(r_i \in S(W'))$ and $n = \ell_{W'}(w)$ then $r_1 \cdots r_n$ is called a reduced expression for w (with respect to S(W')).

Lemma 3.18. Let W' be a reflection subgroup. For each $i \in \{1, 2\}$,

- (i) $N_i(r_x) \cap \widehat{\Phi_i(W')} = \{\widehat{x}\} \text{ for all } x \in \Delta_i(W');$
- (ii) for all $w_1 \in W$ and $w_2 \in W'$

$$N_i(w_1w_2) \cap \widehat{\Phi_i(W')} = w_2^{-1}(N_i(w_1) \cap \widehat{\Phi_i(W')}) \dotplus (N_i(w_2) \cap \widehat{\Phi_i(W')}).$$

Proof. (i) is just the definition of $\Delta_i(W')$.

(ii) Lemma 3.9 (iii) yields that $N_i(w_1w_2) = w_2^{-1}N_i(w_1) + N_i(w_2)$, and hence

$$N_i(w_1w_2) \cap \widehat{\Phi_i(W')} = (w_2^{-1}N_i(w_1) \cap \widehat{\Phi_i(W')}) \dotplus (N_i(w_2) \cap \widehat{\Phi_i(W')}).$$

Since $w_2 \in W'$ it follows from Lemma 3.2 that $w_2^{-1}\widehat{\Phi_i(W')} = \widehat{\Phi_i(W')}$. Thus $w_2^{-1}N_i(w_1) \cap \widehat{\Phi_i(W')} = w_2^{-1}(N_i(w_1) \cap \widehat{\Phi_i(W')})$, giving us

$$N_i(w_1w_2) \cap \widehat{\Phi_i(W')} = w_2^{-1}(N_i(w_1) \cap \widehat{\Phi_i(W')}) \dotplus (N_i(w_2) \cap \widehat{\Phi_i(W')}).$$

Lemma 3.19. Let W' be a reflection subgroup. For each $i \in \{1, 2\}$ and all $w \in W'$, we have

(i) $|N_i(w) \cap \widehat{\Phi_i(W')}| = \ell_{W'}(w)$. Furthermore, if $w = r_{x_1} \cdots r_{x_n}$ (where $x_1, \dots, x_n \in \Delta_i(W')$) is reduced with respect to (W', S(W')) then

$$N_i(w) \cap \widehat{\Phi_i(W')} = \{\widehat{y_1}, \cdots \widehat{y_n}\}$$

where $y_j = (r_{x_n} \cdots r_{x_{j+1}}) x_j$ for all $j = 1, \dots, n$.

(ii)
$$N_i(w) \cap \widehat{\Phi_i(W')} = \{\widehat{x} \in \widehat{\Phi_i(W')} \mid \ell_{W'}(wr_x) < \ell_{W'}(w)\}.$$

Proof. (i): For each $j \in \{1, \dots, n\}$, set $t_j = r_{x_n} \cdots r_{x_{j+1}} r_{x_j} r_{x_{j+1}} \cdots r_{x_n}$, that is, $t_j = r_{y_j}$. If $t_j = t_k$ where j > k then

$$w = r_{x_1} \cdots r_{x_{k-1}} r_{x_{k+1}} \cdots r_{x_n} t_k$$

$$= r_{x_1} \cdots r_{x_{k-1}} r_{x_{k+1}} \cdots r_{x_n} t_j$$

$$= r_{x_1} \cdots r_{x_{k-1}} r_{x_{k+1}} \cdots r_{x_{j-1}} r_{x_{j+1}} \cdots r_{x_n}$$

contradicting $\ell_{W'}(w) = n$. Hence the t_j 's are all distinct and consequently all the \hat{y}_j 's are all distinct. Now by repeated application of

Lemma 3.18 (ii), for each $i \in \{1, 2\}$ we have

$$N_{i}(w) \cap \widehat{\Phi_{i}(W')}$$

$$= (N_{i}(r_{x_{n}} \cap \widehat{\Phi_{i}(W')}) \dotplus r_{x_{n}}(N_{i}(r_{n-1}) \cap \widehat{\Phi_{i}(W')}) \dotplus \cdots$$

$$\dotplus r_{x_{n}} \cdots r_{x_{2}}(N_{i}(r_{x_{1}}) \cap \widehat{\Phi_{i}(W')})$$

$$= \{\widehat{y_{n}}\} \dotplus \{\widehat{y_{n-1}}\} \dotplus \cdots \dotplus \{\widehat{y_{1}}\}$$

$$= \{\widehat{y_{1}}, \cdots, \widehat{y_{n}}\}$$

and consequently $|N_i(w) \cap \widehat{\Phi_i(W')}| = \ell_{W'}(w)$.

(ii): Let $w = r_{x_1} \cdots r_{x_n}$ be a reduced expression for $w \in W'$ with respect to S(W') $(x_1, \dots, x_n \in \Delta_i(W'))$. Then for each $i \in \{1, 2\}$, Part (i) above yields that

$$N_i(w) \cap \widehat{\Phi_i(W')} = \{ \widehat{y_1}, \cdots, \widehat{y_n} \}$$

where $y_j = (r_{x_n} \cdots r_{x_{j+1}}) x_j$, for all $j \in \{1, \cdots, n\}$. Now for each such j,

$$wr_{y_j} = wr_{x_n} \cdots r_{x_{j+1}} r_{x_j} r_{x_{j+1}} \cdots r_{x_n} = r_{x_1} \cdots r_{x_{j-1}} r_{x_{j+1}} \cdots r_{x_n}$$

and so $\ell_{W'}(wr_{y_j}) \leq n-1 < \ell_{W'}(w)$. Hence if $\widehat{x} \in N_i(w) \cap \widehat{\Phi_i(W')}$, then $\ell_{W'}(wr_x) < \ell_{W'}(w)$.

Conversely, suppose that $x \in \Phi_i(W') \cap \Phi_i^+$ and $\widehat{x} \notin N_i(w)$. We are done if we could show that then $\ell_{W'}(wr_x) > \ell_{W'}(w)$. Observe that the given choice of x implies that $\widehat{x} \in N_i(r_x) \cap \widehat{\Phi_i(W')}$, furthermore, $\widehat{x} \notin r_x(N_i(w) \cap \widehat{\Phi_i(W')})$. Therefore

$$\widehat{x} \in r_x(N_i(w) \cap \widehat{\Phi_i(W')}) \dotplus (N_i(r_x) \cap \widehat{\Phi_i(W')}) = N_i(wr_x) \cap \widehat{\Phi_i(W')},$$

and by what has just been proved, this implies that

$$\ell_{W'}(w) = \ell_{W'}((wr_x)r_x) < \ell_{W'}(wr_x),$$

as desired.

The following is a mild generalization of [10, Lemma 3.2]:

Lemma 3.20. Let W' be a reflection subgroup. For each $i \in \{1, 2\}$, let $x, y \in \Delta_i(W')$ such that $r_x \neq r_y$. Let $n = \operatorname{ord}(r_x r_y)$. Then for $0 \leq m < n$

$$\underbrace{\cdots r_y r_x r_y}_{m \ factors} x \in \Phi_i^+ \qquad and \qquad \underbrace{\cdots r_x r_y r_x}_{m \ factors} y \in \Phi_i^+.$$

Proof. It is easily checked that when $0 \le m < n$ we have

$$\ell_{W'}(\underbrace{(\cdots r_y r_x r_y)}_{m \text{ factors}})r_x) = m+1 > m = \ell_{W'}\underbrace{(\cdots r_y r_x r_y)}_{m \text{ factors}},$$

as well as

$$\ell_{W'}(\underbrace{(\cdots r_x r_y r_x)}_{m \text{ factors}})r_y) = m+1 > m = \ell_{W'}\underbrace{(\cdots r_x r_y r_x)}_{m \text{ factors}}.$$

Hence the desired result follows immediately from Lemma 3.19.

In fact we can refine Lemma 3.20 with the following generalization of [10, Lemma 3.3]:

Lemma 3.21. Let W' be a reflection subgroup. For each $i \in \{1, 2\}$, let $x, y \in \Delta_i(W')$ with $r_x \neq r_y$. Let $n = \operatorname{ord}(r_x r_y)$, and let c_m, d_m, c'_m and d'_m be constants such that

$$\underbrace{(\cdots r_y r_x r_y)}_{m \ factors} x = c_m x + d_m y \quad and \quad \underbrace{(\cdots r_x r_y r_x)}_{m \ factors} y = c'_m x + d'_m y.$$

Then $c_m \ge 0$, $d_m \ge 0$, $c'_m \ge 0$ and $d'_m \ge 0$ whenever m < n.

Proof. By symmetry, it will suffice to prove that $d_m \geq 0$ and $d'_m \geq 0$. The proof of this will be based on an induction on $\ell(r_x)$.

Suppose first that $\ell(r_x) = 1$. Then $\lambda x \in \Pi_i$ for some $\lambda > 0$. Write $y = \sum_{z \in \Pi_i} \lambda_z z$ where $\lambda_z \geq 0$ for all $z \in \Pi_i$. In fact, $\lambda_{z_0} > 0$ for some

 $z_0 \in \Pi_i \setminus \{x\}$, since otherwise we would have $y \in \mathbb{R}x$ and so $r_x = r_y$. Now for $0 \le m < n$, Lemma 3.20 yields that

$$(\underbrace{\cdots r_y r_x r_y}_{m \text{ factors}}) x = c_m x + \sum_{z \in \Pi_i} d_m \lambda_z z \in \Phi_i^+.$$

That is

$$c_m x + d_m (\sum_{z \in \Pi_i} \lambda_z z) = \sum_{z \in \Pi_i} \mu_z z$$
, where $\mu_z \ge 0$, for all $z \in \Pi_i$.

Now if $d_m \leq 0$ then the above yields that

$$(c_m - \mu_x)x = (\mu_{z_0} - d_m \lambda_{z_0})z_0 + \sum_{z \in \Pi_i \setminus \{x, z_0\}} (\mu_z - d_m \lambda_z)z,$$

contradicting condition (D2). Therefore, $d_m > 0$ as required. Similarly $d'_m \ge 0$.

Suppose inductively now that the result is true for reflection subgroups W'' of W and x', $y' \in \Delta_i(W'')$ with $r_{x'} \neq r_{y'}$ and $\ell(r_{x'}) < \ell(r_x)$ where $\ell(r_x) \geq 3$. It is well know that there exists $z \in \Pi_i$ such that $\ell(r_z r_x r_z) = \ell(r_x) - 2$. Then $\ell(r_x r_z) < \ell(r_x)$, and thus $\widehat{z} \in N_i(r_x)$. But since $x \in \Delta_i(W')$ and $x \neq z$ (since $\ell(r_x) \geq 3$), it follows that $r_z \notin W'$. Let $W'' = r_z W' r_z$. Lemma 3.16 (i) yields that $\Delta_i(W'') = r_z \Delta_i(W')$ and therefore $r_z x$, $r_z y \in \Delta_i(W'')$. Now

$$r_{(r_z x)} = r_z r_x r_z$$
 and $r_{(r_z y)} = r_z r_y r_z$ (3.7)

and hence $\operatorname{ord}(r_{(r_z x)} r_{(r_z y)}) = \operatorname{ord}(r_x r_y) = n$. Since $\ell(r_{(r_z x)}) = \ell(r_x) - 2$, the inductive hypothesis gives

$$\underbrace{(\cdots r_{(r_z y)} r_{(r_z x)} r_{(r_z y)})}_{m \text{ factors}})(r_z x) = c_m(r_z x) + d_m(r_z y)$$

and

$$\underbrace{(\underbrace{\cdots r_{(r_z x)} r_{(r_z y)} r_{(r_z x)}}_{m \text{ factors}})(r_z y)} = c'_m(r_z x) + d'_m(r_z y)$$

where d_m , $d'_m \ge 0$ for $0 \le m < n$. Finally, by (3.7), the desired result follows on applying r_z to both sides of the last two equations.

Proposition 3.22. Let W' be a reflection subgroup of W. Suppose that $x, y \in \Delta_1(W')$ with $r_x \neq r_y$. Let $n = \operatorname{ord}(r_x r_y) \in \{\infty\} \cup \mathbb{N}$. Then

$$\langle x, \phi(y) \rangle \le 0$$

and

$$\begin{cases} \langle x, \phi(y) \rangle \langle y, \phi(x) \rangle = \cos^2 \frac{\pi}{n} & (n \in \mathbb{N}, n \ge 2) \\ \langle x, \phi(y) \rangle \langle y, \phi(x) \rangle \in [1, \infty) & (n = \infty) \end{cases}$$

Proof. Observe that since $r_{\phi(x)} = r_x \neq r_y = r_{\phi(y)}$, it follows that $\{x,y\}$ and $\{\phi(x),\phi(y)\}$ are both linearly independent, and hence conditions (D1) and (D2) are satisfied. Now let us set $R_1'' = R_2'' := \{r_x,r_y\}$ and $W_1'' = W_2'' := \langle \{r_x,r_y\}\rangle$, and furthermore, $\Phi_1'' := W_1''\{x,y\}$. Observe that Φ_1'' consists of elements of the form $\pm(\underbrace{\cdots r_y r_x r_y})x$ and

 $\pm(\underbrace{\cdots r_x r_y r_x})y$ (where $0 \le m < \operatorname{ord}(r_x r_y)$). Lemma 3.21 then yields

that $\Phi_1'' = \Phi_1''^+ \uplus \Phi_1''^-$, and consequently Theorem 2.3 yields that

$$\begin{cases} \langle x, \phi(y) \rangle \langle y, \phi(x) \rangle = \cos^2 \frac{\pi}{n} & (n \in \mathbb{N}, n \ge 2) \\ \langle x, \phi(y) \rangle \langle y, \phi(x) \rangle \in [1, \infty) & (n = \infty). \end{cases}$$

4. Acknowledgments

A few results presented in this paper are taken from the author's PhD thesis [12] and the author wishes to thank A/Prof. R. B. Howlett for all his help and encouragement throughout the author's PhD candidature. Due gratitude must be paid to Prof. M. Dyer for his penetrating insight and valuable suggestions. The author also wishes to thank Prof. G. I. Lehrer and Prof. R. Zhang for supporting this work.

REFERENCES

 A. Björner and F. Brenti, Combinatorics of Coxeter Groups, Graduate Texts in Mathematics, GTM 231, Springer, 2005

- N. Bourbaki, Groupes et algebras de Lie, Chapitres 4, 5 et 6, Hermann, Paris, 1968
- 3. B. Brink and R. B. Howlett, "A finiteness property and an automatic structure of Coxeter groups", *Math. Ann.*, **296** (1993), 179–190.
- B. Brink, "The set of dominance-minimal roots", J. Algebra, 206 (1998), 371–412
- F. Bergeron, N. Bergeron, R. B. Howlett and D. E. Taylor, "A Decomposition of the Descent Algebra of a Finite Coxeter Group", Journal of Algebraic Combinatorics, 1 (1992), 23–44
- 6. P. E. Caprace and B. Rémy, "Groups with a root group datum", *Innov. Incidence Geom.*, **9** (2009), 5–77.
- 7. W. A. Casselman, 'Computation in Coxeter groups I. Multiplication', *Electron. J. Combin.*, **9** (2002), no.1, Research Paper 25, 22 pp. (electronic).
- 8. W. A.Casselman, 'Computation in Coxeter groups II. Constructing Minimal Roots', *Represent. Theory*, **12**(2008), 260–293.
- 9. V. Deodhar, "On the root system of a Coxeter group", Comm. Algebra, 10 (1982), no. 6, 611–630.
- 10. M. Dyer, *Hecke algebras and reflections in Coxeter groups*, PhD thesis, University of Sydney, 1987.
- M. Dyer, "Reflection Subgroups of Coxeter Systems", J. Algebra, 135 (1990), 57–73.
- 12. X. Fu, Root systems and reflection representations of Coxeter groups, PhD thesis, University of Sydney, 2010.
- 13. X. Fu, "The dominance hierarchy in root systems of Coxeter groups", J. Algebra, **366** (2012), 187–204.
- 14. X.Fu, Non-orthogonal geometric realizations of Coxeter groups, arXiv:1112.3429 [math.RT], preprint, 2011.
- 15. J.-Y. Hée, "Système de racines sur un anneau commutatif totalement ordonné", Geom. Dedicata, 37 (1991), 65–102.
- 16. H. Hiller, *Geometry of Coxeter Groups*. Research Notes in Mathematics 54, Pitman (Advanced Publishing Program), Boston-London, 1981.
- 17. C. Hohlweg, J.P. Labbé and V. Ripoll, Asymptotical behaviour of roots of infinite Coxeter groups I, arXiv:1112.5415 [math.GR], preprint, 2011.
- 18. R. B. Howlett, "Normalizers of parabolic subgroups of reflection groups", *J. London Math. Soc.* (2), **21** (1980), no. 1, 62–80.
- 19. R. B. Howlett, *Introduction to Coxeter groups*, Lectures given at ANU, 1996 (available at http://www.maths.usyd.edu.au/res/Algebra/How/1997-6.html).
- 20. R. B. Howlett, P. J. Rowley and D. E. Taylor, "On Outer Automorphism groups of Coxeter Groups", *Manuscripta Math.*, **93** (1997), 499–513.
- J. Humphreys, Reflection Groups and Coxeter Groups, Cambridge Stud. Adv. Math. vol. 29, Cambridge Univ. Press, 1990.
- V. G. Kac, Infinite-dimensional Lie algebras, third edition, Cambridge University Press, Cambridge, 1990.
- 23. D. Krammer, *The conjugacy problem for Coxeter groups*, PhD thesis, Universiteit Utrecht, 1994.
- 24. G. A. Maxwell, "Sphere packings and hyperbolic reflection groups", *J. Algebra*, **79** (1982), 78–97.
- 25. R. V. Moody and A. Pianzola, "On infinite root systems", Trans. Amer. Math. Soc., 315 (1989), 661–696.
- 26. È. B. Vinberg, "Discrete Linear Groups Generated by Reflections", *Izv. Akad. Nauk SSSR Ser. Mat.*, **35** (1971), 1072–1112.