

OPTIONAL DECOMPOSITION FOR CONTINUOUS SEMIMARTINGALES UNDER ARBITRARY FILTRATIONS

IOANNIS KARATZAS AND CONSTANTINOS KARDARAS

ABSTRACT. We present an elementary treatment of the Optional Decomposition Theorem for continuous semimartingales and general filtrations. This treatment does not assume the existence of equivalent local martingale measure(s), only that of strictly positive local martingale deflator(s).

INTRODUCTION

The *Optional Decomposition Theorem* (ODT) is an important result in the field of Stochastic Analysis, and more particularly in Mathematical Finance. In one of its most “classical” forms, following [Kra96], the ODT can be stated as follows. For some $d \in \mathbb{N}$, let X be a \mathbb{R}^d -valued locally bounded semimartingale on a given filtered probability space $(\Omega, \mathcal{F}, \mathbb{P})$, $\mathbf{F} = \{\mathcal{F}(t)\}_{t \in \mathbb{R}_+}$, and assume that \mathcal{Q} , the collection of probability measures that are equivalent to \mathbb{P} and bestow the local martingale property to X , is non-empty. Then, a given nonnegative process V is a supermartingale under *all* probabilities in \mathcal{Q} , if and only if it admits the “optional” decomposition

$$(OD) \quad V = V(0) + \int_0^\cdot \langle H(t), dX(t) \rangle - C;$$

here H is a predictable X -integrable process, and C is a nondecreasing right-continuous adapted process with $C(0) = 0$.

The representation (OD) is relevant in the setting of Mathematical Finance. Indeed, suppose the components of X represent (discounted) prices of assets in a financial market. If $H = (H_i)_{i \in \{1, \dots, d\}}$ is the investment strategy of an agent in the market, where H_i stands for the units of asset i held in the portfolio for all $i \in \{1, \dots, d\}$, and C measures the agent’s aggregate consumption, then V in (OD) corresponds to the generated wealth-consumption process starting with initial capital $V(0)$. The ODT offers an illuminating “dual” characterization of all such wealth-consumption processes, as supermartingales under all *equivalent local martingale measures* of X . Using this

Date: February 5, 2015.

1991 Mathematics Subject Classification. 60H05; 60H30; 91B28.

Key words and phrases. Semimartingales; optional decomposition; local martingale deflators.

Research supported in part by the National Science Foundation under grant NSF-DMS-14-05210, and the Marie Curie Career Integration Grant (FP7-PEOPLE-2012-CIG), project number 334540. We are greatly indebted to Tomoyuki Ichiba and Johannes Ruf, for their careful reading of the manuscript and their many suggestions.

characterization, the ODT establishes the *superhedging duality* via use of dynamic programming techniques in non-Markovian settings.

Stochastic controllability results, similar to the ODT and obtained via martingale methods, can be traced as far back as [DV73] and, in the context of Mathematical Finance, to [KLSX91]. A version of the ODT when X is driven by Brownian motion under quasi-left-continuous filtrations appears in [EKQ95]. The first paper to treat the ODT for general locally bounded semimartingales is [Kra96], where functional (convex) analytic methods and results from [DS94] were employed. In [FK97], the more general case of constraints on investment is considered, using essentially similar arguments. In [FK98], the assumption of local boundedness on the semimartingale integrator X is dropped; more importantly, the authors avoid infinite-dimensional convex analysis, by following an alternative approach via predictable characteristics; this involves Lagrange multipliers, separating hyperplane arguments in Euclidean space, and measurable selections. Although the treatments of the ODT in the aforementioned papers are quite general, they do require a significant level of sophistication; indeed, they involve either use of difficult functional-analytic results, or deep knowledge of the General Theory of Processes as presented, e.g., in [JS03, Chapters I and II].

The present paper offers a rather elementary proof of the ODT for continuous-path semimartingale integrator X but *arbitrary* filtrations. Instead of assuming that $\mathcal{Q} \neq \emptyset$, it uses the more “localized” assumption that the class \mathcal{Y} of *strictly positive local martingale deflators* is non-empty. This assumption $\mathcal{Y} \neq \emptyset$ is both more general and more descriptive: it allows for an equivalent structural characterization of its validity by inspecting the local drift and local covariation processes of X , as mentioned in Theorem 1.1. (In fact, [SY98] treats the ODT using the condition $\mathcal{Y} \neq \emptyset$.) The important pedagogical element of the paper is that it avoids use of functional analysis and predictable characteristics in order to obtain the ODT. Since arbitrary filtrations support local martingales with potential jumps at both accessible and totally inaccessible times, it is impossible to avoid entirely the use of certain results from the general theory of Stochastic Processes. However, we feel that the path taken here is as elementary as possible. Although some intersection with previous work exists (notably, [EKQ95], as well as [Jac12] which deals with continuous asset prices and continuous filtrations), we believe that the present treatment is more straightforward.

1. THE SETTING

1.1. Preliminaries. We shall work on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, endowed with a filtration $\mathbf{F} = \{\mathcal{F}(t)\}_{t \in \mathbb{R}_+}$ which satisfies the usual hypotheses of right-continuity and augmentation by null sets. We stress that no further assumption is made on the filtration.

Let $X = (X_i)_{i \in \{1, \dots, d\}}$ be a d -dimensional semimartingale with continuous paths. We write $X = A + M$ for the Doob-Meyer decomposition of X ; here A is a d -dimensional process with continuous paths of finite variation and $A(0) = 0$, and M is a d -dimensional local martingale with continuous paths.

For $i \in \{1, \dots, d\}$, we shall denote by \check{A}_i the process of finite first variation associated with A_i . Upon defining $G := \sum_{i=1}^d (\check{A}_i + [M_i, M_i])$, it follows that there exist a d -dimensional predictable process a , and a predictable process c taking values in the set of nonnegative-definite matrices, such that $A = \int_0^\cdot a(t) dG(t)$ and $[M_i, M_j] = \int_0^\cdot c_{ij}(t) dG(t)$ hold for $i \in \{1, \dots, d\}$ and $j \in \{1, \dots, d\}$.

We shall denote by \mathcal{P} the predictable σ -field on $\Omega \times \mathbb{R}_+$, and by $\mathbb{P} \otimes G$ the measure on the product measurable space $(\Omega \times \mathbb{R}_+, \mathcal{P})$ which satisfies $(\mathbb{P} \otimes G)[J] = \mathbb{E} \left[\int_0^\infty \mathbf{1}_J(t) dG(t) \right]$ for all $J \in \mathcal{P}$.

Let $\mathcal{P}(X)$ denote the collection of all d -dimensional, predictable and X -integrable processes. A given d -dimensional predictable process H belongs to $\mathcal{P}(X)$ if and only if both processes $\int_0^\cdot \langle H(t), a(t) \rangle dG(t)$ and $\int_0^\cdot \langle H(t), c(t)H(t) \rangle dG(t)$ are finitely-valued.

We shall be using the notation

$$\mathcal{E}(Z) := \exp \left(Z - \frac{1}{2} [Z, Z]^c \right) \cdot \prod_{t \leq \cdot} (1 + \Delta Z(t)) \exp(-\Delta Z(t))$$

for the *stochastic exponential* of a scalar semimartingale Z with $Z(0) = 0$; note that it satisfies the integral equation $\mathcal{E}(Z) = 1 + \int_0^\cdot \mathcal{E}(Z)(t-) dZ(t)$.

1.2. Strictly positive local martingale deflators. We define \mathcal{Y} as the collection of all *strictly positive* local martingales Y with $Y(0) = 1$, such that YX^i is a local martingale for all $i \in \{1, \dots, d\}$. The next result gives conditions on the drift and local covariance structure of X which are equivalent to the requirement of non-emptiness for \mathcal{Y} .

Theorem 1.1. *In the above set-up, the following two conditions are equivalent:*

- (1) $\mathcal{Y} \neq \emptyset$.
- (2) *There exists a d -dimensional, predictable process ρ , such that $a = c\rho$ holds $(\mathbb{P} \otimes G)$ -a.e. and the process $\int_0^\cdot \langle \rho(t), c(t)\rho(t) \rangle dG(t)$ is finitely-valued.*

The structural conditions in statement (2) of Theorem 1.1 have appeared previously—see, for example, [Sch95] or [KS98, Theorem 4.2, page 12]. A proof of Theorem 1.1 can be found in [Kar10, Section 4]. We shall not repeat it here, but will provide some discussion in order to introduce important quantities that will be used later on.

1.3. Discussion of Theorem 1.1. Let us start by assuming that condition (2) of Theorem 1.1 holds. Since $a = c\rho$ implies that $\langle \rho, a \rangle = \langle \rho, c\rho \rangle = |\langle \rho, a \rangle|$ holds $(\mathbb{P} \otimes G)$ -a.e., it follows that ρ is X -integrable, i.e., $\rho \in \mathcal{P}(X)$. Then, the continuous-path semimartingale

$$(1.1) \quad \widehat{V} := \mathcal{E} \left(\int_0^\cdot \langle \rho(t), dX(t) \rangle \right)$$

is well-defined and satisfies the integral equation

$$(1.2) \quad \widehat{V} = 1 + \int_0^\cdot \widehat{V}(t) \langle \rho(t), dX(t) \rangle.$$

Straightforward computations show now that $(1/\widehat{V})$ is a local martingale, as is (X^i/\widehat{V}) for all $i \in \{1, \dots, d\}$; consequently, $(1/\widehat{V}) \in \mathcal{Y}$. In fact, whenever L is a local martingale with $L(0) = 0$, $\Delta L > -1$ and $[L, M] = 0$, the product $(1/\widehat{V}) \mathcal{E}(L)$ is an element of \mathcal{Y} . Although we shall not make direct use of this fact, let us also note that every element of \mathcal{Y} is of this product form.

The argument of the preceding paragraph establishes the implication (2) \Rightarrow (1) in Theorem 1.1. For completeness, we discuss now briefly, how the failure of condition (2) implies the failure of condition (1) in Theorem 1.1; detailed arguments can be found in [Kar10, Section 4]. Two contingencies need to be considered:

(i) *The vector a fails to be in the range of the matrix c , on a predictable set E of strictly positive $(\mathbb{P} \otimes G)$ -measure.* In this case one can find $\zeta \in \mathcal{P}(X)$, such that $c\zeta = 0$ and the process $\int_0^\cdot \langle \zeta(t), dX(t) \rangle = \int_0^\cdot \langle \zeta(t), a(t) \rangle dG(t)$ is nondecreasing everywhere and eventually strictly positive on E . This implies in a straightforward way that $\mathcal{Y} = \emptyset$.

(ii) *A d -dimensional predictable process ρ exists, so that $a = c\rho$ holds $(\mathbb{P} \otimes G)$ -a.e.; but the event $\{\int_0^T \langle \rho(t), c(t)\rho(t) \rangle dG(t) = \infty\}$ has positive probability for some $T > 0$.* Then, upon noting that $\rho \mathbf{1}_{\{|\rho| \leq n\}} \in \mathcal{P}(X)$ holds for all $n \in \mathbb{N}$, and defining $V_n = \mathcal{E}(\int_0^\cdot \langle \rho(t) \mathbf{1}_{\{|\rho(t)| \leq n\}}, dX(t) \rangle)$, one can show that the collection $\{V_n(T) \mid n \in \mathbb{N}\}$ is unbounded in probability. This again implies $\mathcal{Y} = \emptyset$.

Indeed, if \mathcal{Y} were not empty, then for any $Y \in \mathcal{Y}$ it would be straightforward to check that YV_n is a nonnegative local martingale—thus, a supermartingale—for all $n \in \mathbb{N}$. By Doob’s maximal inequality, this would imply that $\{Y(T)V_n(T) \mid n \in \mathbb{N}\}$ is bounded in probability; and since Y is strictly positive, that $\{V_n(T) \mid n \in \mathbb{N}\}$ is also bounded in probability. But we have already seen that the opposite is true.

Remark 1.2. A Reduction: Under condition (2) of Theorem 1.1, a given d -dimensional and predictable process H belongs to $\mathcal{P}(X)$ if and only if $\int_0^\cdot \langle H(t), c(t)H(t) \rangle dG(t)$ is finitely-valued. Indeed, the Cauchy-Schwartz inequality (see also [KS91, Proposition 3.2.14]) and the $(\mathbb{P} \otimes G)$ -identity $a = c\rho$ imply then

$$\int_0^\cdot |\langle H(t), a(t) \rangle| dG(t) \leq \left(\int_0^\cdot \langle H(t), c(t)H(t) \rangle dG(t) \right)^{1/2} \left(\int_0^\cdot \langle \rho(t), c(t)\rho(t) \rangle dG(t) \right)^{1/2},$$

and show that $\int_0^\cdot |\langle H(t), a(t) \rangle| dG(t)$ is *a fortiori* finitely-valued.

In obvious notation, we have $\mathcal{P}(X) = \mathcal{P}(M)$ under the condition (2) of Theorem 1.1.

Remark 1.3. An Interpretation: It follows from (1.2) that the process \widehat{V} can be interpreted as the strictly positive wealth generated by the predictable process ρ viewed as a “portfolio”, starting with a unit of capital. The components of X represent then the *returns* of the various assets in an equity market; the strictly positive processes $S_i = \mathcal{E}(X_i)$ are the *prices* of these assets; the components of ρ stand for the *proportions* of current wealth invested in each one of these assets;

whereas the scalar processes $\vartheta_i = (\widehat{V}/S_i)\rho_i$ (respectively, $\eta_i = \widehat{V}\rho_i$) keep track of the numbers of *shares* (resp., amounts of currency) invested in the various assets.

1.4. Optional Decomposition Theorem. The following is the main result of this work. It is proved in Section 2.

Theorem 1.4. *Assume that $\mathcal{Y} \neq \emptyset$. Let V be an adapted càdlàg process, locally bounded from below. Then, the following statements are equivalent:*

- (1) *The product YV is a local supermartingale, for all $Y \in \mathcal{Y}$.*
- (2) *The process V is of the form*

$$(1.3) \quad V = V(0) + \int_0^\cdot \langle H(t), dX(t) \rangle - C,$$

where $H \in \mathcal{P}(X)$ and C is a nondecreasing, right-continuous and adapted process with $C(0) = 0$, which is locally bounded from above.

Remark 1.5. On Uniqueness: In the present setting, the decomposition (1.3) of V is unique in the following sense: If $V = V(0) + \int_0^\cdot \langle H'(t), dX(t) \rangle - C'$ holds along with (1.3) for $H' \in \mathcal{P}(X)$ and C' a nondecreasing, right-continuous and adapted process with $C'(0) = 0$, then $C = C'$ and $\int_0^\cdot \langle H(t), dX(t) \rangle = \int_0^\cdot \langle H'(t), dX(t) \rangle$ hold modulo evanescence.

Indeed, let $\widehat{Y} := 1/\widehat{V}$, $D = C' - C$, $F := H' - H$, and note that $\widehat{Y}D = \widehat{Y} \int_0^\cdot \langle F(t), dX(t) \rangle$ is a continuous-path local martingale. This follows from integration-by-parts on the right-hand side of the last equality, the equation (1.2) for \widehat{V} , and property (2) in Theorem 1.1 for ρ . (For the latter local martingale property, see also the proof of the implication (2) \Rightarrow (1) in Theorem 1.4 in the beginning of §2.2.) Integrating by parts again, we see that $\int_0^\cdot \widehat{Y}(t) dD(t) = \widehat{Y}D - \int_0^\cdot D(t-) d\widehat{Y}(t)$ is both a continuous-path local martingale and a finite-variation process, which implies $\int_0^\cdot \widehat{Y}(t) dD(t) = 0$ modulo evanescence. Since \widehat{Y} is strictly positive, the last fact implies that $D = 0$ holds modulo evanescence, completing the argument.

Finally, let us note that for the integrands H and H' we may only conclude that $H = H'$ holds in the $(\mathbb{P} \otimes G)$ -a.e. sense.

2. PROOF OF THEOREM 1.4

We shall denote by \mathcal{L}^c the collection of all local martingales L with continuous paths and $L(0) = 0$; furthermore, we shall denote by \mathcal{L}^d the collection of all local martingales L with $\Delta L > -1$ and $L(0) = 0$ which are purely discontinuous, i.e., satisfy $[L, \Lambda] \equiv 0$ for all $\Lambda \in \mathcal{L}^c$.

2.1. An intermediate result. In order to prove Theorem 1.4, we first isolate the result that will enable us eventually to deal only with continuous-path local martingales.

Lemma 2.1. *Let B be a locally bounded from above semimartingale with the following properties:*

- $B + [B, L]$ is a local submartingale, for every $L \in \mathcal{L}^d$.
- $[B, L] = 0$, for every $L \in \mathcal{L}^c$.

Then, B is actually non-decreasing.

Proof. The first property implies that B itself is a local submartingale (just take $L \equiv 0$ there). Replacing B by $B - B(0)$, we may assume that $B(0) = 0$. Furthermore, standard localization arguments imply that we may take B to be bounded from above; therefore, we shall assume in the proof the existence of $b \in \mathbb{R}_+$ such that $B \leq b$. This means that B is an actual submartingale with last element $B(\infty)$, and that $\mathbb{E}[B(T)] \geq \mathbb{E}[B(0)] = 0$ holds for all stopping times T .

Consider now a countable collection $(\tau_n)_{n \in \mathbb{N}}$ of *predictable* stopping times that exhaust the accessible jump-times of B . Defining also the predictable set

$$J := \bigcup_{n \in \mathbb{N}} \llbracket \tau_n, \tau_n \rrbracket,$$

we note that the process

$$B^J := \int_{(0, \cdot]} \mathbf{1}_J(t) dB(t) = \sum_{n \in \mathbb{N}} \Delta B(\tau_n) \mathbf{1}_{\{\tau_n \leq \cdot\}}$$

is a local submartingale.

- We shall first show that B^J is nondecreasing, which amounts to showing that $\Delta B(\tau_n) \geq 0$ holds for all $n \in \mathbb{N}$. To this end, we define for each $n \in \mathbb{N}$ the $\mathcal{F}(\tau_n -)$ -measurable random variable

$$p_n := \mathbb{P}[\Delta B(\tau_n) < 0 \mid \mathcal{F}(\tau_n -)] \mathbf{1}_{\{\tau_n < \infty\}}.$$

On $\{\tau_n < \infty, p_n = 0\}$, we have $\Delta B(\tau_n) \geq 0$. For $n \in \mathbb{N}$ and $k \in \mathbb{N}$, we define $L_{n,k} \in \mathcal{L}^d$ to be the local martingale with $L_{n,k}(0) = 0$ and a single jump at τ_n , such that

$$\Delta L_{n,k}(\tau_n) = (1 - (1/k)) \mathbf{1}_{\{p_n > 0\}} ((1/p_n) \mathbf{1}_{\{\Delta B(\tau_n) < 0\}} - 1).$$

We note that $B + [B, L_{n,k}] = B$ holds on $\llbracket 0, \tau_n \llbracket$, while on the event $\{\tau_n < \infty, p_n > 0\}$ we have

$$\Delta(B + [B, L_{n,k}])(\tau_n) = \Delta B(\tau_n)(1 + \Delta L_{n,k}(\tau_n)) = \frac{\Delta B(\tau_n)}{k} - \frac{1 - (1/k)}{p_n} (\Delta B(\tau_n))^-.$$

The properties imposed on B imply, in particular, that $B + [B, L_{n,k}]$ is a local submartingale, bounded from above by $b + 1/k$ on $[0, \tau_n]$. It follows that on the event $\{\tau_n < \infty, p_n > 0\}$ we have $\mathbb{E}[\Delta B(\tau_n)(1 + \Delta L_{n,k}(\tau_n)) \mid \mathcal{F}(\tau_n -)] \geq 0$, which translates into

$$\mathbb{E}[(\Delta B(\tau_n))^- \mid \mathcal{F}(\tau_n -)] \leq \frac{\mathbb{E}[(\Delta B(\tau_n))^+ \mid \mathcal{F}(\tau_n -)]}{1 + ((k-1)/p_n)} \quad \text{on the event } \{\tau_n < \infty, p_n > 0\}$$

for all $k \in \mathbb{N}$. Sending $k \rightarrow \infty$, it follows that $\Delta B(\tau_n) \geq 0$ holds on $\{\tau_n < \infty, p_n > 0\}$.

• It remains to show that $B' := B - B^J = \int_{(0, \cdot]} \mathbf{1}_{(\Omega \times \mathbb{R}_+) \setminus J}(t) dB(t)$ is also a nondecreasing process. We note that B' inherits some of the properties of B : in particular, we have $B'(0) = 0$ and for every $L \in \mathcal{L}^d$, the process

$$B' + [B', L] = + \int_0^\cdot \mathbf{1}_{(\Omega \times \mathbb{R}_+) \setminus J}(t) (dB(t) + d[B, L](t))$$

is a local submartingale. We also have that $B' \leq B \leq b$, and additionally B' has jumps only at totally inaccessible stopping times. To ease the notation we write B instead of B' for the remainder of this proof, assuming from now onwards that B has jumps only at totally inaccessible stopping times. For each $n \in \mathbb{N}$, we define the local martingale

$$L_n := n \sum_{t \leq \cdot} \mathbf{1}_{\{\Delta B(t) \leq -1/n\}} - D_n,$$

where D_n is a suitable nondecreasing process with continuous paths (since the jumps of B are totally inaccessible). Note that $L_n \in \mathcal{L}^d$, which implies that $B + [B, L_n]$ is a local submartingale for all $n \in \mathbb{N}$. Furthermore, we note that $B + [B, L_n] = B + n \sum_{t \leq \cdot} \Delta B(t) \mathbf{1}_{\{\Delta B(t) \leq -1/n\}} \leq b$, which implies that $B + [B, L_n]$ is a true submartingale. Therefore, it follows that

$$n \mathbb{E} \left[- \sum_{t \leq n} \Delta B(t) \mathbf{1}_{\{\Delta B(t) \leq -1/n\}} \right] \leq \mathbb{E} [B(n) - B(0)] = \mathbb{E} [B(n)] \leq b, \quad \forall n \in \mathbb{N}.$$

The monotone convergence theorem gives now $\mathbb{E} \left[\sum_{t \in \mathbb{R}_+} (\Delta B(t))^- \right] = 0$, which implies $\Delta B \geq 0$. Continuing, we define for each $n \in \mathbb{N}$ a new local martingale

$$\tilde{L}_n := \tilde{D}_n - (1 - (1/n)) \sum_{t \leq \cdot} \mathbf{1}_{\{\Delta B(t) \geq 1/n\}},$$

where \tilde{D}_n is an appropriate continuous and nondecreasing process. Note that we have $\tilde{L}_n \in \mathcal{L}^d$ for all $n \in \mathbb{N}$, which implies that the processes

$$B + [B, \tilde{L}_n] = B - (1 - 1/n) \sum_{t \leq \cdot} \Delta B(t) \mathbf{1}_{\{\Delta B(t) \geq 1/n\}}, \quad n \in \mathbb{N}$$

are local submartingales, uniformly bounded from above by b . Thus, it follows that $\sum_{t \leq \cdot} \Delta B(t)$ is finitely-valued; and that $\tilde{B} := B - \sum_{t \leq \cdot} \Delta B(t)$ is a local submartingale. Recalling that the jumps of B occur only at totally inaccessible stopping times, we see that this last process \tilde{B} has continuous paths and is strongly orthogonal to all continuous-path local martingales, which means that it is of finite variation. Since it is a local submartingale, it has to be nondecreasing. It follows from this reasoning that the process $B = \left(B - \sum_{t \leq \cdot} \Delta B(t) \right) + \sum_{t \leq \cdot} \Delta B(t)$ is nondecreasing, and this concludes the argument. \square

2.2. Proof of Theorem 1.4. The implication (2) \Rightarrow (1) is straightforward. Indeed, we fix $Y \in \mathcal{Y}$ and note that

$$Z := \int_0^\cdot Y(t-) dX(t) + [Y, X] = YX - X(0) - \int_0^\cdot X(t) dY(t)$$

is a d -dimensional local martingale. Then, if $V = V(0) + \int_0^\cdot \langle H(t), dX(t) \rangle - C$, one computes

$$YV = V(0) + \int_0^\cdot V(t-) dY(t) + \int_0^\cdot \langle H(t), dZ(t) \rangle - \int_0^\cdot Y(t) dC(t),$$

which shows that YV is a local supermartingale.

• For the implication (1) \Rightarrow (2), let us assume that V is such that YV is a local supermartingale for all $Y \in \mathcal{Y}$. In particular, recalling the notation of (1.1), we note that (V/\widehat{V}) is a local supermartingale and write this process in its Kunita-Watanabe / Doob-Meyer representation

$$(2.1) \quad U := V/\widehat{V} = V(0) + \int_0^\cdot \langle \theta(t), dM(t) \rangle + N - B.$$

Here $\theta \in \mathcal{P}(X)$ (see Remark 1.2), and $N \in \mathcal{L}^c$ satisfies $[N, M] = 0$, whereas B is a local submartingale with $B(0) = 0$ and “purely discontinuous”, in the sense that

$$(2.2) \quad [B, L] = 0 \quad \text{holds for every } L \in \mathcal{L}^c.$$

In particular, we note that $[B, N] = 0$.

(i): The first item of business is to show that the process B in (2.1) is actually *nondecreasing*; for this we shall use Lemma 2.1. Since $N + \int_0^\cdot \langle \theta(t), dM(t) \rangle$ is continuous and U locally bounded from below, the process $B = V(0) + N + \int_0^\cdot \langle \theta(t), dM(t) \rangle - U$ is locally bounded from above.

Let us fix $L \in \mathcal{L}^d$. From $(1/\widehat{V}) \mathcal{E}(L) \in \mathcal{Y}$ we observe – e.g., using the product rule – that the process

$$V \cdot (1/\widehat{V}) \mathcal{E}(L) = U \mathcal{E}(L) = \left(V(0) + N + \int_0^\cdot \langle \theta(t), dM(t) \rangle - B \right) \mathcal{E}(L)$$

is a local supermartingale. Furthermore, the process $(N + \int_0^\cdot \langle \theta(t), dM(t) \rangle) \mathcal{E}(L)$ is a local martingale, and it follows from these two observations that

$$B \mathcal{E}(L) = \int_0^\cdot \mathcal{E}(L)(t-) d(B + [B, L])(t) + \int_0^\cdot (B \mathcal{E}(L))(t-) dL(t)$$

is a local submartingale. This, in turn, implies that

$$B + [B, L] = \int_0^\cdot \frac{1}{\mathcal{E}(L)(t-)} d(BL)(t) - \int_0^\cdot B(t-) dL(t)$$

is also a local submartingale. Recalling the property (2.2) and invoking Lemma 2.1, we conclude from this observation that the local submartingale B in the decomposition (2.1) is indeed nondecreasing.

(ii): The second item of business is to show that $N \equiv 0$ holds in (2.1). The crucial observation here is that, because of $[N, M] = 0$, the product $(1/\widehat{V})\mathcal{E}(nN)$ is an element of \mathcal{Y} for all $n \in \mathbb{N}$. As a consequence, $U\mathcal{E}(nN) = V \cdot (1/\widehat{V})\mathcal{E}(nN)$ is a local supermartingale for all $n \in \mathbb{N}$.

Since $[\mathcal{E}(nN), B] = 0$ and $[\mathcal{E}(nN), M] = 0$, it follows that $\mathcal{E}(nN)(B - N)$ is a local submartingale for all $n \in \mathbb{N}$. Now we observe

$$\mathcal{E}(nN)(B - N) = \int_0^\cdot (B - N)(t-) d\mathcal{E}(nN)(t) + \int_0^\cdot \mathcal{E}(nN)(t-) d(B - N - [nN, N])(t),$$

from which it follows that $B - n[N, N]$ is a local submartingale for all $n \in \mathbb{N}$. This is only possible if $[N, N] = 0$ which, since $N(0) = 0$, implies $N \equiv 0$; as a result, (2.1) becomes

$$(2.3) \quad U = (V/\widehat{V}) = V(0) + \int_0^\cdot \langle \theta(t), dM(t) \rangle - B.$$

(iii): We are now in a position to conclude. Yet another application of the integration-by-parts formula on $V = \widehat{V}U$ gives, in conjunction with (1.2), (2.3) and Theorem 1.1, the decomposition

$$\begin{aligned} V &= V(0) + \int_0^\cdot U(t-) d\widehat{V}(t) + \int_0^\cdot \widehat{V}(t) dU(t) + [\widehat{V}, U] \\ &= V(0) + \int_0^\cdot U(t-) d\widehat{V}(t) + \int_0^\cdot \widehat{V}(t) \langle \theta(t), dX(t) \rangle - \int_0^\cdot \widehat{V}(t) dB(t) \\ &= V(0) + \int_0^\cdot \widehat{V}(t) \langle U(t-)\rho(t) + \theta(t), dX(t) \rangle - \int_0^\cdot \widehat{V}(t) dB(t). \end{aligned}$$

Defining $U_-(t) := U(t-)$ for $t > 0$, as well as

$$H := \widehat{V}(U_- \rho + \theta) \in \mathcal{P}(X) \quad \text{and} \quad C := \int_0^\cdot \widehat{V}(t) dB(t),$$

we obtain the decomposition (1.3) as claimed. \square

REFERENCES

- [DS94] F. Delbaen and W. Schachermayer. A general version of the fundamental theorem of asset pricing. *Math. Ann.*, 300(3):463–520, 1994.
- [DV73] M. H. A. Davis and P. Varaiya. Dynamic programming conditions for partially observable stochastic systems. *SIAM J. Control*, 11:226–261, 1973.
- [EKQ95] Nicole El Karoui and Marie-Claire Quenez. Dynamic programming and pricing of contingent claims in an incomplete market. *SIAM J. Control Optim.*, 33(1):29–66, 1995.
- [FK97] H. Föllmer and D. Kramkov. Optional decompositions under constraints. *Probab. Theory Related Fields*, 109(1):1–25, 1997.
- [FK98] H. Föllmer and Yu. M. Kabanov. Optional decomposition and Lagrange multipliers. *Finance Stoch.*, 2(1):69–81, 1998.
- [Jac12] S. Jacka. A simple proof of Kramkov’s result on uniform supermartingale decompositions. *Stochastics*, 84(5-6):599–602, 2012.
- [JS03] J. Jacod and A. N. Shiryaev. *Limit theorems for stochastic processes*, volume 288 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 2003.

- [Kar10] C. Kardaras. Finitely additive probabilities and the fundamental theorem of asset pricing. In *Contemporary quantitative finance*, pages 19–34. Springer, Berlin, 2010.
- [KLSX91] I. Karatzas, J. P. Lehoczky, S. E. Shreve, and G.-L. Xu. Martingale and duality methods for utility maximization in an incomplete market. *SIAM J. Control Optim.*, 29(3):702–730, 1991.
- [Kra96] D. O. Kramkov. Optional decomposition of supermartingales and hedging contingent claims in incomplete security markets. *Probab. Theory Related Fields*, 105(4):459–479, 1996.
- [KS91] I. Karatzas and S. E. Shreve. *Brownian motion and stochastic calculus*, volume 113 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1991.
- [KS98] I. Karatzas and S. E. Shreve. *Methods of mathematical finance*, volume 39 of *Applications of Mathematics (New York)*. Springer-Verlag, New York, 1998.
- [Sch95] M. Schweizer. On the minimal martingale measure and the Föllmer-Schweizer decomposition. *Stochastic Anal. Appl.*, 13(5):573–599, 1995.
- [SY98] C. Stricker and J. A. Yan. Some remarks on the optional decomposition theorem. In *Séminaire de Probabilités, XXXII*, volume 1686 of *Lecture Notes in Math.*, pages 56–66. Springer, Berlin, 1998.

IOANNIS KARATZAS: DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, NEW YORK, NY 10027, USA;
INTECH INVESTMENT MANAGEMENT, ONE PALMER SQUARE, SUITE 441, PRINCETON, NJ 08542, USA
E-mail address: ik@math.columbia.edu, ik@enhanced.com

CONSTANTINOS KARDARAS: DEPARTMENT OF STATISTICS, LONDON SCHOOL OF ECONOMICS, 10 HOUGHTON
STREET, LONDON, WC2A 2AE, ENGLAND
E-mail address: k.kardaras@lse.ac.uk