THE DERIVED CATEGORY OF A GRADED GORENSTEIN RING

JESSE BURKE AND GREG STEVENSON

Abstract. We give an exposition and generalization of Orlov's theorem on graded Gorenstein rings. We show the theorem holds for non-negatively graded rings which are Gorenstein in an appropriate sense and whose degree zero component is an arbitrary non-commutative right noetherian ring of finite global dimension. A short treatment of some foundations for local cohomology and Grothendieck duality at this level of generality is given in order to prove the theorem. As an application we give an equivalence of the derived category of a commutative complete intersection with the homotopy category of graded matrix factorizations over a related ring.

1. INTRODUCTION

Let A be a graded Gorenstein ring. In [\[17\]](#page-23-0) Orlov related the bounded derived category of coherent sheaves on $\text{Proj } A$ and the singularity category of graded A modules via fully faithful functors; the exact relation depends on the a-invariant of A. This is a striking theorem that has found applications in physics, algebraic geometry and representation theory. To give an idea of the scope of the theorem: in the limiting case that A has finite global dimension (so the singularity category is trivial), it recovers (and generalizes to non-commutative rings) Beilinson's result [\[5\]](#page-22-0) that the derived category of Proj A is generated by a finite sequence of twists of the structure sheaf.

There has been much work related to this theorem. The idea of Orlov's construction perhaps first appears in Van Den Bergh's paper on non-commutative crepant resolutions [\[24\]](#page-23-1) where he described functors similar to those considered by Orlov, in the case of torus invariants. After Orlov's paper appeared, the idea was further explored by the physicists Herbst, Hori and Page in [\[12\]](#page-23-2). In turn these ideas were the inspiration for two papers on the derived category of GIT quotients [\[3,](#page-22-1) [11\]](#page-23-3). Segal and then Shipman gave geometric proofs of Orlov's theorem for commutative hypersurfaces in [\[20\]](#page-23-4) and [\[21\]](#page-23-5). Related results are [\[4\]](#page-22-2) and [\[13\]](#page-23-6). The theorem has been used in similar ways in [\[2,](#page-22-3) [14\]](#page-23-7). Finally, it has been used in representation theory; especially in the study of weighted projective lines, see e.g. [\[15\]](#page-23-8).

Orlov assumed that A_0 a field. In this paper, which we consider largely expository, we generalize his result to show that the same relation holds when A_0 is a non-commutative noetherian ring of finite global dimension. This has an immediate application to commutative complete intersection rings and we expect there to be further applications, for instance to (higher) preprojective algebras. The structure of our proof is very close to Orlov's original arguments. We give many details and we hope that these details may help the reader (even one only interested in algebras defined over a field) to better understand Orlov's work.

The main tool in the proof is a semiorthogonal decomposition; this separates a triangulated category into an admissible subcategory and its orthogonal. Derived global sections gives an embedding of $D^b(\text{Proj } A)$ into $D^b(\text{gr}_{\geq i} A)$ as an admissible subcategory. When A_0 is a field, Orlov showed that there is an embedding of the singularity category $D^b_{sg}(\text{gr }A)$ into $D^b(\text{gr}_{\geq i}A)$. The existence of such an embedding is rather remarkable and constitutes perhaps the key insight required to prove the theorem. Orlov then used Grothendieck duality in a very clever way to compare the orthogonals of $\mathsf{D}^{\mathsf{b}}(\text{Proj } A)$ and $\mathsf{D}^{\mathsf{b}}_{\mathsf{sg}}(\text{gr } A)$ inside of $\mathsf{D}^{\mathsf{b}}(\text{gr}_{\geq i} A)$. For example when A is Calabi-Yau, the orthogonals coincide and so there is an equivalence between $\mathsf{D}^{\mathsf{b}}(\operatorname{Proj} A)$ and $\mathsf{D}^{\mathsf{b}}_{\mathsf{sg}}(\operatorname{gr} A)$.

The arguments we present here follow those of Orlov. The main addition is the observation that, when A_0 has finite global dimension, one can construct particularly nice resolutions of complexes of graded modules with bounded finitely generated cohomology. This allows us to prove Orlov's embedding $D_{sg}^{b}(gr A) \rightarrow$ $D^{b}(gr_{\geq i} A)$ is valid in this more general setting. We also need to develop some foundations concerning local cohomology and Grothendieck duality over non-commutative rings to prove analogues of the other steps of Orlov's proof. These foundations do not seem to be contained in the literature in the form and generality that we need, although the arguments we give here are relatively straightforward generalizations of arguments by Artin and Zhang [\[1\]](#page-22-4).

Let us give a quick summary of the paper. The second section contains some categorical background, especially on semiorthogonal decompositions. The third section is devoted to the derived category of graded modules over a graded ring, and some standard semiorthogonal decompositions that appear there. This section contains the key observation, Lemma [3.10,](#page-5-0) needed to prove the embedding of the singularity category works for the rings we work with. The fourth section deals with local cohomology and the semiorthogonal decomposition it gives, while the fifth deals with the embedding of the singularity category, Grothendieck duality, and the semiorthogonal decomposition these give. The sixth section contains the proof of the main result as well as a sufficient condition for a Gorenstein ring to satisfy Artin and Zhang's condition χ , which is necessary for the proof. Finally, in the last section, we apply the main theorem to give a description of the bounded derived category of a complete intersection ring in terms of graded matrix factorizations.

Acknowledgements. Ragnar-Olaf Buchweitz gave two beautiful lectures on Orlov's theorem at Bielefeld University in July 2011 that inspired this and from which we learned a lot. The existence of the crucial left adjoint to the projection onto the singularity category over general bases of finite global dimension is implicitly contained in Ragnar's exposition of Orlov's work. We thank Mark Walker for several helpful conversations on the contents of the paper.

2. Background

We recall here some standard results on semiorthogonal decompositions of triangulated categories which we will need. Throughout this section $\mathcal T$ denotes a triangulated category.

Definition 2.1. For \mathcal{D} a triangulated subcategory of \mathcal{T} , define \mathcal{D}^{\perp} to be the full subcategory with objects those $X \in \mathcal{T}$ such that $\text{Hom}_{\mathcal{T}}(D, X) = 0$ for all objects D of D. Similarly, $\pm \mathcal{D}$ has objects those X with $\text{Hom}_{\mathcal{T}}(X, D) = 0$ for all D in D. Both \mathcal{D}^{\perp} and $^{\perp} \mathcal{D}$ are thick subcategories of \mathcal{T} i.e., they are triangulated subcategories which are closed under taking direct summands.

Definition 2.2. A triangulated subcategory \mathcal{D} of \mathcal{T} is *left admissible* in \mathcal{T} if the inclusion functor $i : \mathcal{D} \to \mathcal{T}$ has a left adjoint; \mathcal{D} is *right admissible* if i has a right adjoint.

The following criterion for admissibility can be found as [\[6,](#page-22-5) Lemma 3.1].

Lemma 2.3. Let D be a triangulated subcategory of T .

(1) *The category* D *is left admissible if and only if for every* X *in* T *there is a triangle:*

$$
E_X \to X \to D_X \to \Sigma E_X
$$

with D_X *in* D *and* E_X *in* $\perp D$ *.*

(2) *The category* D *is right admissible if and only if for every* X *in* $\mathcal T$ *there is a triangle:*

$$
D_X \to X \to E_X \to \Sigma D_X
$$

with D_X *in* \mathcal{D} *and* E_X *in* \mathcal{D}^{\perp} *.*

Corollary 2.4. *A subcategory* D *of* T *is left admissible if and only if* $\perp D$ *is right admissible.* In this case $({}^{\perp}D)^{\perp} = D$.

Definition 2.5. A *semiorthogonal decomposition* of T is a pair of subcategories A and B such that A is left admissible and $\mathcal{B} = \perp \mathcal{A}$ (equivalently, B is right admissible and $\mathcal{A} = \mathcal{B}^{\perp}$. We write this as

$$
\mathcal{T}=(\mathcal{A},\mathcal{B})\,.
$$

The following useful lemma essentially follows from Lemma [2.3](#page-2-0) in a straightforward way.

Lemma 2.6. *There is a semiorthogonal decomposition* $\mathcal{T} = (\mathcal{A}, \mathcal{B})$ *if and only if* $\mathcal{B} \subset \perp \mathcal{A}$ *and for every* X *in* T *there is a triangle*

$$
B_X \to X \to A_X \to
$$

with $B_X \in \mathcal{B}$ *and* $A_X \in \mathcal{A}$ *. We will call such a triangle the* localization triangle *for* X*.*

Orlov generalized the definition of semiorthogonal decomposition in [\[17\]](#page-23-0) to:

Definition 2.7. A sequence of full triangulated subcategories $(\mathcal{D}_1, \ldots, \mathcal{D}_n)$ of \mathcal{T} is a *semiorthogonal decomposition* if for each $i = 1, \ldots, n - 1$, the thick subcategory generated by $\mathcal{D}_1, \ldots, \mathcal{D}_i$, which we denote $\langle \mathcal{D}_1, \ldots, \mathcal{D}_i \rangle$, is left admissible and

$$
^{\perp}\left\langle \mathcal{D}_{1},\ldots,\mathcal{D}_{i}\right\rangle =\langle\mathcal{D}_{i+1},\ldots,\mathcal{D}_{n}\rangle.
$$

We can (and will) construct semi-orthogonal decompositions inductively:

Lemma 2.8. Let $\mathcal{T} = (\mathcal{A}, \mathcal{B})$, $\mathcal{A} = (\mathcal{D}_1, \dots, \mathcal{D}_i)$, and $\mathcal{B} = (\mathcal{D}_{i+1}, \dots, \mathcal{D}_n)$ be *semiorthogonal decompositions. Then*

$$
\mathcal{T}=(\mathcal{D}_1,\ldots,\mathcal{D}_n)
$$

is a semiorthogonal decomposition.

3. The bounded derived category of graded modules

We now provide some preliminary results on derived categories of graded modules. We begin by exhibiting some semiorthogonal decompositions of the bounded derived category which we will need in the sequel (and which are relatively straightforward generalisations of those in Orlov's work). We also prove the main technical results concerning graded projectives and graded projective resolutions which we will need for our extension of Orlov's result.

In this section $A = \bigoplus_{i \geq 0} A_i$ is a positively graded noetherian ring with A_0 a ring of finite global dimension. All modules will be right modules unless otherwise stated. We denote by $g \circ A$ the abelian category of finitely generated graded Amodules and degree zero homogeneous maps. If $M = \bigoplus M_i$ is a graded A-module, then $M(1)$ is the graded A-module with $M(1)_i = M_{i+1}$.

We denote by $gr_{\geq i} A$ the full subcategory of gr A consisting of objects M such that $M_j = 0$ for all $j < i$. This is an abelian subcategory of gr A and there is an adjoint pair of functors

$$
\operatorname{gr}_{\geq i} A \xrightarrow[(-)_{\geq i}]{\operatorname{inc}} \operatorname{gr} A
$$

where $M_{\geq i} = \bigoplus_{j \geq i} M_j$ is right adjoint to the inclusion.

We denote by $\overline{D}^b(-)$ the bounded derived category of an abelian category. Both functors of the above adjoint pair are exact and so induce functors

$$
\mathsf{D}^{\mathsf{b}}(\operatorname{gr}_{\geq i} A) \xrightarrow[(-)_{\geq i}]{\operatorname{inc}} \mathsf{D}^{\mathsf{b}}(\operatorname{gr} A)
$$

which also form an adjoint pair. The functor induced by inclusion is fully faithful and the essential image is the full subcategory of D^b (gr A) consisting of objects M such that $H^j(M) \in \text{gr}_{\geq i} A$ for all $j \in \mathbb{Z}$. We denote this subcategory also by $D^{\mathsf{b}}(\mathrm{gr}_{\geq i} A)$; it is a right admissible subcategory.

Definition 3.1. Define $S_{\leq i}$ to be the thick subcategory generated by the objects $A_0(e)$, for all $e > -i$ and $S_{\geq i}$ to be the thick subcategory generated by $A_0(e)$, for all $e \leq -i$.

Lemma 3.2. An object M of $D^b(\text{gr }A)$ is in $\mathcal{S}_{\leq i}$ if and only if $M_{\geq i} \simeq 0$.

Proof. The full subcategory with objects those M satisfying $M_{\geq i} \simeq 0$ is thick by virtue of being the kernel of an exact functor. Since $A_0(e)_{\geq i} = 0$ for all $e > -i$, we see that $\mathcal{S}_{\leq i}$ is contained in this thick subcategory. Thus if M is in $\mathcal{S}_{\leq i}$ we must have $M_{\geq i} \simeq 0$.

For the converse, we first assume M is a module, i.e. concentrated in homological degree 0, and that there is an integer $e < i$ with $M_j = 0$ for all $j \neq e$. Since M_e is a finitely generated A_0 -module and A_0 has finite global dimension, M_e has a finite projective resolution over A_0 . Over A this says that M_e is in the thick subcategory generated by $A_0(-e)$, which is contained in $\mathcal{S}_{\leq i}$.

Now suppose M is a non-zero finitely generated graded A-module with $M_{\geq i} \simeq 0$. As M is finitely generated there is an integer j with $M_{\geq j} = M$ and we may as well choose a maximal such j , which is necessarily less than i . Consider the triangle

$$
M_{\geq j+1} \to M \to M_j \to
$$

By the previous argument M_j is in $\mathcal{S}_{\leq i}$ and arguing inductively on the number of degrees in which M is non-zero we see that $M_{\geq j+1}$ is in $\mathcal{S}_{\leq i}$, and hence M is in $\mathcal{S}_{\leq i}$.

For an arbitrary non-zero object $M \in \mathsf{D}^{\mathsf{b}}(\text{gr } A)$, with $M_{\geq i} \simeq 0$ the result follows from induction on the number of non-vanishing cohomology modules, using the triangle

$$
M^{
$$

where $n = \max\{j | H^j(M) \neq 0\}$ and $M^{ is the truncation with respect to the$ standard t-structure.

Remark 3.3. It follows from the definition that $S_{\geq i}$ is contained in $D^b(\text{gr}_{\geq i} A)$. We will show in Lemma [4.17](#page-13-0) that $\mathcal{S}_{\geq i}$ is the full subcategory of $\mathsf{D}^{\mathsf{b}}(\mathrm{gr}_{\geq i} A)$ whose objects have torsion cohomology.

Lemma 3.4. *There is a semiorthogonal decomposition*

$$
\mathsf{D}^{\mathsf{b}}(\operatorname{gr} A) = \left(\mathcal{S}_{\leq i}, \mathsf{D}^{\mathsf{b}}(\operatorname{gr}_{\geq i} A) \right).
$$

The localization triangle for $M \in D^b(\text{gr } A)$ *is given by the canonical maps*

$$
M_{\geq i} \to M \to M/M_{\geq i} \to.
$$

Proof. Let M be in $\mathcal{S}_{\leq i}$ and N be in $D^{\mathsf{b}}(\mathrm{gr}_{\geq i} A)$. Then we have that

$$
\operatorname{Hom}\nolimits_{\mathsf{D}^{\mathsf{b}}(\operatorname{gr}\nolimits A)}(N,M)\cong \operatorname{Hom}\nolimits_{\mathsf{D}^{\mathsf{b}}(\operatorname{gr}\nolimits_{\geq i}A)}(N,M_{\geq i})=0
$$

by right adjointness of $(-)_{\geq i}$ and since $M_{\geq i} \simeq 0$. Thus $D^{\mathsf{b}}(\mathrm{gr}_{\geq i} A) \subseteq {}^{\perp} \mathcal{S}_{\leq i}$. If M is any object in $\mathsf{D}^{\mathsf{b}}(\text{gr}\,A)$ we have the triangle

$$
M_{\geq i}\to M\to M/M_{\geq i}\to
$$

with $M_{\geq i}$ in $\mathsf{D}^{\mathsf{b}}(\mathrm{gr}_{\geq i} A)$ and $M/M_{\geq i}$ in $\mathcal{S}_{\leq i}$. Thus we may apply Lemma [2.6.](#page-2-1) \Box

Definition 3.5. Define $P_{\leq i}$ to be the thick subcategory generated by the objects $A(e)$ for all $e > -i$ and $\mathcal{P}_{\geq i}$ to the thick category generated by $A(e)$ for all $e \leq -i$.

Remark 3.6. It follows from the definition that $\mathcal{P}_{\geq i}$ is contained in $D^b(\text{gr}_{\geq i} A)$. In fact, $\mathcal{P}_{\geq i}$ is the full subcategory of $\mathsf{D}^{\mathsf{b}}(\mathrm{gr}_{\geq i} A)$ whose objects are perfect complexes of A-modules.

Lemma 3.7. *There is a semiorthogonal decomposition*

$$
\mathsf{D}^{\mathsf{b}}(\operatorname{gr} A) = (\mathsf{D}^{\mathsf{b}}(\operatorname{gr}_{\geq i} A), \mathcal{P}_{< i}).
$$

Before proving this lemma, we need two results on graded projective A-modules and graded projective resolutions over A.

Lemma 3.8. *Let* P *be a finitely generated graded projective* A*-module. Then there is an isomorphism, for some integers* n, m_1, \ldots, m_n ,

$$
P \cong \bigoplus_{i=1}^n P_i \otimes_{A_0} A(m_i)
$$

where the P_i *are projective right* A_0 *-modules.*

Proof. Let P be a non-zero finitely generated graded projective A-module. Consider the graded projective A_0 -module $\overline{P} = P \otimes_A A_0$ which is non-zero by the graded Nakayama lemma. We obtain a graded projective A-module $\overline{P} \otimes_{A_0} A$ fitting into a commutative diagram

where the vertical morphism is the canonical one and the dashed arrow exists by projectivity of $\overline{P} \otimes_{A_0} A$. By construction the morphism $\overline{P} \otimes_{A_0} A \to P$ is surjective so it splits. But

$$
\overline{P}\otimes_{A_0}A\otimes_A A_0\cong \overline{P}=P\otimes_A A_0
$$

and so by another application of the graded Nakayama lemma we see $P \cong \overline{P} \otimes_{A_0} A$ is induced up from a graded projective A_0 -module proving the lemma.

Definition 3.9. For every graded projective A -module Q , we define summands $Q_{\prec i}$ and $Q_{\succ i}$ with $Q_{\prec i}$ in $\mathcal{P}_{\leq i}$ and $Q_{\succ i}$ in $\mathcal{P}_{\geq i}$ via the unique up to isomorphism split exact sequence of graded projective modules

$$
0 \to Q_{\prec i} \to Q \to Q_{\succcurlyeq i} \to 0
$$

which exists by the previous lemma.

The next lemma is a key technical observation concerning the structure of resolutions over A.

Lemma 3.10. *Every object* M *in* D b (gr A) *is quasi-isomorphic to a complex of finitely generated graded projective* A*-modules*

$$
P = \cdots \to P^j \to P^{j+1} \to \cdots
$$

such that $P^j = 0$ *for all* $j \gg 0$ *and for any* $i \in \mathbb{Z}$ *there exists a* k_i *with* P^{-k} *in* $\mathcal{P}_{\geq i}$ *for all* $k \geq k_i$ *.*

Proof. It is sufficient to prove the result for graded A-modules as the condition is closed under suspensions and taking cones, and every object of $D^b(\text{gr }A)$ can be written as an iterated extension of suspensions of modules using the standard tstructure. Let us introduce notation local to this proof. Given a finitely generated A-module M define the integer

$$
\min(M) = \min\{i \in \mathbb{Z} \mid M_i \neq 0\}.
$$

Let M be a finitely generated graded A-module and set

$$
M = M \otimes_A A_0 = M/A_{\geq 1}M,
$$

which we consider as a graded A_0 -module. We may assume M has infinite projective dimension as the result is trivial in the finite projective dimension case. We will construct a projective resolution of the desired form. If \overline{M} is zero then, by Nakayama, so is M and thus we may suppose $\overline{M} \neq 0$. We choose an epimorphism from a graded projective A_0 -module $\overline{P^0} \to \overline{M}$ by writing $\overline{M} \cong \bigoplus_{i=1}^n M_i(a_i)$, taking epimorphisms $P_i^0(a_i) \to M_i(a_i)$ where the P_i^0 are projective A_0 -modules and

setting $\overline{P^0} = \bigoplus_{i=1}^n P_i^0(a_i)$ with the obvious morphism to \overline{M} . This gives rise to an exact sequence of graded A-modules

$$
0 \to Z^0 \to P^0 \to M \to 0,
$$

where $P^0 = \overline{P^0} \otimes_{A_0} A$, with the property that

$$
\min(Z^0) \ge \min(P^0) = \min(M).
$$

We have assumed A_0 has finite global dimension, say d. Proceeding as above we may find projectives P^i for $i = 1, ..., d - 1$ and exact sequences

$$
0 \to Z^i \to P^i \to Z^{i-1} \to 0
$$

with $\min(Z^i) \ge \min(P^i) = \min(Z^{i-1})$. Thus, restriction to the graded components in degree $j = min(M)$ gives an exact sequence

$$
0 \to Z_j^{d-1} \to P_j^{d-1} \to \cdots \to P_j^0 \to M_j \to 0
$$

of A_0 -modules with the P_j^i projective. As A_0 has global dimension d we see Z_j^{d-1} must be projective. Hence $\overline{Z^{d-1}}$ can be written as $Z_j^{d-1} \oplus X$ with X living in degrees strictly greater than j. As before we can pick an epimorphism $Q \to X$ from a graded projective A_0 -module Q which lives in the same degrees as X. Setting $P^{\bar{d}} = (Z_j^{d-1} \oplus Q) \otimes_{A_0} A$ we get a short exact sequence

$$
0 \to Z^d \to P^d \to Z^{d-1} \to 0
$$

with $\min(Z^d) > \min(M)$; thus our recipe guarantees projectives with generators in degrees less than or equal to $\min(M)$ cannot occur beyond the dth step of the resolution. We can now repeat this procedure starting at Z^d to obtain a resolution satisfying the desired properties.

Remark 3.11*.* It is easy to construct examples showing this lemma is no longer true if A_0 does not have finite global dimension. Indeed, let $A = k[x, y]/(x^2, y^2)$, with $|x| = 0$ and $|y| = 1$. The resolution of $A/(x)$ is

$$
\dots \to A \xrightarrow{x} A \xrightarrow{x} A \to 0
$$

which does not satisfy the conclusion of the previous lemma.

Proof of [3.7.](#page-4-0) Given an object M in $D^b(\text{gr }A)$, let $P \xrightarrow{\simeq} M$ be a quasi-isomorphism where P is a complex of projectives as in the previous lemma. Apply the decom-position [3.9](#page-5-1) degree-wise to P to get a triangle

$$
P_{\prec i} \to P \to P_{\succcurlyeq i} \to
$$

where $P_{\prec i}$ is the subcomplex of P consisting of all projective summands generated in degrees less than i and $P_{\geq i}$ is the quotient complex consisting of all projective summands generated in degree at least *i*. Since P^{-k} is in $\mathcal{P}_{\geq i}$ for all $k \gg 0$, we see that $P_{\prec i}$ is bounded, and hence in $\mathcal{P}_{\leq i}$. Note that $P_{\succ i}$ has bounded finitely generated cohomology by the triangle, and so must be in $\mathsf{D}^{\mathsf{b}}(\mathrm{gr}_{\geq i} A)$.

Now observe that there are no non-zero maps from objects in $\mathcal{P}_{\leq i}$ to any module M in $\operatorname{gr}_{\geq i} A$. Thus $\operatorname{gr}_{\geq i} A$ is contained in $\mathcal{P}_{\leq i}^{\perp}$ and hence so is $\mathsf{D}^{\mathsf{b}}(\operatorname{gr}_{\geq i} A)$ since it is generated by $gr_{\geq i} A$ and $\mathcal{P}_{\leq i}^{\perp}$ is thick. Thus $\mathcal{P}_{\leq i} \subseteq {}^{\perp}D^{\mathsf{b}}(gr_{\geq i} A)$. We can now apply Lemma [2.6.](#page-2-1) \Box

Remark 3.12. Let M be an object of $D^b(\text{gr }A)$ and P a projective resolution of M satisfying the conditions of Lemma [3.10.](#page-5-0) The proof shows that the localization triangle for M is given by

$$
P_{\prec i} \to P \simeq M \to P_{\succcurlyeq i} \to.
$$

Remark 3.13*.* Although we have chosen to work throughout with the grading group $\mathbb Z$, the results are valid more generally. One can replace $\mathbb Z$ by any totally ordered abelian group and work with graded rings concentrated in degrees greater than or equal to the identity.

This will also be the case for the majority of the results that follow. However, there are instances in which one does need additional hypotheses. For example in Lemma [6.5](#page-18-0) (and the main theorem [6.4\)](#page-17-0) one must assume the order admits successors.

4. Non-commutative Proj and local cohomology

For a graded non-commutative ring A, Artin and Zhang in [\[1\]](#page-22-4) defined the category of quasi-coherent sheaves on the non-commutative projective scheme $\text{Proj } A$ as the category of graded modules modulo the full subcategory of torsion modules (here and throughout torsion means torsion with respect to the two-sided ideal $A_{\geq 1}$). In this section we recall some of their definitions and results, in particular concerning local cohomology functors. When A is Gorenstein, these give a semiorthogonal decomposition of $D^{\mathsf{b}}(\mathrm{gr}_{\geq i} A)$ that is a key step in the proof of Orlov's theorem.

We assume $A = \bigoplus_{i \geq 0} A_i$ is a positively graded right noetherian ring. We consider Gr A , the abelian category of graded right A -modules. This contains gr A , the category of finitely generated graded A-modules, as a full abelian subcategory.

Definition 4.1. Let M be a graded A-module. An element $m \in M$ is *torsion* if

$$
m \cdot (A_{\geq n}) = 0
$$

for some $n \geq 1$. Denote by $\tau(M)$ the submodule of M consisting of all torsion elements. The module M is *torsion* if $\tau(M) = M$ and *torsion-free* if $\tau(M) = 0$. Denote by Tors A the full subcategory of Gr A consisting of torsion modules and set tors $A = Tors A \cap \text{gr } A$.

The subcategory $Tors A$ (respectively, $tors A$) satisfies the property that for a short exact sequence

$$
0 \to X' \to X \to X'' \to 0
$$

in Gr A (gr A), we have X in Tors A (tors A) if and only if X' and X'' are in Tors A $(\text{tors }A)$ i.e., they are Serre subcategories. Moreover, Tors A is closed under colimits. Thus we can form the quotient categories

$$
\text{Qcoh } X = \text{Gr } A / \text{Tors } A \qquad \text{and} \qquad \text{coh } X = \text{gr } A / \text{tors } A,
$$

see e.g. [\[19,](#page-23-9) §4.3] for the construction. The relevant features here are that:

- (1) the categories $Q \text{coh } X$ and $\text{coh } X$ have the same objects as $\text{Gr } A$ and $\text{gr } A$, respectively;
- (2) the categories $Q\text{coh } X$ and coh X are abelian and there are canonical exact functors $\operatorname{Gr} A \to \operatorname{Qcoh} X$ and $\operatorname{gr} A \to \operatorname{coh} X$;

(3) a map f in $\operatorname{Gr} A$ is an isomorphism in $\operatorname{Qcoh} X$ if and only if ker f and coker f are in Tors A; in particular the image of every object in Tors A is isomorphic to zero in $Q\text{coh } X$. The analogous statement holds for gr A.

For an object M in Gr A, we denote by \overline{M} the image of M in Qcoh X. For future reference, we note that as $Tors A$ is closed under the grading shifts, the shifts induce automorphisms of Qcoh X and coh X which we also denote by $(-)(i)$.

Remark 4.2. The notation Qcoh X and coh X reflects that these categories should be thought of as sheaves of modules on the noncommutative projective scheme $X = \text{Proj } A$. If A is commutative and generated in degree 1, then by a famous result of Serre, the category $Q\text{coh } X$ (respectively coh X) is equivalent to the category of quasi-coherent (respectively coherent) sheaves on the scheme $X = \text{Proj } A$. If A is generated in higher degrees, then $\cosh X$ is equivalent to the category of coherent sheaves on the Deligne-Mumford stack ProjA.

Definition 4.3. For M, N in Gr A, denote by $\underline{\text{Hom}}_{\text{Gr }A}(M, N)$ the graded abelian group

$$
\underline{\mathrm{Hom}}_{\mathrm{Gr}\,A}(M,N)=\bigoplus_{i\in\mathbb{Z}}\mathrm{Hom}_{\mathrm{Gr}\,A}(M(-i),N)\cong\bigoplus_{i\in\mathbb{Z}}\mathrm{Hom}_{\mathrm{Gr}\,A}(M,N(i)).
$$

If M is an A-A-bimodule, e.g. $M = A$, then $\underline{Hom}_{\text{Gr }A}(M, N)$ is a graded right A-module and so is in Gr A.

For any integer $p \geq 0$, we have a short exact sequence of A-bimodules:

$$
0 \to A_{\geq p} \to A \to A/A_{\geq p} \to 0.
$$

Applying $\underline{\text{Hom}}_{G_r A}(-, -)$, we have an exact sequence of endofunctors on Gr A:

$$
0 \to \underline{\text{Hom}}_{\text{Gr }A}(A/A_{\geq p}, -) \to \underline{\text{Hom}}_{\text{Gr }A}(A, -) \to \underline{\text{Hom}}_{\text{Gr }A}(A_{\geq p}, -).
$$

We may take the colimit of these sequences as $p \to \infty$ to get another exact sequence of functors; the sequence remains exact as both the abelian structure and colimits for endofunctors are inherited value-wise from Gr A and Gr A has exact filtered colimits. Note that for any M in $\mathbb{G}r A$ we have isomorphisms in $\mathbb{G}r A$:

$$
\underset{p\to\infty}{\text{colim }} \underline{\text{Hom}}_{\text{Gr }A}(A/A_{\geq p}, M) \cong \tau(M) \text{ and } \underline{\text{Hom}}_{\text{Gr }A}(A, M) \cong M.
$$

This gives a functorial exact sequence

(4.4)
$$
0 \to \tau(M) \to M \to \underset{p \to \infty}{\text{colim }} \underline{\text{Hom}}_{\text{Gr }A}(A_{\geq p}, M).
$$

Proposition 4.5. *The inclusion of* TorsA *into* Gr A *and the corresponding quotient functor have right adjoints* $\tau(-)$ *and* Γ_* *, respectively:*

$$
\operatorname{Tors} A \xrightarrow[\tau(-)]{} \operatorname{Gr} A \xrightarrow[\tau_{*}]{\widetilde{(-)}} \operatorname{Qcoh} X
$$

where for M *in* Gr A, $\tau(M)$ *is the torsion submodule of* M *and*

$$
\Gamma_*(M) = \underset{p \to \infty}{\text{colim}} \underline{\text{Hom}}_{\text{Gr }A}(A_{\geq p}, M).
$$

The functor Γ[∗] *and the inclusion of* TorsA *are fully faithful so the corresponding counit and unit respectively are isomorphisms. The remaining counit and unit are given by* [\(4.4\)](#page-8-0)*.*

Proof. It is easy to see that $\tau(-)$ is a right adjoint to the inclusion of Tors $A \rightarrow$ Gr A and it follows from abstract nonsense, see [\[19,](#page-23-9) $\S 4.4$], that there exists a right adjoint $\Gamma_*: \text{Qcoh } X \to \text{Gr } A$. We give here a direct proof that the functor colim $\underline{\text{Hom}}_{\text{Gr }A}(A_{\geq p},-)$ induces a right adjoint, and that the unit is given by [\(4.4\)](#page-8-0).

We first prove two claims:

Claim 1: If M is in Tors A, then $\operatorname{colim}_{p\to\infty} \underline{\text{Hom}}_{\text{Gr }A}(A_{\geq p}, M) = 0.$

Proof of claim: To see this, let ϕ be an element of $\underline{\text{Hom}}_{\text{Gr }A}(A_{\geq p}, M)$, for some $p \geq 0$. As A is right noetherian, $A_{\geq p}$ is finitely generated as a right ideal by some x_1, \ldots, x_k . We can find $m \geq 0$ so that $\phi(x_i) \cdot A_{\geq m} = 0$ for all i, using that M is torsion. Since $\phi(x_i \cdot A_{\geq m}) = \phi(x_i) \cdot A_{\geq m} = 0$ and, picking m larger if necessary, $A_{\geq m+p} = (x_1, \ldots, x_k)A_{\geq m}$, we have that $\phi|_{A_{\geq m+p}} = 0$ and so $\phi = 0$ in $\operatorname*{colim}_{p\rightarrow\infty}\underline{\mathrm{Hom}}_{\mathrm{Gr}\,A}(A_{\geq p},M).$

Claim 2: There are no non-zero morphisms from torsion modules to modules in the image of colim $\underline{\text{Hom}}_{\text{Gr }A}(A_{\geq p}, -)$.

Proof of claim: Let T be in Tors A and let $g: T \to \text{colim}_{\mathbb{P} \to \infty} \underline{\text{Hom}}_{\text{Gr }A}(A_{\geq p}, N)$ be a map. For $x \in T$, let $\phi \in \text{Hom}_{\text{Gr }A}(A_{\geq p}, N)$ be a representative of $g(x)$, for some p. Pick m such that $x \cdot A_{\geq m} = 0$. We have that $g(x) \cdot A_{\geq m} = g(x \cdot A_{\geq m}) = 0$, and that $\phi \cdot A_{\geq m}$ represents $g(x) \cdot A_{\geq m}$. However, picking a larger m if necessary, we see $\phi \cdot A_{\geq m}$ is the image of ϕ under the map $\underline{\text{Hom}}_{\text{Gr }A}(A_{\geq p}, N) \to \underline{\text{Hom}}_{\text{Gr }A}(A_{\geq m+p}, N)$ and so $\phi = 0$ in colim $\underline{\text{Hom}}_{\text{Gr }A}(A/A_{\geq p}, N)$, i.e. $g(x) = 0$.

To see that $\operatorname{colim}_{p\to\infty} \underline{\text{Hom}}_{\text{Gr }A}(A_{\geq p},-)$ induces a functor $\Gamma_*: \operatorname{Qcoh} X \to \operatorname{Gr }A$ it is enough to show that $\operatorname*{colim}_{p\to\infty} \underline{\text{Hom}}_{\text{Gr }A}(A/A_{\geq p},-)$ takes morphisms f with ker f and coker f in Tors A to invertible morphisms. This follows from Claims 1 and 2, and two applications of the snake lemma. To show that Γ_* is right adjoint to the quotient and [\(4.4\)](#page-8-0) is the unit, it is enough to show that any map $f : M \to \Gamma_*(N)$ factors through $M \to \Gamma_*(M)$. Note that by construction, we may extend [\(4.4\)](#page-8-0) to an exact sequence

(4.6)
$$
0 \to \operatorname{colim}_{p \to \infty} \underline{\text{Hom}}_{\text{Gr }A}(A/A_{\geq p}, M) \to M \to
$$

\n $\operatorname{colim}_{p \to \infty} \underline{\text{Hom}}_{\text{Gr }A}(A_{\geq p}, M) \to \operatorname{colim}_{p \to \infty} \underline{\text{Ext}}_{\text{Gr }A}^1(A/A_{\geq p}, M) \to 0.$

Since $\underline{\text{Hom}}_{\text{Gr }A}(A/A_{\geq p},-) \cdot A_{\geq p} = 0$, subobjects and quotients of torsion modules are torsion, and the colimit of torsion modules is torsion, we see that the last term $\operatorname{colim}_{p\to\infty} \underline{\operatorname{Ext}}^1_{\operatorname{Gr}A}(A/A_{\geq p},M)$ is in Tors A. To see that the map $f: M \to \Gamma_*(\widetilde{N}) =$ colim $\underline{\text{Hom}}_{\text{Gr }A}(A_{\geq p}, N)$ factors through $M \to \operatorname{colim}_{p\to\infty} \underline{\text{Hom}}_{\text{Gr }A}(A_{\geq p}, M)$, by [\[19,](#page-23-9) 4.1], it is enough to show that there are no non-zero morphisms from torsion modules to modules in the image of Γ_* , which was shown in Claim 2.

We now show Γ_* is fully faithful. Let $\eta_M : M \to \Gamma_* \tilde{M}$ be the unit of the adjunction, which is the center arrow of [\(4.6\)](#page-9-0). Since the outer two terms of that sequence are torsion, it follows that $\widetilde{\eta_M}$ is an isomorphism. Let $\epsilon_{\widetilde{M}} : \widetilde{(\Gamma_*\widetilde{M})} \to \widetilde{M}$

be the counit of the adjunction. By definition, the composition

$$
\widetilde{M} \xrightarrow{\widetilde{\eta_M}} \widetilde{\left(\Gamma_*\widetilde{M}\right)} \xrightarrow{\epsilon_{\widetilde{M}}} \widetilde{M}
$$

is the identity. Thus $\epsilon_{\widetilde{M}}$ is an isomorphism and so Γ_* is fully faithful.

Remark 4.7*.* As the notation suggests, if A is a commutative ring generated in degree 1, then $\Gamma_*(\tilde{\to})$ is isomorphic to $\bigoplus_{i\in\mathbb{Z}}\Gamma(\text{Proj }A,\widetilde{(-)}(i))$ as they are both right adjoint to sheafification Gr $A \to \text{Qcoh } \bar{X}$.

It is clear from the definition that the functor $\tau(-)$ takes gr A to tors A. However, Γ_* does not necessarily take objects of coh X to gr A:

Example 4.8. Let $A = k[x]$ with k a field, graded by $|x| = 1$. The A-module structure on $\operatorname{colim}_{p\to\infty} \frac{\operatorname{Ext}^1_{\mathbf{Gr}}_A(A/A_{\geq p}, A)$ is easily computed: it has a k-basis e_1, \ldots, e_n, \ldots with $|e_i| = -i$ and $xe_i = e_{i-1}$. In particular it is not finitely generated over A and so from [\(4.6\)](#page-9-0) we see that $\Gamma_*(\widetilde{A})$ is not either.

In the example above, $(\Gamma_*(A))_{\geq i}$ is finitely generated (in fact of finite length) for any $i \in \mathbb{Z}$. Artin and Zhang gave a criterion for A-modules that is equivalent to this fact being true. It is often easy to check. For instance, it holds for all modules over commutative rings.

Definition 4.9 (Artin, Zhang). An object M in gr A satisfies $\chi_j(M)$ if there exists an integer n_0 such that $\underline{\operatorname{Ext}}_{\operatorname{Gr} A}^k(A/A_{\geq n},M)_{\geq i}$ is a finitely generated A-module for all $i \in \mathbb{Z}$, $k \leq j$ and all $n \geq n_0$. The ring satisfies condition χ_j if $\chi_j(M)$ holds for all $M \in \operatorname{gr} A$.

If M satisfies $\chi_1(M)$, then [\[1,](#page-22-4) 3.8.3] shows that $\text{colim}_{p\to\infty} \frac{\text{Ext}_{\text{Gr }A}^1(A/A_{\geq p}, M)_{\geq i}}{p\to\infty}$ is a finitely generated A-module for all $i \in \mathbb{Z}$. Thus by [\(4.6\)](#page-9-0), we see that $\Gamma_{\geq i}(\widetilde{M}) :=$ $(\Gamma_*(M))_{\geq i}$ is finitely generated.

Remark 4.10*.* As [\[1,](#page-22-4) 3.1.4] shows, if A is commutative, then every module M satisfies $\chi_j(M)$. Indeed, we can compute the A-module $\mathrm{Ext}^j_{\mathrm{gr}\, A}(A/A_{\geq p}, M)$ using a graded free resolution of $A/A_{\geq p}$, which we can assume to be finite in each degree. If A is not commutative then we must use the bimodule structure on $A/A_{\geq p}$ to compute the A-module structure on $\text{Ext}^j_{\text{gr }A}(A/A_{\geq p}, M)$, i.e. in this case we must look at the derived functor of $\text{Hom}_{gr\,A}(A)A_{\geq p},-)$ (rather than deriving in the first variable) and so we cannot necessarily use a free resolution of $A/A_{\geq p}$ to compute the A-module structure of $\text{Ext}^j_{\text{gr }A}(A/A_{\geq p}, M)$. In [\[22\]](#page-23-10), an example is given of a non-commutative graded noetherian domain A such that $\chi_j(A)$ does not hold for any $j > 0$.

Recall that $gr_{\geq i} A$ is the full subcategory of gr A with objects those M with $M =$ $M_{\geq i}$. We denote by $\operatorname{Gr}_{\geq i} A$ the analogous subcategory of $\operatorname{Gr} A$. Let $\operatorname{Tors}_{\geq i} A =$ $Gr_{\geq i} A \cap Tors A$ and $tors_{\geq i} A = gr_{\geq i} A \cap tors A$. The functor $\tau(-)$ restricted to Gr $\geq_i A$ (respectively, $\operatorname{gr}_{\geq i} A$) is a right adjoint of the inclusion $\operatorname{Tors}_{\geq i} A \to$ $Gr_{\geq i} A$ (respectively, $tors_{\geq i} A \to gr_{\geq i} A$). Moreover, it is easy to check that the composition of the functors $\text{Gr}_{\geq i} A \to \text{Gr} A \to \text{Qcoh} X$ induces an equivalence $\text{Gr}_{\geq i} A/\text{Tors}_{\geq i} A \stackrel{\cong}{\to} \text{Qcoh } X \text{ and } \Gamma_{\geq i} = (\Gamma_*(-))_{\geq i}$ is a right adjoint to the quotient map. There is also an equivalence $gr_{\geq i} A/\operatorname{tors}_{\geq i} A \stackrel{\cong}{\to} \coth X$.

Assume that A satisfies the condition χ_1 . Then, using the above, we have the following diagram where the vertical arrows are inclusions and the horizontal arrows form adjoint pairs with the left adjoint on top:

(4.11)
$$
\operatorname{Tors}_{\geq i} A \xrightarrow{\operatorname{inc}} \operatorname{Gr}_{\geq i} A \xrightarrow{\overbrace{(\overline{-})}^{\operatorname{inc}}} \operatorname{Qcoh} X
$$

$$
\operatorname{tors}_{\geq i} A \xrightarrow{\operatorname{inc}} \operatorname{gr}_{\geq i} A \xrightarrow{\overbrace{(\overline{-})}^{\operatorname{fc}}}^{\operatorname{inc}} \operatorname{coh} X
$$

$$
\operatorname{tors}_{\geq i} A \xrightarrow{\operatorname{inc}} \operatorname{gr}_{\geq i} A \xrightarrow{\overbrace{\overline{(\overline{-})}^{\operatorname{fc}}}^{\operatorname{fc}}} \operatorname{coh} X
$$

For any M in Gr $>i$ A, one counit and one unit are isomorphisms and the other two are given by

(4.12)
$$
0 \to \tau(M) \to M \cong \underline{\text{Hom}}_{\text{Gr }A}(A, M) \to (\underset{p \to \infty}{\text{colim }} \underline{\text{Hom}}_{\text{Gr }A}(A_{\geq p}, M))_{\geq i}
$$

as in the case of Gr A. Also note that $\Gamma_{\geq i}$ is fully faithful, as it is the right adjoint of a quotient functor.

We wish to extend this diagram to functors between the bounded derived categories of the above abelian categories. The existence of a corresponding localization sequence involving the derived categories is standard but we provide some details. We start with a simple lemma.

Lemma 4.13. Let M be an object of $\text{Gr}_{\geq i} A$ and let $M \to I$ be an injective reso*lution in* Gr A. Then $M \to I_{\geq i}$ *is an injective resolution in* Gr $\geq_i A$. In particular $Gr_{\geq i}$ A has enough injectives.

Proof. The functor $(-)_{\geq i}$ is exact and $M_{\geq i} = M$, thus $M \to I_{\geq i}$ is a quasiisomorphism. So to complete the proof it is sufficient to show $I_{\geq i}$ is a complex of injectives.

Let J be an injective object in $\mathbb{G}r A$. By adjunction there is an isomorphism of functors $\text{Hom}_{\text{Gr }A}(\text{inc}(-), J) \cong \text{Hom}_{\text{Gr }_{\geq i}A}(-, J_{\geq i})$. The former functor is exact as J is injective and the inclusion is exact, and thus so is the latter showing $J_{\geq i}$ is injective in $\mathrm{Gr}_{\geq i} A$.

Thus $(-)_{\geq i}$ preserves injectives and so the quasi-isomorphism $M \to I_{\geq i}$ is an injective resolution.

The functors to the right in [\(4.11\)](#page-11-0) are exact and those to the left are left exact (since they are right adjoints). Since $\text{Gr}_{\geq i} A$ has enough injectives by the above lemma and $Q\text{coh } X$ has enough injectives by $[1, 7.1]$ (in fact, by standard abstract nonsense both of these categories are Grothendieck categories and so have enough injectives), we may form $\mathbf{R}\tau(-)$ and $\mathbf{R}\Gamma_{\geq i}$, the right derived functors of $\tau(-)$ and $\Gamma_{\geq i}$, respectively. This gives two pairs of adjoint functors

$$
(4.14) \qquad D_{Tors_{\geq i}A}(Gr_{\geq i}A) \xrightarrow{\text{inc}} D(Gr_{\geq i}A) \xrightarrow{\widetilde{(-)}} D(\text{Qcoh } X)
$$

where $D_{Tors_{\geq i} A}(Gr_{\geq i} A)$ is the full subcategory of $D(Gr_{\geq i} A)$ consisting of complexes with torsion cohomology.

Since $\Gamma_{\geq i}$ sends injectives to injectives and is fully faithful one checks easily that $\mathbf{R}\Gamma_{\geq i}$ is also fully faithful. In particular, we have that $(-)$ is a quotient functor. As $(-)$ at the level of the abelian categories is exact, the kernel of this functor at the level of derived categories consists of precisely those complexes whose cohomology is annihilated by (-) i.e., it is exactly $D_{Tors_{\geq i}} A(\text{Gr}_{\geq i} A)$. This proves the above functors give a localization sequence of triangulated categories.

It follows that for every $M \in D(\text{Gr}_{\geq i} A)$ there is a localization triangle

(4.15)
$$
\mathbf{R}\tau(M) \to M \to \mathbf{R}\Gamma_{\geq i}(M) \to.
$$

where the first map is the counit of the first adjunction of (4.14) and the second map is the unit of the second adjunction of [\(4.14\)](#page-11-1).

Remark. Note that when A is commutative, a triangle such as [4.15](#page-12-0) can be explicitly constructed using the Cech complex.

For the above adjoint pairs to restrict to the bounded derived categories of complexes of finitely generated modules, we need to place two further restrictions on A. Let $\mathbb{R}\Gamma_*\colon D(\mathrm{Qcoh}\,X)\to D(\mathrm{Gr}\,A)$ be the right derived functor of the left exact functor Γ∗.

Definition 4.16 (Artin-Zhang). The *cohomological dimension* of A is

$$
\operatorname{cd}(A) := \sup \{ d \mid H^d \mathbf{R} \Gamma_*(\widetilde{A}) \neq 0 \}.
$$

By [\[1,](#page-22-4) 7.10], if cd(A) is finite, then $\mathbb{R}\Gamma_*(\widetilde{M})$ is a bounded complex for every $\widetilde{M} \in$ $\operatorname{Qcoh} X$ and so restricts to a functor

$$
\mathbf{R}\Gamma_*: \mathsf{D}^{\mathsf{b}}(\operatorname{Qcoh} X) \to \mathsf{D}^{\mathsf{b}}(\operatorname{Gr} A).
$$

Since $\Gamma_{\geq i}$ is the composition of Γ_* and the exact functor $(-)_{\geq i}$, we see that $\mathbf{R}\Gamma_{\geq i}$ $(R\Gamma_*)_{\geq i}$ and so $R\Gamma_{\geq i}$ restricts to a functor

$$
\mathbf{R}\Gamma_{\geq i} : \mathsf{D}^{\mathrm{b}}(\mathrm{Qcoh}\, X) \to \mathsf{D}^{\mathrm{b}}(\mathrm{Gr}_{\geq i}\, A).
$$

By the long exact sequence in homology induced by [\(4.15\)](#page-12-0), we see that $\mathbf{R}\tau$ also restricts to a functor between bounded derived categories.

Now we consider finiteness. We want to compute the cohomology of $\mathbf{R}\tau(M)$. We view $\tau(-) = \operatorname*{colim}_{p \to \infty} \underline{\text{Hom}}_{\text{Gr }A}(A/A_{\geq p}, -)$ as a functor from $\operatorname{gr}_{\geq i} A \to \operatorname{tors}_{\geq i} A$. For M in $gr_{\geq i} A$, let $M \to I_M$ be an injective resolution in Gr A. Then $(I_M)_{\geq i}$ is an injective resolution in $\mathrm{Gr}_{\geq i} A$. Thus we have

$$
\mathbf{R}\tau(M) = \operatorname*{colim}_{p \to \infty} \underline{\operatorname{Hom}}_{\operatorname{Gr} A}(A/A_{\geq p}, (I_M)_{\geq i}) = \operatorname*{colim}_{p \to \infty} \underline{\operatorname{Hom}}_{\operatorname{Gr} A}(A/A_{\geq p}, I_M)_{\geq i}
$$

where the second equality follows from the commutativity of the square of inclusions

$$
\operatorname{Tors}_{\geq i} A \longrightarrow \operatorname{Gr}_{\geq i} A
$$
\n
$$
\downarrow \qquad \qquad \downarrow
$$
\n
$$
\operatorname{Tors} A \longrightarrow \operatorname{Gr} A
$$

by taking right adjoints. This shows that

$$
H^k \mathbf{R} \tau(M) \cong \underset{p \to \infty}{\text{colim}} \underline{\operatorname{Ext}}_{\operatorname{Gr} A}^k (A/A_{\geq p}, M)_{\geq i}
$$

for all $k \geq 0$. By [\[1,](#page-22-4) 3.8.3], if M satisfies $\chi_j(M)$, then $\text{colim}_{p\to\infty} \underline{\text{Ext}}_{\text{gr }A}^k(A/A_{\geq p}, M)_{\geq i}$ is a finitely generated A-module for all $k \leq j$ and all $p \in \mathbb{Z}$.

Assume now that A has finite cohomological dimension and satisfies χ_i for all $j \geq 0$. The above shows that **R** τ (−) restricts to a functor

$$
\mathbf{R}\tau(-):{\mathsf D}^{\mathsf{b}}({\mathsf{gr}}_{\geq i}A)\to{\mathsf D}^{\mathsf{b}}_{\operatorname{tors}_{\geq i}A}({\mathsf{gr}}_{\geq i}A).
$$

By the long exact sequence in cohomology coming from [\(4.15\)](#page-12-0), we see that we also have a functor

$$
\mathbf{R}\Gamma_{\geq i} : \mathsf{D}^{\mathsf{b}}(\operatorname{coh} X) \to \mathsf{D}^{\mathsf{b}}(\operatorname{gr}_{\geq i} A).
$$

This gives the following diagram of adjoint functors, where the top functors are the left adjoints:

$$
\mathsf{D}_{\mathsf{tors}_{\geq i}^{\mathsf{b}}\,A}^{\mathsf{b}}(\mathsf{gr}_{\geq i}A) \xrightarrow{\mathsf{inc}} \mathsf{D}^{\mathsf{b}}(\mathsf{gr}_{\geq i}A) \xrightarrow{\widetilde{(-)}} \mathsf{D}^{\mathsf{b}}(\mathsf{coh}\,X) .
$$

The functor $\mathbf{R}\Gamma_{\geq i}$ is fully faithful and its image is left admissible. Also, any object in this image is contained in $\left(\mathsf{D}^{\mathsf{b}}_{\text{tors}_{\geq i}} A(\text{gr}_{\geq i} A)\right)^{\perp}$. Indeed, for M an object with torsion cohomology and any $N \in D^{\mathsf{b}}(\mathrm{gr}_{\geq i} A)$, we have that

$$
\operatorname{Hom}\nolimits_{\mathsf{D}^{\mathrm{b}}(\operatorname{gr}\nolimits_{\geq i}A)}(M,\mathbf{R}\Gamma_{\geq i}\widetilde{N})\cong \operatorname{Hom}\nolimits_{\mathsf{D}^{\mathrm{b}}(\operatorname{coh}\nolimits X)}(\widetilde{M},\widetilde{N})=0
$$

since $\widetilde{M} \simeq 0$. From this containment and the triangle [\(4.15\)](#page-12-0), we may apply [2.6](#page-2-1) to see that there is a semiorthogonal decomposition

$$
\mathsf{D}^{\mathsf{b}}(\operatorname{gr}_{\geq i}A)=\left(\mathbf{R}\Gamma_{\geq i}\mathsf{D}^{\mathsf{b}}(\operatorname{coh} X),\mathsf{D}^{\mathsf{b}}_{\operatorname{tors}_{\geq i}A}(\operatorname{gr}_{\geq i}A)\right).
$$

Recall that $S_{\geq i}$ is the thick subcategory generated by $A_0(e)$ for all $e \leq -i$.

Lemma 4.17. *There is an equality* $S_{\geq i} = D^{\mathsf{b}}_{\text{tors}_{\geq i}} A(\text{gr}_{\geq i} A)$.

Proof. It's clear that $A_0(e)$ is in tors_{≥i} A for all $e \leq -i$, so $S_{\geq i}$ is contained in $D^{\mathsf{b}}_{\text{tors}_{\geq i}} A(\text{gr}_{\geq i} A)$. Given M in $D^{\mathsf{b}}_{\text{tors}_{\geq i}} A(\text{gr}_{\geq i} A)$, we have that $H^*(M)$ is finitely generated and torsion, thus M must have cohomology in only finitely many degrees. Analogously to the proof of [3.2,](#page-3-0) this shows that M is in $S_{\geq i}$. .

The above shows the following:

Proposition 4.18. *Let* A *be a positively graded right noetherian ring that satisfies condition* χ *and has finite cohomological dimension. Then there is a semiorthogonal decomposition*

$$
\mathsf{D}^{\mathsf{b}}(\mathrm{gr}_{\geq i}\,A)=\left(\mathbf{R}\Gamma_{\geq i}\mathsf{D}^{\mathsf{b}}(\mathrm{coh}\,X),\mathcal{S}_{\geq i}\right).
$$

The corresponding localization triangle is given by [\(4.15\)](#page-12-0)*.*

5. Singularity category of a Gorenstein ring

In this section we assume that $A = \bigoplus_{i \geq 0} A_i$ is a positively graded (two-sided) noetherian ring with A_0 of finite global dimension, but not necessarily commutative.

In the following, we denote by $\mathrm{id}_A M$ the graded injective dimension of a graded A-module M.

Definition 5.1. The ring A is *(Artin-Schelter) Gorenstein* if $id_A A < \infty$, $id_{A^{\text{op}}} A <$ ∞ and

 $\mathbf{R} \underline{\text{Hom}}_{gr\,A}(A_0, A) \cong A_0[n](a)$ for some $n, a \in \mathbb{Z}$

in both D^b (gr A) and D^b (gr A^{op}). The unique integer a is the a-invariant of A.

Remark 5.2*.* In [\[16\]](#page-23-11) a different definition of Artin-Schelter Gorenstein ring is given under the restriction that A_0 is a finite dimensional algebra over a fixed base field k. Their definition differs from ours in two ways: Minamoto and Mori require the shift occurring to match the injective dimension of A i.e., $n = -id_A A$, and that rather than $\mathbf{R} \underline{\text{Hom}}_{gr\,A}(A_0, A) \cong A_0[n](a)$ one asks for an isomorphism

$$
\mathbf{R}\underline{\mathrm{Hom}}_{\mathrm{gr}\,A}(A_0,A)\cong \mathrm{Hom}_k(A_0,k)[n](a).
$$

We note both definitions restrict to the classical one in the case $A_0 = k$.

As an example, if R is a commutative regular ring of positive Krull dimension and we set $A = R[x]/(x^n)$ with x in degree 1 then A is AS-Gorenstein in our sense but not according to [\[16\]](#page-23-11). On the other hand the definition of Minamoto-Mori covers certain (higher) preprojective algebras which are in general excluded by our definition.

From this point forward we will use the term *Gorenstein ring* to refer to a ring that is Gorenstein either in the sense of Definition [5.1](#page-13-1) or [\[16\]](#page-23-11). Our results hold for both definitions. We will work with the definition we give and, when necessary, point out what changes in the arguments are necessary if one uses the definition of Minamoto and Mori. In fact, the only place in which the arguments do not go through verbatim are Lemmas [6.2](#page-17-1) and [6.6](#page-18-1) which require minor tweaking.

The most important feature of Gorenstein rings for us is the duality given below. We will make a standard abuse of notation and not differentiate between the two duality functors notationally.

Lemma 5.3. *Assume that* A *is a Gorenstein ring. Then the functors*

$$
D = \mathbf{R} \underline{\mathrm{Hom}}_{\mathrm{gr}\,A}(-, A) : \mathsf{D}^{\mathsf{b}}(\mathrm{gr}\,A) \to \mathsf{D}^{\mathsf{b}}(\mathrm{gr}\,A^{op})^{op}
$$

$$
D = \mathbf{R} \underline{\mathrm{Hom}}_{\mathrm{gr}\,A^{op}}(-, A) : \mathsf{D}^{\mathsf{b}}(\mathrm{gr}\,A^{op}) \to \mathsf{D}^{\mathsf{b}}(\mathrm{gr}\,A)^{op}
$$

are quasi-inverse equivalences.

Proof. We first observe that D does indeed take $D^b(\text{gr }A)$ to $D^b(\text{gr }A^{\text{op}})^\text{op}$. Since A has finite injective dimension as both a left and a right module over itself it is clear that D preserves boundedness of cohomology. It is also clear that D sends complexes with finitely generated cohomology groups to the same as we can resolve any object of $D^b(\text{gr }A)$ by a complex of finitely generated projectives and A is noetherian.

These functors are adjoint so we can consider the unit of this adjunction

$$
\eta\colon\text{Id}\to D^2
$$

and we need to show it is an equivalence. But this is again clear: for a bounded above complex of finitely generated projectives the map η is just componentwise the natural map to the double dual and finitely generated projectives are reflexive. \Box

Recall that $\mathcal{P}_{\geq i}$ is the thick subcategory of $\mathsf{D}^{\mathsf{b}}(\text{gr }A)$ generated by those $A(e)$ with $e \leq -i$ and $\mathcal{P}_{\leq i}$ is the thick subcategory generated by the $A(e)$ with $e > -i$.

Lemma 5.4. *If* A *is Gorenstein, there is a semiorthogonal decomposition*

$$
\mathsf{D}^{\mathsf{b}}(\operatorname{gr}_{\geq i} A) = \left(\mathcal{P}_{\geq i}, \left({}^{\perp}\mathcal{P}_{\geq i}\right) \cap \mathsf{D}^{\mathsf{b}}(\operatorname{gr}_{\geq i} A)\right).
$$

 $For M \in D^b(\text{gr}_{\geq i} A)$ *, the localization triangle is given by*

$$
D(G)_{\prec i} \to M \cong D(G) \to D(G)_{\succcurlyeq i} \to
$$

where the notation is as in Definition [3.9,](#page-5-1) and $G \to D(M)$ *is a projective resolution of the dual of* M *as in Lemma [3.10.](#page-5-0)*

Proof. Let M be an object of $D^b(\mathrm{gr}_{\geq i} A)$ and let $G \to D(M)$ be a projective resolution as in Lemma [3.10,](#page-5-0) where $D(M) = \mathbf{R} \underline{Hom}_{grA}(M, A)$ is the image of M under the duality functor. As in the proof of Lemma [3.7,](#page-4-0) there is a triangle

$$
G_{\prec-i+1} \to G \to G_{\succcurlyeq-i+1} \to
$$

where $G_{\prec-i+1}$ is an object of $\mathcal{P}_{\prec-i+1}$ and every component of $G_{\succ_{i+1}}$ is generated in degree at least $-i + 1$. If $P = P_0 \otimes_{A_0} A(e)$ is any indecomposable graded projective A-module, then $D(P) \cong \underline{\text{Hom}}_{gr\,A}(P, A) \cong (P_0)^* \otimes_{A_0} A(-e)$, where $(P_0)^*$ is the A_0 -dual of P_0 . Thus $D(G_{\prec-i+1}) = D(G)_{\succ i}$ and $D(G_{\succ i+1}) = D(G)_{\prec i}$. Applying D to the triangle above gives a triangle

$$
D(G)_{\prec i} \to D(G) \to D(G)_{\succcurlyeq i} \to.
$$

Note that $D(G)_{\geq i}$ is in $\mathcal{P}_{\geq i}$ and that there are isomorphisms $M \xrightarrow{\simeq} D(D(M)) \xrightarrow{\simeq} D(D(M))$ D(G). We can now apply Lemma [2.6,](#page-2-1) once we show that $D(G)_{\prec i}$ is in $({}^{\perp}P_{\geq i})$ $D^{b}(gr_{\geq i} A)$. It follows from the long exact sequence in homology of the above triangle that each of the homology groups of $D(G)_{\prec i}$ is generated in degrees at least i and thus $D(G)_{\prec i}$ is in $D^{\mathsf{b}}(\mathrm{gr}_{\geq i} A)$. That $D(G)_{\prec i}$ is in $\perp(\mathcal{P}_{\geq i})$ follows from the fact that $\text{Hom}_{\text{gr }A}(A(e), A(f)) = 0$ for $e > f$.

Let us denote by \mathcal{B}_i the subcategory $({}^{\perp}\mathcal{P}_{\geq i}) \cap D^{\mathsf{b}}(\text{gr}_{\geq i} A)$, which by the above lemma is a right admissible subcategory of $D^{\mathsf{b}}(\mathrm{gr}_{\geq i} A)$. There is a description of \mathcal{B}_i using the well-known singularity category of A.

Definition 5.5. Let A be a graded ring.

- (1) An object M in D^b (gr A) is *perfect* if M is in the thick subcategory generated by $A(e)$ for all $e \in \mathbb{Z}$ i.e., it is quasi-isomorphic to a bounded complex of projectives. We denote the subcategory of perfect complexes by $\text{perf}(A)$. We see from the definitions that $\text{perf}(A) = \langle \mathcal{P}_{\geq i}, \mathcal{P}_{\leq i} \rangle$.
- (2) The *singularity category* of A is

$$
\mathsf{D}^{\mathsf{b}}_{\mathsf{sg}}(\operatorname{gr} A) := \mathsf{D}^{\mathsf{b}}(\operatorname{gr} A) / \operatorname{perf}(A).
$$

Orlov showed that when A is a connected graded Gorenstein algebra over a field, there is an embedding of $D_{sg}^{b}(gr A)$ in $D^{b}(gr A)$ for every $i \in \mathbb{Z}$, and the image is equal to \mathcal{B}_i . We now show this holds in the generality in which we are working. First we recall a lemma whose proof is left to the reader.

Lemma 5.6. Let A be a left admissible subcategory in a triangulated category T *with* $i_L : \mathcal{T} \to \mathcal{A}$ *the left adjoint to the inclusion* $i : \mathcal{A} \to \mathcal{T}$ *. Then* i_L *induces an equivalence*

$$
\mathcal{T}/{}^{\perp}\mathcal{A}\to\mathcal{A}
$$

with inverse equivalence the composition $A \to \mathcal{T} \to \mathcal{T}/\mathcal{L} A$. The analogous state*ment holds for right admissible subcategories.*

Applying the above lemma to [3.7](#page-4-0) shows that there is an equivalence

$$
\psi_i \colon \mathsf{D}^{\mathsf{b}}(\operatorname{gr} A)/\mathcal{P}_{< i} \xrightarrow{\cong} \mathsf{D}^{\mathsf{b}}(\operatorname{gr}_{\geq i} A).
$$

Remark [3.12](#page-7-0) shows that $\psi_i(M) = P_{\geq i}$, where $P \to M$ is a projective resolution as in Lemma [3.10.](#page-5-0) If we apply Lemma [5.6](#page-15-0) again to the semiorthogonal decomposition $D^{\mathsf{b}}(\mathrm{gr}_{\geq i}, A) = (\mathcal{P}_{\geq i}, \mathcal{B}_i)$, we have an equivalence

$$
\phi_i \colon \mathsf{D}^{\mathsf{b}}(\mathsf{gr}_{\geq i} \, A)/\mathcal{P}_{\geq i} \xrightarrow{\cong} \mathcal{B}_i
$$

with $\phi_i(N) = D(Q)_{\prec i}$, where $Q \to D(N)$ is a projective resolution as in Lemma [3.10.](#page-5-0) Let us set $\mathbf{b}_i = \phi_i \circ \pi \circ \psi_i$ where $\pi \colon \mathsf{D}^{\mathsf{b}}(\mathrm{gr}_{\geq i} A) \to \mathsf{D}^{\mathsf{b}}(\mathrm{gr}_{\geq i} A)/\mathcal{P}_{\geq i}$ is the quotient functor. This gives an equivalence

(5.7)
$$
\mathbf{b}_i \colon \mathsf{D}^{\mathsf{b}}_{\mathsf{sg}}(\operatorname{gr} A) = \mathsf{D}^{\mathsf{b}}(\operatorname{gr} A) / \langle \mathcal{P}_{\leq i}, \mathcal{P}_{\geq i} \rangle \xrightarrow{\cong} \mathcal{B}_i
$$

with $\mathbf{b}_i(M) = D(Q)_{\prec i}$, where $Q \to D(P_{\succ i})$ and $P \to M$ are projective resolutions as in Lemma [3.10.](#page-5-0) The inverse of the equivalence is given by the composition of the inclusion and quotient $\mathcal{B}_i \to \mathsf{D}^{\mathsf{b}}(\text{gr } A) \to \mathsf{D}^{\mathsf{b}}(\text{gr } A)/\operatorname{perf}(A)$. Moreover, we have that \mathbf{b}_i followed by the inclusion $\mathcal{B}_i \to \mathsf{D}^{\mathsf{b}}(\mathrm{gr}_{\geq i} A)$ is left adjoint to the quotient functor $D^b(\operatorname{gr}_{\geq i} A) = D^b(\operatorname{gr} A)/\mathcal{P}_{\leq i} \to D^b(\operatorname{gr} A) / \langle \mathcal{P}_{\leq i}, \mathcal{P}_{\geq i} \rangle = D^b_{\text{sg}}(\operatorname{gr} A).$

To sum up, we have shown the following:

Proposition 5.8. *If* A *is a graded Gorenstein ring, the quotient* $D^b(g_{{\bf r}_{\geq i}} A) \rightarrow$ D b sg(gr A) *has a fully faithful left adjoint*

$$
b_i: \mathsf{D}^{\mathsf{b}}_{\mathsf{sg}}(\text{gr}\, A) \to \mathsf{D}^{\mathsf{b}}(\text{gr}_{\geq i} A).
$$

The image of \mathbf{b}_i *is the subcategory* $\mathcal{B}_i = (\perp \mathcal{P}_{\geq i}) \cap D^b(\mathrm{gr}_{\geq i} A)$ *and there is a semiorthogonal decomposition:*

$$
\mathsf{D}^{\mathsf{b}}(\mathrm{gr}_{\geq i} A) = (\mathcal{P}_{\geq i}, \mathcal{B}_i).
$$

The localization triangle is described in [5.4.](#page-14-0)

6. Relating the bounded derived category of coherent sheaves and THE SINGULARITY CATEGORY

In this section we prove the main theorem by comparing the semiorthogonal decompositions constructed in the previous sections. We assume that $A = \bigoplus_{i \geq 0} A_i$ is a positively graded noetherian Gorenstein ring with A_0 a ring of finite global dimension, but not necessarily commutative.

Gorenstein rings often satisfy the two properties we need to apply [4.18.](#page-13-2)

Lemma 6.1. *If* A *is a Gorenstein ring, then* A *has finite cohomological dimension.*

Proof. We need to show

$$
\operatorname{cd}(A) = \sup \{ d \mid H^d \mathbf{R} \Gamma_*(\widetilde{A}) \neq 0 \} < \infty.
$$

Since A is Gorenstein we can choose a bounded injective resolution I for A as a right A-module. Hence $\mathbf{R}\tau(A) = \tau(I)$ has bounded cohomology and the localization triangle

$$
\mathbf{R}\tau(A) \to A \to \mathbf{R}\Gamma_*(A) \to
$$

then implies $\mathbf{R}\Gamma_*(\widetilde{A})$ also has bounded cohomology.

We have remarked earlier that any commutative ring satisfies condition χ . The next lemma gives some noncommutative and not necessarily graded connected examples.

Lemma 6.2. *Let* k *be a commutative ring and* A *a flat Gorenstein* k*-algebra. Then* A *satisfies condition* χ*.*

Proof. As A is flat over k it follows that the enveloping algebra $A \otimes_k A^{op}$ is flat over both A and A^{op} . Thus the restriction of scalars functors induced by the maps $A \to A \otimes_k A^{op}$ and $A^{op} \to A \otimes_k A^{op}$ preserve injectives. Taking an injective resolution I of A over $A \otimes_k A^{op}$ thus gives a bimodule resolution of A which is an injective resolution as both a complex of left and of right A-modules.

We may use such a resolution to compute $D = \mathbf{R} \underline{\text{Hom}}_{\text{gr }A}(-, A)$ as $\underline{\text{Hom}}_{\text{gr }A}(-, I)$ and obtain the correct A^{op} -module structure and similarly for the inverse duality functor; this is just a consequence of the fact that I and A are quasi-isomorphic as complexes of bimodules. Given a complex of injectives $M \in D^b(\text{gr } A)$ we now compute, using the duality of Lemma [5.3,](#page-14-1) that there are quasi-isomorphisms of right A_0 -modules

$$
\underline{\operatorname{Hom}}_{{\operatorname{gr}\,} A}(A_0,M)\cong\operatorname{\mathbf{R}}\underline{\operatorname{Hom}}_{{\operatorname{gr}\,} A^{\operatorname{op}}}(\underline{\operatorname{Hom}}_{{\operatorname{gr}\,} A}(M,I),\underline{\operatorname{Hom}}_{{\operatorname{gr}\,} A}(A_0,I))\\\cong \underline{\operatorname{Hom}}_{{\operatorname{gr}\,} A^{\operatorname{op}}}(P,\underline{\operatorname{Hom}}_{{\operatorname{gr}\,} A}(A_0,I))\\\cong \underline{\operatorname{Hom}}_{{\operatorname{gr}\,} A^{\operatorname{op}}}(P,\nu A_0[n](a)),
$$

where P is a projective resolution of $\underline{\text{Hom}}_{gr\,A}(M,I)$ over A^{op} and ν is a twist by some, possibly non-trivial, automorphism which needs to be accounted for as we view $\underline{\text{Hom}}_{\text{gr }A}(A_0, I)$ as a bimodule rather than just a right module (see for example [\[16,](#page-23-11) Lemma 2.9]). Now $\underline{\text{Hom}}_{\text{gr }A^{\text{op}}}(P, \Sigma^n {\scriptstyle_{\nu}} A_0(a))$ is a complex of finitely generated A_0 -modules and so in particular has finitely generated cohomology over A_0 and hence over A. In particular, if M is an injective resolution of a right A module N this shows $\underline{\operatorname{Ext}}_{gr\,A}^i(A_0, N)$ is finitely generated over A for all $i \in \mathbb{Z}$.

It only remains to observe that $A/A_{\geq n}$ has a filtration, as bimodules, by copies of $A_0(j)$ for $j \in \mathbb{Z}$ and considering the corresponding long exact sequences shows $\mathbf{R} \underline{\text{Hom}}_{\text{gr }A}(A/A_{\geq n}, M)$ has finitely generated cohomology for all $M \in \mathsf{D}^{\mathsf{b}}(\text{gr }A)$. Hence A satisfies condition χ .

$$
\Box
$$

Remark 6.3*.* In the above lemma if A is AS-Gorenstein in the sense of [\[16\]](#page-23-11) then one has to replace $_{\nu}A_0(a)$ by $\text{Hom}_k({}_{\nu}A_0(a), k)$ but this does not alter the argument as $\underline{\text{Hom}}_{\text{gr }A^{\text{op}}}(P, \Sigma^n \text{Hom}_k(\nu A_0(a), k))$ is still a complex of finitely generated A_0 modules.

Theorem 6.4. Let $A = \bigoplus_{i \geq 0} A_i$ be a positively graded noetherian Gorenstein ring with A_0 of finite global dimension, but not necessarily commutative. We assume in *addition that* A *satisfies condition* χ*. Let* a *be the* a*-invariant of* A *defined in [5.1.](#page-13-1)*

(1) *If* $a > 0$ *, then for any* $i \in \mathbb{Z}$ *there is a semiorthogonal decomposition*

 $D^b(\operatorname{coh} X) = \left(\mathcal{O}(-i - a + 1), \ldots, \mathcal{O}(-i), \widetilde{\mathcal{B}}_i\right),$

where $\mathcal{O}(j)$ *is the image of* $A(j)$ *in* coh X *and* \mathcal{B}_i *is the image of* $D_{sg}^b(\text{gr }A)$ *under the fully faithful functor* $\mathbf{b}_i : \mathsf{D}^{\mathsf{b}}_{\mathsf{sg}}(\text{gr } A) \to \mathsf{D}^{\mathsf{b}}(\text{gr}_{\geq i} A)$ *described in* [\(5.7\)](#page-16-0)*.*

(2) If $a < 0$, then for any $i \in \mathbb{Z}$ there is a semiorthogonal decomposition

$$
D_{\text{sg}}^{b}(\text{gr }A) = (pA_0(-i), \ldots, pA_0(-i+a+1), pR\Gamma_{\geq i-a}D^{b}(\text{coh }X)),
$$

where $p : D^b(\mathrm{gr}_{\geq i} A) \to D^b_{sg}(\mathrm{gr} A)$ *is the canonical quotient.*

(3) If $a = 0$, then for any $i \in \mathbb{Z}$ the functors $\overline{(-)}b_i : D^b_{sg}(\text{gr } A) \to D^b(\text{coh } X)$ $and \ p\mathbf{R}\Gamma_{\geq i} : \mathsf{D}^{\mathsf{b}}(\operatorname{coh} X) \to \mathsf{D}^{\mathsf{b}}_{\mathsf{sg}}(\operatorname{gr} A)$ *are inverse equivalences.*

Before beginning the proof, we need two lemmas. For the rest of the section we rely heavily on notation introduced earlier: recall that $\mathcal{S}_{\leq i}$ (respectively $\mathcal{S}_{\geq i}$) is the thick subcategory generated by the objects $A_0(e)$, for all $e > -i$ (respectively $e \leq -i$) and $\mathcal{P}_{\leq i}$ (respectively $\mathcal{P}_{\geq i}$) is the thick subcategory generated by the objects $A(e)$ for all $e > -i$ (respectively $e \leq -i$).

Lemma 6.5. *Let* A *be a graded ring.*

(1) *For any* $i \in \mathbb{Z}$ *, there is a semiorthogonal decomposition*

$$
\mathcal{P}_{\geq i} = (\mathcal{P}_{\geq i+1}, A(-i)).
$$

(2) For any $i \in \mathbb{Z}$, there is a semiorthogonal decomposition

$$
\mathcal{S}_{\leq i+1} = (\mathcal{S}_{\leq i}, A_0(-i))\,.
$$

Proof. It is clear that $A(-i) \subseteq {}^{\perp}P_{\geq i+1}$. We know that any object in $P_{\geq i}$ is isomorphic in $\mathsf{D}^{\mathsf{b}}(\mathrm{gr}_{\geq i} A)$ to a bounded complex X of finitely generated graded projective modules and so we may restrict ourselves to working with such complexes. As in the proof of [3.7,](#page-4-0) using the structure of graded projectives given in [3.8](#page-4-1) and the notation of Definition [3.9,](#page-5-1) we see that there is a short exact sequence of complexes, split in each degree

$$
0 \to X_{\prec i+1} \to X \to X_{\succcurlyeq i+1} \to 0
$$

where $X_{\prec i+1}$ is the subcomplex of X which is termwise the projective summands generated in degree i and $X_{\geq i+1}$ is the quotient complex which is termwise all those projective summands generated in degree at least $i + 1$. This gives a triangle

$$
X_{\prec i+1} \to X \to X_{\succcurlyeq i+1} \to
$$

with $X_{\prec i+1}$ in the thick subcategory generated by $A(-i)$ and $X_{\succ i+1}$ in $P_{\succ i+1}$. By Lemma [2.6](#page-2-1) we have proved part 1.

We have that $A_0(-i) \in {}^{\perp}S_{\leq i}$, since **R** Hom_{gr} $_A(A_0(e), A_0(f)) \simeq 0$ for all $e < f$. Indeed, we may find a graded free resolution of $A_0(e)$ that exists entirely in degrees at least e. For any $X \in \mathcal{S}_{\leq i+1}$, we have the triangle

$$
X_{\geq i} \to X \to X/X_{\geq i} \to
$$

as in Lemma [3.4.](#page-4-2) Since $X_{\geq i+1} = 0$ we see $X_{\geq i}$ has cohomology concentrated in grading degree i and so is in the thick subcategory generated by $A_0(-i)$. On the other hand $X/X_{\geq i}$ is killed by $(-)_{\geq i}$ so is in $\mathcal{S}_{\leq i}$ by Lemma [3.2.](#page-3-0) Applying Lemma [2.6](#page-2-1) now proves part 2.

For the sake of clarity we introduce the following notation for the next lemma. We denote by $\mathcal{S}_{\leq i}(A)$ and $\mathcal{S}_{\leq i}(A^{\text{op}})$ the thick subcategories generated by the $A_0(e)$, for all $e > -i$, in $D^b(\text{gr }A)$ and $D^b(\text{gr }A^{\text{op}})$ respectively. We use similar notation for $\mathcal{S}_{\geq i}, \mathcal{P}_{\geq i},$ and $\mathcal{P}_{\leq i}$ in order to indicate in which category we are working.

Lemma 6.6. *Under the hypothesis of Theorem [6.4,](#page-17-0)* we have $S_{\geq i}^{\perp} = {}^{\perp} \mathcal{P}_{\geq i+a}$ as $subcategories \ of \ D^b(\text{gr}_{\geq i} A).$

Proof. As A is Gorenstein we have Grothendieck duality by Lemma [5.3.](#page-14-1) We note that restricting the duality functor $D = \mathbf{R} \underline{Hom}_{grA}(-, A)$ to $\mathcal{S}_{\geq i}$ gives an equivalence

$$
D \colon \mathcal{S}_{\geq i}(A) \xrightarrow{\cong} (\mathcal{S}_{\leq -i-a+1}(A^{\text{op}}))^{\text{op}}.
$$

Indeed, one can see this simply by computing D applied to the generators and observing equivalences send thick subcategories to thick subcategories. Similarly we can also restrict D to get an equivalence

$$
D \colon \mathcal{P}_{\leq -i-a+1}(A) \xrightarrow{\cong} (\mathcal{P}_{\geq i+a}(A^{\text{op}}))^{\text{op}}.
$$

By [3.4,](#page-4-2) [3.7,](#page-4-0) and the definition of a semi-orthogonal decomposition, we have in $D^{\mathsf{b}}(\operatorname{gr} A)$

$$
\perp S_{<-i-a+1}(A) = \mathsf{D}^{\mathsf{b}}(\text{gr } A_{\geq -i-a+1}) = \mathcal{P}_{<-i-a+1}(A)^{\perp}.
$$

We thus have

$$
\mathcal{S}_{\geq i}(A)^{\perp} \xrightarrow{\cong} ((\mathcal{S}_{<-i-a+1}(A^{\text{op}}))^{\text{op}})^{\perp} = ({}^{\perp}\mathcal{S}_{<-i-a+1}(A^{\text{op}}))^{\text{op}} = (\mathcal{P}_{<-i-a+1}(A^{\text{op}})^{\perp})^{\text{op}}
$$

$$
= {}^{\perp}((\mathcal{P}_{<-i-a+1}(A^{\text{op}}))^{\text{op}}) \xrightarrow{\cong} {}^{\perp}\mathcal{P}_{\geq i+a}(A),
$$

i.e. the functor D^2 , which is isomorphic to the identity functor, takes $\mathcal{S}_{\geq i}(A)^{\perp}$ to $^{\perp}P_{\geq i+a}(A)$ and hence these categories are equal.

Remark 6.7*.* If A is AS-Gorenstein in the sense of [\[16\]](#page-23-11) then one needs a minor additional argument to prove the above lemma. We need to check $D(\mathcal{S}_{\geq i}(A)),$ the thick subcategory of $D^{b}(gr A^{op})^{op}$ generated by the $D(A_0(e))$ for $e \leq -i$, is $(S_{*i-a+1*}(A^{op}))^{op}$. By definition

$$
\mathbf{R}\underline{\operatorname{Hom}}_{\operatorname{gr} A}(A_0(e), A) \cong \operatorname{Hom}_k(A_0, k)[-n](-e+a)
$$

and it is sufficient to check this object generates the same thick subcategory as $A_0(-e + a)$ (of course we can ignore the degree shift). This follows essentially immediately from the equivalence

$$
\operatorname{Hom}_k(-,k) \colon \mathsf{D}^{\mathsf{b}}(\operatorname{mod} A_0) \xrightarrow{\cong} \mathsf{D}^{\mathsf{b}}(\operatorname{mod} A_0^{\operatorname{op}})^{\operatorname{op}}
$$

which sends the generator A_0 to $\text{Hom}_k(A_0, k)$.

Proof of Theorem [6.4.](#page-17-0) Combining the decompositions of [3.4,](#page-4-2) [5.8](#page-16-1) via [2.8](#page-2-2) there is a semiorthogonal decomposition

(6.8)
$$
D^{b}(\text{gr } A) = (\mathcal{S}_{\leq i}, \mathcal{P}_{\geq i}, \mathcal{B}_{i}).
$$

Similarly, by [3.4,](#page-4-2) [4.18](#page-13-2) and [2.8,](#page-2-2) there is a semiorthogonal decomposition

$$
\mathsf{D}^{\mathsf{b}}(\operatorname{gr} A) = \left(\mathcal{S}_{\leq i}, \mathbf{R} \Gamma_{\geq i} \mathsf{D}^{\mathsf{b}}(\operatorname{coh} X), \mathcal{S}_{\geq i} \right).
$$

Using Lemma [6.6,](#page-18-1) we see that ${}^{\perp}\mathcal{P}_{\geq i+a} = \mathcal{S}_{\geq i}{}^{\perp} = (\mathcal{S}_{\leq i}, \mathbf{R}\Gamma_{\geq i}\mathsf{D}^{\mathsf{b}}(\text{coh } X)),$ and thus there is a semiorthogonal decomposition

(6.9)
$$
\mathsf{D}^{\mathsf{b}}(\text{gr }A) = (\mathcal{P}_{\geq i+a}, \mathcal{S}_{\leq i}, \mathbf{R}\Gamma_{\geq i}\mathsf{D}^{\mathsf{b}}(\text{coh }X)).
$$

The rest of the proof boils down to comparing the decompositions [\(6.8\)](#page-19-0) and (6.9) , depending on the sign of a.

Assume first that $a \geq 0$. Then $\mathcal{P}_{\geq i+a} \subseteq \mathsf{D}^{\mathsf{b}}(\mathrm{gr}_{\geq i} A)$ by definition, and $\mathsf{D}^{\mathsf{b}}(\mathrm{gr}_{\geq i} A) =$ $\pm \mathcal{S}_{\leq i}$ by [3.4.](#page-4-2) Hence the first two factors of (6.9) are mutually orthogonal and we may swap them to get

(6.10)
$$
D^{b}(\text{gr } A) = \left(\mathcal{S}_{\leq i}, \mathcal{P}_{\geq i+a}, \mathbf{R} \Gamma_{\geq i} D^{b}(\text{coh } X) \right).
$$

Comparing with [\(6.8\)](#page-19-0) we see that

$$
\mathsf{D}^{\mathsf{b}}(\mathrm{gr}_{\geq i} A) = (\mathcal{P}_{\geq i}, \mathcal{B}_i) = (\mathcal{P}_{\geq i+a}, \mathbf{R}\Gamma_{\geq i}\mathsf{D}^{\mathsf{b}}(\mathrm{coh}\, X)) .
$$

By [6.5](#page-18-0) there is a decomposition

$$
\mathcal{P}_{\geq i} = (\mathcal{P}_{\geq i+a}, A(-i-a+1), \ldots, A(-i+1), A(-i)).
$$

It follows there is an equality in $\mathsf{D}^{\mathsf{b}}(\mathrm{gr}_{\geq i} A)$:

$$
(A(-i-a+1),..., A(-i+1), A(-i), B_i) =
$$
R $\Gamma_{\geq i}$ **D**^b(coh X).

Applying $\widetilde{(-)}$ to both sides gives the semiorthogonal decomposition

$$
\mathsf{D}^{\mathsf{b}}(\operatorname{coh} X) = \left(\mathcal{O}(-i - a + 1), \ldots, \mathcal{O}(-i), \widetilde{\mathcal{B}}_i \right).
$$

Assume now that $a \leq 0$. In this case, $\mathcal{P}_{\geq i} \subseteq \mathcal{S}_{\leq i}^{\perp}$, i.e. $\text{Hom}_{\mathsf{D}^{\mathsf{b}}(\text{gr }A)}(\mathcal{S}_{\leq i}, \mathcal{P}_{\geq i})$ 0. To see this, it is enough to check that $\text{Hom}_{\mathsf{D}^b(\text{gr }A)}(A_0(e)[m], A(f)) = 0$ for all $e > -i, f \leq -i$ and all $m \in \mathbb{Z}$. But we have that

$$
\begin{aligned} \operatorname{Hom}\nolimits_{\mathsf{D}^{\mathsf{b}}(\operatorname{gr}\nolimits A)}(A_0(e)[m], A(f)) &\cong \operatorname{Hom}\nolimits_{\mathsf{D}^{\mathsf{b}}(\operatorname{gr}\nolimits A)}(A_0, A)(f - e)[-m] \\ &= \left(H^{-m} \mathbf{R} \underline{\operatorname{Hom}\nolimits}_{\operatorname{gr}\nolimits A}(A_0, A) \right)_{f - e} . \end{aligned}
$$

By the definition of Gorenstein, this is nonzero if and only if $-m = n$ and $f - e = -a$, however $f - e < 0$ and $-a \ge 0$. Thus we may switch the order of the first two factors of [\(6.8\)](#page-19-0) and we have a semi-orthogonal decomposition

(6.11)
$$
\mathsf{D}^{\mathsf{b}}(\text{gr }A)=(\mathcal{P}_{\geq i},\mathcal{S}_{\leq i},\mathcal{B}_i).
$$

We also have, substituting $i - a$ for i in [\(6.9\)](#page-19-1),

$$
\mathsf{D}^{\mathsf{b}}(\operatorname{gr} A) = (\mathcal{P}_{\geq i}, \mathcal{S}_{\leq i-a}, \mathbf{R}\Gamma_{\geq i-a}\mathsf{D}^{\mathsf{b}}(\operatorname{coh} X)).
$$

This shows that $(\mathcal{S}_{\leq i}, \mathbf{b} \mathsf{D}^{\mathsf{b}}_{\mathsf{sg}}(\text{gr }A)) = (\mathcal{S}_{\leq i-a}, \mathbf{R}\Gamma_{\geq i-a} \mathsf{D}^{\mathsf{b}}(\text{coh }X)).$ By [6.5](#page-18-0) we have $S_{\leq i-a} = (S_{\leq i}, A_0(-i), A_0(-i-1), \ldots, A_0(-i+a+1)).$ Thus we have

$$
\mathcal{B}_i = (A_0(-i), A_0(-i-1), \dots, A_0(-i+a+1), \mathbf{R}\Gamma_{\geq i-a}\mathsf{D}^{\mathsf{b}}(\cosh X))
$$

and applying the functor $p: D^{\mathsf{b}}(\text{gr}_{\geq i} A) \to D^{\mathsf{b}}_{\text{sg}}(\text{gr } A)$ gives the desired decomposition. \square

7. Complete intersection rings and matrix factorizations

In this section we apply the main theorem to relate the derived category of a commutative complete intersection ring to the homotopy category of graded matrix factorizations over a "generic hypersurface."

7.1. Graded matrix factorizations. Let $S = \bigoplus_{i \geq 0} S_i$ be a commutative noetherian graded ring and let $W \in S_d$, for some $d \geq 1$. A *graded matrix factorization* of W is a pair of graded projective S-modules E_1, E_0 and morphisms in gr S,

$$
e_1: E_1 \to E_0 \qquad e_0: E_0 \to E_1(d)
$$

such that $e_0e_1 = W \cdot 1_{E_1}$ and $e_1(d)e_0 = W \cdot 1_{E_0}$. A morphism h between $\mathbb{E} = (E_1 \stackrel{e_1}{\longrightarrow}$ $E_0 \xrightarrow{e_0} E_1(d)$ and $\mathbb{F} = (F_1 \xrightarrow{f_1} F_0 \xrightarrow{f_0} F_1(d))$ is a pair of maps $h_1 : E_1 \to F_1$ and $h_0 :$ $E_0 \rightarrow F_0$ making the obvious diagrams commute. One defines a homotopy between two such maps analogously to the case of a map of complexes. The category with objects graded matrix factorizations of W and morphisms homotopy equivalence classes of morphisms of matrix factorizations is called the *homotopy category of matrix factorizations* and denoted $[\text{gr-mf}(S, W)].$

Now assume that S_0 is a regular commutative ring and S is a polynomial ring over S_0 . Set $A = S/(W)$ and consider the singularity category $\mathsf{D}^{\mathsf{b}}_{\mathsf{sg}}(\text{gr }A)$ as defined

in [5.5.](#page-15-1) The assignment that sends $\mathbb{E} = (E_1 \stackrel{e_1}{\to} E_0 \stackrel{e_0}{\to} E_1(d))$ to the image of coker e_1 in $\mathsf{D}^{\mathsf{b}}_{\mathsf{sg}}(\text{gr }A)$ induces a functor

(7.1)
$$
\operatorname{coker} : [\operatorname{gr-mf}(S,W)] \to \mathsf{D}^{\mathsf{b}}_{\mathsf{sg}}(\operatorname{gr} A).
$$

It follows from work of Eisenbud [\[10\]](#page-23-12) and and Buchweitz [\[8\]](#page-22-6), but seems to have first been written down by Orlov in [\[17,](#page-23-0) §3] that this functor is an equivalence of categories.

7.2. Generic hypersurface. Let $R = Q/(f)$, where Q is a commutative regular ring of finite Krull dimension, and $f = f_1, \ldots, f_c$ is a Q-regular sequence. Define $S = Q[T_1, \ldots, T_c]$ to be the graded polynomial ring over Q with $|T_i| = 1$. Let $W = f_1 T_1 + \ldots + f_c T_c \in S_1$ and set $A = S/(W)$.

Let $Y = \text{Proj } A$ and note that there is a diagram

(7.2)
$$
\mathbb{P}_R^{c-1} = \text{Proj}(S \otimes_Q R) \xrightarrow{\beta} Y \longrightarrow \text{Proj } S = \mathbb{P}_Q^{c-1}
$$

$$
\uparrow \qquad \qquad \downarrow
$$

$$
\text{Spec } R \longrightarrow \text{Spec } Q
$$

where the vertical arrows are the canonical proper maps and each horizontal arrow is a regular closed immersion and thus has finite Tor dimension. In particular the map $\beta : \mathbb{P}^{c-1}_R \to Y$ is a regular closed immersion of codimension $c-1$. Orlov used this setup in [\[18\]](#page-23-13) to show that there is an equivalence between the singularity categories of R and Y. This equivalence was used in [\[9\]](#page-22-7) and [\[23\]](#page-23-14).

Lemma 7.3. *The functor* $\beta_* \pi^* : D^b(R) \to D^b(\text{coh } Y)$ *is fully faithful and has a right adjoint. Thus the image* $\mathcal R$ *is a right admissible subcategory of* $D^b(\text{coh } Y)$ *equivalent to* $D^b(R)$ *. Moreover, the right orthogonal of* R *is*

$$
(\beta_* \pi^* \mathsf{D}^{\mathsf{b}}(R))^{\perp} = \langle \mathcal{O}_Y(-c+2), \ldots, \mathcal{O}_Y(-1), \mathcal{O}_Y \rangle.
$$

Proof. Orlov shows in [\[18,](#page-23-13) 2.2] that the functor $\beta_* \pi^* : D^b(R) \to D^b(\text{coh } Y)$ is fully faithful and has a right adjoint (the existence of a right adjoint to β_* is one formulation of Grothendieck duality in this context). He also shows in [\[18,](#page-23-13) 2.10] that the left orthogonal of the image is $\langle \mathcal{O}_Y(1), \ldots, \mathcal{O}_Y(c-1)\rangle$; a slight reworking of this argument shows the right orthogonal is as claimed. \square

7.3. The equivalence. We continue to assume that R is a complete intersection of the form $Q/(f)$, where Q is a commutative regular ring of finite Krull dimension, and $f = f_1, \ldots, f_c$ is a Q-regular sequence. Recall that $A = Q[T_1, \ldots, T_c]/(f_1T_1 +$ $\ldots + f_cT_c = S/(W)$. We wish to apply Theorem [6.4](#page-17-0) to this ring. We first must show A is Gorenstein. This holds by "graded local duality" as in [\[7,](#page-22-8) $\S 3.4$].

Lemma 7.4. *There is an isomorphism in* $D^b(\text{gr }A)$ *,*

 $\mathbf{R} \underline{\text{Hom}}_{\text{gr }A}(A_0, A) \cong A_0[-n](c-1),$

where $n = \dim A$ *. In particular* A *is a Gorenstein ring with a-invariant* $c - 1$ *.*

We now have:

Theorem 7.5. *There is an equivalence*

$$
\Psi: \mathsf{D}^{\mathsf{b}}(R) \xrightarrow{\cong} [\operatorname{gr-mf}(S,W)]
$$

 $given by \Psi = q(\mathbf{R}\Gamma_{\geq 0})\beta_*\pi^*, where q is the composition \mathsf{D}^{\mathsf{b}}(\text{gr }A_{\geq 0}) \stackrel{p}{\rightarrow} \mathsf{D}^{\mathsf{b}}_{\mathsf{sg}}(\text{gr }A) \stackrel{\cong}{\rightarrow}$ $\left[\operatorname{gr-mf}(S,W)\right]$.

Proof. By Theorem [6.4](#page-17-0) applied to A with $i = 0$, we know that $D^b(\text{coh } Y)$ has a semiorthogonal decomposition $(\mathcal{O}_Y(-c+2), \ldots, \mathcal{O}_Y, \widetilde{\mathcal{B}}_i)$ where \mathcal{B}_i is the image of $D^b_{sg}(gr A)$ under the fully faithful functor $\mathbf{b}_i: D^b_{sg}(gr A) \to D^b(gr_{\geq i} A)$ described in [\(5.7\)](#page-16-0). Thus $\widetilde{\mathcal{B}}_i^{\perp} = (\mathcal{O}_Y(-c+2), \ldots, \mathcal{O}_Y)$, which by Lemma [7.3](#page-21-0) is also equal to \mathcal{R}^{\perp} , where R is the image of $D^b(R)$ under $\beta_* \pi^*$. Thus $\mathcal{R} = \widetilde{\mathcal{B}}_i$ and applying $qR\Gamma_{\geq 0}$ to both sides we have an equivalence

$$
\mathsf{D}^{\mathsf{b}}(R) \xrightarrow{q\mathbf{R}\Gamma_{\geq 0}\beta_*\pi^*} \mathsf{D}^{\mathsf{b}}_{\mathsf{sg}}(\textup{gr}\,A).
$$

Finally, the equivalence (7.1) finishes the proof.

In [\[9\]](#page-22-7), it was shown that there is an equivalence

$$
\mathsf{D}^{\mathsf{b}}_{\mathsf{sg}}(R) \cong [MF(\mathbb{P}^{\mathsf{c}-1}_{Q},\mathcal{O}(1), W)],
$$

where $[MF(\mathbb{P}_Q^{c-1}, \mathcal{O}(1), W)]$ is the homotopy category of matrix factorizations of locally free sheaves on \mathbb{P}_Q^{c-1} . This category has objects pairs of locally free sheaves $(\mathcal{E}_1, \mathcal{E}_0)$ on \mathbb{P}_Q^{c-1} and maps $e_1 : \mathcal{E}_1 \to \mathcal{E}_0$ and $e_0 : \mathcal{E}_0 \to \mathcal{E}_1(1)$ such that composition is multiplication by W . Morphisms are defined analogously as in the affine case above, however there is a further localization at objects that are locally contractible. There is an obvious functor $\widetilde{(-)} : [\text{gr-mf}(S,W)] \to [MF(\mathbb{P}_Q^{c-1}, \mathcal{O}(1), W)]$. This equivalence fits into the following commutative diagram, where the left hand arrow is the natural projection onto the singularity category.

$$
\begin{array}{ccc} \mathsf{D}^{\mathsf{b}}(R) \longrightarrow & \cong & \mathsf{[gr\text{-}mf}(S,W)] \\ & & \downarrow & & \downarrow \\ \mathsf{D}^{\mathsf{b}}_{\mathsf{sg}}(R) \longrightarrow & \mathsf{M}F(\mathbb{P}^{\mathsf{c}-1}_Q,\mathcal{O}(1),W)]. \end{array}
$$

REFERENCES

- [1] M. Artin and J. J. Zhang. Noncommutative projective schemes. *Adv. Math.*, 109(2):228–287, 1994.
- [2] Matthew Ballard, David Favero, and Ludmil Katzarkov. Orlov spectra: bounds and gaps. *Invent. Math.*, 189(2):359–430, 2012.
- [3] Matthew Ballard, David Favero, and Ludmil Katzarkov. Variation of geometric invariant theory quotients and derived categories, 2012.
- [4] Vladimir Baranovsky and Jeremy Pecharich. On equivalences of derived and singular categories. *Cent. Eur. J. Math.*, 8(1):1–14, 2010.
- [5] A. A. Beilinson. Coherent sheaves on P ⁿ and problems in linear algebra. *Funktsional. Anal. i Prilozhen.*, 12(3):68–69, 1978.
- [6] A. I. Bondal. Representations of associative algebras and coherent sheaves. *Izv. Akad. Nauk SSSR Ser. Mat.*, 53(1):25–44, 1989.
- [7] Winfried Bruns and J¨urgen Herzog. *Cohen-Macaulay rings*, volume 39 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1993.
- [8] Ragnar-Olaf Buchweitz. Maximal Cohen-Macaulay modules and Tate-cohomology over Gorenstein rings. Unpublished manuscript, 1987.
- [9] Jesse Burke and Mark E. Walker. Matrix factorizations in higher codimension. arXiv:1205.2552.

- [10] David Eisenbud. Homological algebra on a complete intersection, with an application to group representations. *Trans. Amer. Math. Soc.*, 260(1):35–64, 1980.
- [11] Daniel Halpern-Leistner. The derived category of a GIT quotient, 2012.
- [12] Manfred Herbst, Kentaro Hori, and David Page. Phases of n=2 theories in 1+1 dimensions with boundary, 2008.
- [13] M. Umut Isik. Equivalence of the derived category of a variety with a singularity category. arXiv:1011.1484.
- [14] Bernhard Keller, Daniel Murfet, and Michel Van den Bergh. On two examples by Iyama and Yoshino. *Compos. Math.*, 147(2):591–612, 2011.
- [15] Helmut Lenzing. Weighted projective lines and applications. In *Representations of algebras and related topics*, EMS Ser. Congr. Rep., pages 153–187. Eur. Math. Soc., Zürich, 2011.
- [16] Hiroyuki Minamoto and Izuru Mori. The structure of AS-Gorenstein algebras. *Adv. Math.*, 226(5):4061–4095, 2011.
- [17] Dmitri Orlov. Derived categories of coherent sheaves and triangulated categories of singularities. In *Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II*, volume 270 of *Progr. Math.*, pages 503–531. Birkhäuser Boston Inc., Boston, MA, 2009.
- [18] Dmitri O. Orlov. Triangulated categories of singularities, and equivalences between Landau-Ginzburg models. *Mat. Sb.*, 197(12):117–132, 2006.
- [19] N. Popescu. *Abelian categories with applications to rings and modules*. Academic Press, London, 1973. London Mathematical Society Monographs, No. 3.
- [20] Ed Segal. Equivalence between GIT quotients of Landau-Ginzburg B-models. *Comm. Math. Phys.*, 304(2):411–432, 2011.
- [21] Ian Shipman. A geometric approach to Orlov's theorem. *Compos. Math.*, 148(5):1365–1389, 2012.
- [22] J. T. Stafford and J. J. Zhang. Examples in non-commutative projective geometry. *Math. Proc. Cambridge Philos. Soc.*, 116(3):415–433, 1994.
- [23] Greg Stevenson. Subcategories of singularity categories via tensor actions. arXiv:1105.4698, to appear in Compos. Math.
- [24] Michel van den Bergh. Non-commutative crepant resolutions. In *The legacy of Niels Henrik Abel*, pages 749–770. Springer, Berlin, 2004.

Mathematics Department, UCLA, Los Angeles, CA, 90095-1555, USA *E-mail address*: jburke@math.ucla.edu

UNIVERSITAT BIELEFELD, FAKULTÄT FÜR MATHEMATIK, BIREP GRUPPE, POSTFACH 10 01 31, 33501 Bielefeld, Germany.

E-mail address: gstevens@math.uni-bielefeld.de

