

PDR Emission from the Arched-Filaments and Nearby Positions

Pablo García¹, Markus Röllig², Nicholas Abel³, Martin Steinke²,
Michael Burton⁴, and Rebecca Blackwell⁵

¹Instituto de Radioastronomía Milimétrica IRAM
Av. Divina Pastora 7, 18012 Granada, Spain
email: pgarcia@iram.es

²I. Physikalisches Institut, University of Cologne, Germany

³University of Cincinnati, USA

⁴Armagh Observatory and Planetarium, Northern Ireland

⁵School of Physical Sciences, University of Adelaide, Australia

Abstract. We investigate the physical conditions of the gas, atomic and molecular, in the filaments in the context of Photo-Dissociation Regions (PDRs) using the KOSMA-PDR mode of clumpy clouds. We also compare the [CII] vs. [NII] integrated intensity predictions in Abel et al. 2005 for HII regions and adjacent PDRs in the Galactic disk, and check for their applicability under the extreme physical conditions present in the GC. Our preliminary results show that observed integrated intensities are well reproduced by the PDR model. The gas is exposed to a relatively low Far-UV field between 10^2 - 10^3 Draine fields. The total volume hydrogen density is well constrained between 10^4 - 10^5 cm^{-3} . The hydrogen ionization rate due to cosmic-rays varies between 10^{-15} and 4×10^{-15} s^{-1} , with the highest value $\sim 10^{-14}$ s^{-1} found towards G0.07+0.04. Our results show that the line-of-sight contribution to the total distance of the filaments to the Arches Cluster is not negligible. The spatial distribution of the [CII]/[NII] ratio shows that the integrated intensity ratios are fairly homogeneously distributed for values below 10 in energy units. Calculations including variation on the [C/N] abundance ratio show that tight constraints on this ratio are needed to reproduce the observations.

Keywords. Galaxy: center, ISM: atoms, ISM: molecules, ISM: clouds, ISM: structure

1. Introduction

In the Galactic Center (GC), the Arched-Filaments are thought to originate from the interaction of the gas with the radiation field of the massive Arches Cluster [Simpson et al.(2007)]. Using the sub-mm data in García et al.(2016), and complementary millimeter data from the Mopra archive [Jones et al.(2012)], and unpublished CO(1-0) and ¹³CO(1-0) rotational lines from the ongoing Mopra CMZ Survey [R. Backwell et al., in prep.], we investigate the physical conditions of the gas, both atomic and molecular, in the Arched-Filaments region in the context of Photo-Dissociation Regions (PDRs) using the KOSMA-PDR model of clumpy clouds [Stoerzer et al.(1996), Cubick et al.(2008), and Röllig et al.(2013)]. We also compare the [CII] vs. [NII] integrated intensity predictions in Abel et al.(2005) for HII regions and adjacent PDRs in the Galactic disk, and check for their applicability under the extreme physical conditions present in the GC.

2. Overview

We have selected seven positions within the Arched-Filaments following the peaks of 20 cm continuum emission in Yusef-Zadeh & Morris(1987) and two positions close to them

tracing high density gas in Serabyn & Guesten(1987). From the KOSMA-PDR model of clumpy clouds, the following quantities are obtained from the fit to the observed integrated intensities: hydrogen nucleus number density $n(\text{H})_{s,tot}$; total clump mass within the beam M_{cl} ; FUV field in Draine units χ ; hydrogen ionization rate due to cosmic-rays $\eta(\text{H})$; clump ensemble mass lower limit M_{low} ; and clump ensemble mass upper limit M_{upper} . The model intensities can reproduce the observed intensities extremely well in almost all cases within error bars, except for the CO(1-0) transition where the model systematically underestimates observations by a factor of $\sim 2 - 3$. Calculations for the contribution of HII regions to the observed [CII] intensities as traced by [NII] emission for Galactic plane sources have been presented in Abel et al.(2005). When compared with the observed intensities in and around the Arched-Filaments location, their predictions for solar and under-solar metallicities do not match the data. The observations show that the great majority of the integrated intensity [CII]/[NII] ratios (calculated between -70 km s^{-1} and $+20 \text{ km s}^{-1}$) are fairly homogeneously distributed over the map for values below 10. Calculations including variations in the [C/N] abundance ratio and stellar clusters SEDs can explain the observed scatter in the [CII] vs. [NII] plot and suggest that, in order to use the [NII] intensity to disentangle the [CII] component arising from H II regions, tight constraints are needed on the [C/N] abundance ratio, significantly tighter than previous abundance measurements have discerned.

3. Implications

Our preliminary results show that the gas is exposed to a relatively low χ , between $10^2 - 10^3$ Draine fields, with $n(\text{H})_{s,tot}$ well constrained between $10^4 - 10^5 \text{ cm}^{-3}$. The $\eta(\text{H})$ values vary between 10^{-15} and $4 \times 10^{-15} \text{ s}^{-1}$, with the highest value $\sim 10^{-14} \text{ s}^{-1}$ found towards G0.07+0.04. The model results (specially for $n(\text{H})_{s,tot}$ and χ), for all positions as a function of projected distance from the Arches Cluster, show a large dispersion suggesting that the true distance to the cluster must have a significant line of sight component. Combining all these results puts constraints on the 3-D spatial distribution, 3-D density structure, and hydrogen ionization rate due to cosmic-rays in the Arched-Filaments. We discuss this further in an upcoming publication.

References

- Abel, N. P., Ferland, G. J., Shaw, G., & van Hoof, P. A. M. 2005, *ApJS*, 161, 65
 Cubick, M., Stutzki, J., Ossenkopf, V., Kramer, C., Röllig, M. 2008, *A&A*, 488, 623
 García, P., Simon, R., Stutzki, J., R.Güsten, M. A. Requena-Torres, & R. Higgins 2016, *A&A*, 588, A131
 Jones, P. A., Burton, M. G., Cunningham, M. R., M. A. Requena-Torres, K. M. Menten, P. Schilke, A. Belloche, S. Leurini, J. Martín-Pintado, J. Ott & A. J. Walsh 2012, *MNRAS*, 419, 2961
 Röllig, M., Szczerba, R., Ossenkopf, V., & Glück, C. 2013, *A&A*, 549, A85
 Rodríguez-Fernández, N. J., Martín-Pintado, J., & de Vicente, P. 2001, *A&A*, 377, 631
 Serabyn, E., & Guesten, R. 1987, *A&A*, 184, 133
 Simpson, J. P., Colgan, S. W. J., Cotera, A. S, Erickson, E. F., Hollenbach, D. J., Kaufman M. J., & Rubin, R. H. 2007, *ApJ*, 670, 1115
 Stoerzer, H., Stutzki, J., & Sternberg, A. 1996, *A&A*, 310, 592
 Yusef-Zadeh, F., & Morris, M. 1987, *ApJ*, 320, 545