BPS Wilson loops in $N \ge 2$ superconformal Chern–Simons–matter theories

Andrea Mauri a , Hao Ouyang bc , Silvia Penati a , Jun-Bao Wu de , Jiaju Zhang a

- ^aDipartimento di Fisica, Università degli Studi di Milano-Bicocca, & INFN, Sezione di Milano-Bicocca, Piazza della Scienza 3, I-20126 Milano, Italy
- ^bInstitute of High Energy Physics, & Theoretical Physics Center for Science Facilities, Chinese Academy of Sciences, 19B Yuquan Road, Beijing 100049, China
- ^cSchool of Physical Sciences, University of Chinese Academy of Sciences, 19A Yuquan Road, Beijing 100049, China
- ^d Center for Joint Quantum Studies, School of Science, Tianjin University, 135 Yaguan Road, Tianjin 300350, China
- ^eCenter for High Energy Physics, Peking University, 5 Yiheyuan Road, Beijing 100871, China E-mail: andrea.mauri@mi.infn.it, ouyangh@ihep.ac.cn, silvia.penati@mib.infn.it, junbao.wu@tju.edu.cn, jiaju.zhang@unimib.it

ABSTRACT: In $\mathcal{N} \geq 2$ superconformal Chern–Simons–matter theories we construct the infinite family of Bogomol'nyi-Prasad-Sommerfield (BPS) Wilson loops featured by constant parametric couplings to scalar and fermion matter, including both line Wilson loops in Minkowski spacetime and circle Wilson loops in Euclidean space. We find that the connection of the most general BPS Wilson loop cannot be decomposed in terms of double-node connections. Moreover, if the quiver contains triangles, it cannot be interpreted as a supermatrix inside a superalgebra. However, for particular choices of the parameters it reduces to the well-known connections of 1/6 BPS Wilson loops in Aharony-Bergman-Jafferis-Maldacena (ABJM) theory and 1/4 BPS Wilson loops in $\mathcal{N}=4$ orbifold ABJM theory. In the particular case of $\mathcal{N}=2$ orbifold ABJM theory we identify the gravity duals of a subset of operators. We investigate the cohomological equivalence of fermionic and bosonic BPS Wilson loops at quantum level by studying their expectation values, and find strong evidence that the cohomological equivalence holds quantum mechanically, at framing one. Finally, we discuss a stronger formulation of the cohomological equivalence, which implies non-trivial identities for correlation functions of composite operators in the defect CFT defined on the Wilson contour and allows to make novel predictions on the corresponding unknown integrals that call for a confirmation.

KEYWORDS: Supersymmetry, Wilson loops, Chern-Simons theories, M-theory

Contents		
1	Introduction	2
2	BPS WLs in $\mathcal{N}=2$ CS-matter theories	4
	2.1 Line BPS WLs in Minkowski spacetime	4
	2.2 Circle BPS WLs in Euclidean space	10
	2.3 More general $1/2$ BPS WLs	10
	2.4 The case of matter fields with non-canonical dimensions	11
3	$\mathcal{N}=2$ orbifold ABJM theory	14
	3.1 1/2 BPS WLs	14
	3.2 Gravity duals	18
4	BPS WLs at quantum level	20
	4.1 The perturbative analysis	21
	4.2 A conjecture	28
5	Conclusions	29
A	Spinor conventions in three dimensions	31
	A.1 Minkowski spacetime	31
	A.2 Euclidean space	31
В	BPS WLs in ABJ(M) theory	32
	B.1 - 1/6 BPS line WLs in Minkowski spacetime	32
	$B.2 ext{ } 1/2 ext{ BPS line WLs and their gravity duals}$	35
	B.3 Circle WLs in Euclidean space	38
\mathbf{C}	BPS WLs in $\mathcal{N}=4$ orbifold ABJM theory	39
D	Connections for 1/2 BPS WL in $\mathcal{N}=2$ orbifold ABJM	43
${f E}$	Lagrangian and Feynman rules	44
\mathbf{F}	Details on the perturbative computation	46

1 Introduction

The study of Bogomol'nyi-Prasad-Sommerfield Wilson loops (BPS WLs) have yielded tremendous insights into supersymmetric gauge theories. In particular, vacuum expectation values of BPS WLs can be computed exacly using localization techniques [1, 2], so providing functions of the coupling constants that interpolate between weak and strong coupling regimes. Therefore, for theories admitting holographic dual descriptions BPS WLs represent one of the most important tests of the AdS/CFT correspondence [3-7].

In three-dimensional superconformal Chern–Simons-matter (SCSM) theories many interesting results on BPS WLs have been obtained in recent years. One of the most important aspects is that one can construct BPS operators either generalizing the gauge connection to include couplings solely to matter bosons (bosonic WLs) [8–12] or including couplings both to bosonic and fermionic fields (fermionic WLs) [13]. While the construction of BPS WLs in $\mathcal{N} \geq 2$ SCSM quiver theories has been extensively investigated [8–17], most of the results on fermionic BPS WLs have been limited to operators with connections that can be written as 2×2 block diagonal matrices. When taking the trace, these operators correspond to linear combinations of WLs connecting adjacent nodes. Recently, new BPS WLs in $\mathcal{N}=4$ circular quiver SCSM theories with alternating levels have been constructed in [18], which are described by more general connections that cannot be decomposed as linear combinations of double–node connections. This result suggests that the general form of BPS fermionic WLs may have a richer structure waiting to be explored.

The main goal of this paper is to investigate the most general BPS WL in $\mathcal{N} \geq 2$ SCSM theories featured by parametric couplings to scalar and fermion matter¹. For a generic quiver $\mathcal{N}=2$ SCSM theory with (anti)bifundamental and/or (anti)fundamental matter fields, we write the most general expression for a WL containing arbitrary couplings to bosons and fermions and study under which conditions the operator preserves half of the supersymmetries. It turns out that with fixed preserved supercharges there is only one bosonic 1/2 BPS WL, while there is an infinite family of parametric fermionic 1/2 BPS WLs whose connection is in general a non-block-diagonal matrix. In addition, in the $\mathcal{N}=2$ case the connection does not have necessarily the structure of a superconnection of a given supergroup. This is the main novelty of our classification.

As a check of our construction we reproduce the already known operators for the Aharony–Bergman–Jafferis(–Maldacena) (ABJ(M)) and $\mathcal{N}=4$ orbifold ABJM theories [16–18]. As a new result we provide the classification of fermionic 1/2 BPS WLs for the $\mathcal{N}=2$ orbifold ABJM theory [22]. For the subset of operators that can be obtained by an orbifold quotient of the 1/2 BPS WL in ABJ(M) theory, we identify the corresponding gravity duals following the orbifold decomposition strategy in [18].

For the new infinite family of fermionic BPS WLs that we construct, it is mandatory to investigate how they behave at quantum level and understand how the parametric dependence enters their expectation values. At classical level, generalizing what happens in ABJ(M) theory and $\mathcal{N}=4$ SCSM theory [13–15, 17], we prove that

¹In this paper we consider only line and circle WLs with constant couplings to scalars, although more general WLs with contour dependent couplings have been also studied [19–21].

all the fermionic WLs, independently of their couplings, are cohomologically equivalent to the bosonic one, i.e. $W_{\text{fer}} = W_{\text{bos}} + \mathcal{Q}(\text{something})$ where \mathcal{Q} is a supercharge preserved by all the operators. Whether and how this relation gets promoted at quantum level is a crucial question to be answered when comparing fermionic and bosonic expectation values. In fact, if the cohomological equivalence survives quantum corrections, it implies that the expectation values of all the fermionic BPS WLs are equal to the expectation value of the bosonic operator, and therefore they can all be computed by the same matrix model [2, 23, 24]. However, one important subtlety that we have to take into account when comparing expectation values is framing [2]. In fact, since the localization procedure always leads to framing—one results, we expect this to be the correct regularization scheme where the classical cohomological equivalence translates into $\langle W_{\text{fer}} \rangle_1 = \langle W_{\text{bos}} \rangle_1$. This problem has been already investigated at first few perturbative orders for particular kinds of WLs both in ABJ(M) and $\mathcal{N}=4$ models [2, 23–31]. In this paper we extend this analysis to general $\mathcal{N}=2$ SCSM theories. Up to two loops, using the arguments and speculations in [30] we find that the bosonic and fermionic BPS WLs have the same framing—one expectation values not only in ABJ(M) theory but also in a generic $\mathcal{N} = 2$ SCSM theory.

This is already a strong indication that the cohomological equivalence might be valid at quantum level, at framing one, in any $\mathcal{N}=2$ SCSM theory, although a truly non-trivial check would come at higher orders where the particular choice of the superpotential characterizing the model would enter.

At classical level the expansion of W_{fer} in powers of its bosonic and fermionic couplings leads to a stronger cohomological equivalence that translates into an infinite number of non-trivial \mathcal{Q} -identities [14]. For any $\mathcal{N} \geq 2$ SCSM theory, up to two loops and at framing one these identities lead to non-trivial vanishing conditions for the correlators of the corresponding bosonic and fermionic operators (see eq. (4.8)). We make the educated conjecture that these identities survive at higher orders and discuss the novel constraints that follow for three–loop, framing–one integrals, and the implications for the defect CFT defined on the Wilson contour.

As a by–product of our analysis, we provide the two–loop expression for the framing–zero and framing–one expectation values of all the fermionic BPS WLs. While the framing–one result is parameter independent, being equal to the bosonic expectation value, the framing–zero result exhibits a non–trivial dependence on the parameters that feature the couplings to matter fields.

The rest of the paper is organized as follows. In section 2, we provide the general classification of 1/2 BPS WLs in a generic $\mathcal{N}=2$ SCSM theory, both on a line in Minkowski spacetime and on a circle in Euclidean space. We distinguish the cases of matter with canonical and non–canonical conformal dimensions. In section 3 we apply the previous results to the construction of BPS WLs in $\mathcal{N}=2$ orbifold ABJM theory and study the gravity duals of operators that can be obtained as orbifold quotients of 1/2 BPS WLs in ABJ(M) theory. Section 4 is devoted to the perturbative calculation of the WL expectation values, the discussion of the cohomological equivalence, its stronger version and its non–trivial consequences. Our conclusions are then collected in section 5. In appendix A we give spinor conventions in Minkowski and Euclidean signatures. In appendices B and C

we review the WLs in ABJ(M) theory and $\mathcal{N}=4$ orbifold ABJ(M) theory, together with their gravity duals. As a check of our classification we also reproduce the known WLs by applying our general recipe. Appendix D contains some details of section 2. Finally, in appendix E we give the Lagrangian and Feynman rules of the general $\mathcal{N}=2$ SCSM theory that would be useful to the calculations of the integrals in appendix F.

2 BPS WLs in $\mathcal{N}=2$ CS-matter theories

For a generic $\mathcal{N}=2$ SCSM quiver theory we construct the most general class of 1/2 BPS WLs featured by constant parametric couplings to matter fields. In subsection 2.1, we give full details of the classification for the case of line BPS WLs in Minkowski spacetime. We discuss cohomological equivalence between bosonic and fermionic operators and provide a toy-model example of a quiver theory to make manifest the novel features of WLs in $\mathcal{N}=2$ SCSM models. In subsection 2.2, we introduce the circle BPS WLs in Euclidean spacetime signature. These operators are the relevant observables in the context of localization and their non-trivial expectation value can be generically captured by matrix model integrals. In subsection 2.3, the classification is slightly generalized to the case of connections involving repeated nodes. Finally, in subsection 2.4 we briefly discuss the case of fields with non-canonical conformal dimensions.

2.1 Line BPS WLs in Minkowski spacetime

In three–dimensional Minkowski spacetime, we consider a generic quiver $\mathcal{N}=2$ SCSM theory with gauge group $\prod_{a=1}^n U(N_a)_{k_a}$, where k_a indicate the CS levels that can be non-vanishing or vanishing. The gauge sector of the theory is organized into $n \mathcal{N}=2$ vector multiplets, which in the Wess–Zumino gauge read (we refer to appendix A for spinor conventions)

$$\mathcal{V}^{(a)} = 2i\bar{\theta}\theta\sigma^{(a)} + 2\bar{\theta}\gamma^{\mu}\theta A_{\mu}^{(a)} + \sqrt{2}i\theta^{2}\bar{\theta}\bar{\chi}^{(a)} - \sqrt{2}i\bar{\theta}^{2}\theta\chi^{(a)} + \theta^{2}\bar{\theta}^{2}D^{(a)} \qquad a = 1,\dots, n \quad (2.1)$$

whereas the matter sector is described by sets of N_{ab} chiral multiplets in the bifundamental representation of arbitrary pairs of nodes $U(N_a) \times U(N_b)$

$$\mathcal{Z}^{i}_{(ab)} = Z^{i}_{(ab)} + i\bar{\theta}\gamma^{\mu}\theta\partial_{\mu}Z^{i}_{(ab)} - \frac{1}{4}\bar{\theta}^{2}\theta^{2}\partial^{\mu}\partial_{\mu}Z^{i}_{(ab)} + \sqrt{2}\theta\zeta^{i}_{(ab)} - \frac{i}{\sqrt{2}}\theta^{2}\bar{\theta}\gamma^{\mu}\partial_{\mu}\zeta^{i}_{(ab)} + \theta^{2}F^{i}_{(ab)} \quad (2.2)$$

with $i = 1, ..., N_{ab}$. In general, for a different (ab) pair, the i index in eq. (2.2) varies in a different range. For a = b, $\mathcal{Z}^i_{(aa)}$ describe N_{aa} matter chiral multiplets in the adjoint representation of $U(N_a)$. We also allow for the presence of N_{a0} matter multiplets in the fundamental representation of $U(N_a)$ and N_{0a} multiplets in the antifundamental representation, denoted by $\mathcal{Z}^i_{(a0)}$ and $\mathcal{Z}^i_{(0a)}$ respectively.

Complex conjugated matter fields, belonging to the anti-bifundamental representation of $U(N_a) \times U(N_b)$, are defined as $[Z^i_{(ab)}]^{\dagger} = \bar{Z}^{(ba)}_i$, $[\zeta^i_{(ab)}]^{\dagger} = \bar{\zeta}^{(ba)}_i$, $[F^i_{(ab)}]^{\dagger} = \bar{F}^{(ba)}_i$, with $i = 1, 2, \dots, N_{ab}$.

In superspace language the lagrangian of the theory is given by

$$\mathcal{L} = \mathcal{L}_{CS} + \mathcal{L}_k + \mathcal{L}_{sn} \tag{2.3}$$

where

$$\mathcal{L}_{CS} = -\sum_{a} \frac{k_{a}}{8\pi i} \int_{0}^{1} dt \operatorname{Tr}\left(\bar{D}^{\alpha} \mathcal{V}^{(a)} e^{t\mathcal{V}^{(a)}} D_{\alpha} e^{-t\mathcal{V}^{(a)}}\right) \Big|_{\theta^{2}\bar{\theta}^{2}}$$

$$\mathcal{L}_{k} = -\sum_{ab} \operatorname{Tr}\left[\bar{\mathcal{Z}}_{i}^{(ba)} e^{-\mathcal{V}^{(a)}} \mathcal{Z}_{(ab)}^{i} e^{\mathcal{V}^{(b)}}\right] \Big|_{\theta^{2}\bar{\theta}^{2}}$$

$$(2.4)$$

whereas \mathcal{L}_{sp} is the superpotential term that we do not write explicitly, as it is not relevant for the construction of BPS WLs and for the perturbative investigation at the order we work. Here we have defined $D_{\alpha} = \partial_{\alpha} + i\bar{\theta}^{\beta}\gamma^{\mu}_{\beta\alpha}\partial_{\mu}$, $\bar{D}^{\alpha} = \bar{\partial}^{\alpha} + i\gamma^{\mu\alpha\beta}\theta_{\beta}\partial_{\mu}$.

For non-vanishing k_a levels, writing (2.4) in components and extracting the equations of motion for the auxiliary fields of the vector multiplets we obtain ²

$$\sigma^{(a)} = \frac{2\pi}{k_a} \sum_b (Z_{(ab)}^i \bar{Z}_i^{(ba)} - \bar{Z}_i^{(ab)} Z_{(ba)}^i)$$

$$\chi^{(a)} = -\frac{4\pi}{k_a} \sum_b (\zeta_{(ab)}^i \bar{Z}_i^{(ba)} - \bar{Z}_i^{(ab)} \zeta_{(ba)}^i) \qquad a = 1, \dots, n$$

$$\bar{\chi}^{(a)} = -\frac{4\pi}{k_a} \sum_b (Z_{(ab)}^i \bar{\zeta}_i^{(ba)} - \bar{\zeta}_i^{(ab)} Z_{(ba)}^i) \qquad (2.5)$$

General superconformal transformations of the component fields read

$$\delta A_{\mu}^{(a)} = \frac{1}{2} (\bar{\chi}^{(a)} \gamma_{\mu} \Theta + \bar{\Theta} \gamma_{\mu} \chi^{(a)}), \qquad \delta \sigma^{(a)} = -\frac{i}{2} (\bar{\chi}^{(a)} \Theta + \bar{\Theta} \chi^{(a)})
\delta Z_{(ab)}^{i} = i \bar{\Theta} \zeta_{(ab)}^{i}, \qquad \delta \bar{Z}_{i}^{(ab)} = i \bar{\zeta}_{i}^{(ab)} \Theta
\delta \zeta_{(ab)}^{i} = (-\gamma^{\mu} D_{\mu} Z_{(ab)}^{i} - \sigma^{(a)} Z_{(ab)}^{i} + Z_{(ab)}^{i} \sigma^{(b)}) \Theta - Z_{(ab)}^{i} \vartheta + i F_{(ab)}^{i} \bar{\Theta}
\delta \bar{\zeta}_{i}^{(ab)} = \bar{\Theta} (\gamma^{\mu} D_{\mu} \bar{Z}_{i}^{(ab)} - \bar{Z}_{i}^{(ab)} \sigma^{(b)} + \sigma^{(a)} \bar{Z}_{i}^{(ab)}) - \bar{\vartheta} \bar{Z}_{i}^{(ab)} - i \bar{F}_{i}^{(ab)} \Theta$$
(2.6)

where $\Theta \equiv \theta + x^{\mu}\gamma_{\mu}\vartheta$, $\bar{\Theta} \equiv \bar{\theta} - \bar{\vartheta}x^{\mu}\gamma_{\mu}$ are linear combinations of $(\theta, \bar{\theta})$ spinors parametrizing Poincaré supersymmetry transformations, and $(\vartheta, \bar{\vartheta})$ ones parameterizing superconformal transformations ³. The definition of covariant derivative can be found in (E.3).

The 1/2 BPS WLs. We construct WLs defined along the timelike infinite straight line $x^{\mu} = (\tau, 0, 0)$, which preserve half of the supersymmetries. Decomposing the spinorial charges as in (A.6), without loss of generality we choose the preserved supercharges to be $Q_+, \bar{Q}_-, S_+, \bar{S}_-$, i.e. we require the operator to be invariant under

$$\delta = \bar{\theta}_{-}Q_{+} + \bar{Q}_{-}\theta_{+} + \bar{\vartheta}_{-}S_{+} + \bar{S}_{-}\vartheta_{+} \tag{2.7}$$

In the rest of the paper we will shortly identify the preserved supercharges with the corresponding $\theta_+, \bar{\theta}_-, \vartheta_+, \bar{\vartheta}_-$ parameters.

²Through the paper we use the convention that repeated flavor i indices are summed, while summations on node indices $a, b, c \cdots$ are explicitly indicated. Repeated node indices with no explicit sum are meant to be fixed.

³Here we only consider the case where matter fields in the chiral multiplets have canonical conformal dimensions. The more general case will be discussed in section 2.4. The difference does not appear when we focus on Poincaré supercharges.

The first kind of 1/2 BPS operator is the so-called bosonic WL defined as [8]

$$W_{\text{bos}} = \mathcal{P}e^{-i\int d\tau L_{\text{bos}}(\tau)}$$
 (2.8)

where the $n \times n$ connection matrix is given by

$$L_{\text{bos}} = \text{diag}(A_0^{(1)} - \sigma^{(1)}, A_0^{(2)} - \sigma^{(2)}, \cdots, A_0^{(n)} - \sigma^{(n)})$$
(2.9)

It is easy to check that this operator is invariant under (2.7).

Using equations of motion (2.5) for $\sigma^{(a)}$, $a = 1, 2, \dots, n$, this generalized connection ends up including quadratic couplings to matter scalars. However, as usually happens in three dimensions, we can look for more general operators with connections containing couplings also to fermions. We then consider the fermionic operator

$$W_{\text{fer}} = \mathcal{P}e^{-i\int d\tau L_{\text{fer}}(\tau)}$$
 (2.10)

with a $n \times n$ connection [13]

$$L_{\text{fer}} = L_{\text{bos}} + B + F \tag{2.11}$$

where L_{bos} is given in (2.9), whereas the B and F entries

$$B_{(ab)} = \sum_{c} (R_{abi}^{c}{}^{j} Z_{(ac)}^{i} \bar{Z}_{j}^{(cb)} + R_{abij}^{c} Z_{(ac)}^{i} Z_{(cb)}^{j} + S_{ab}^{c}{}^{i}{}_{j} \bar{Z}_{i}^{(ac)} Z_{(cb)}^{j}) + S_{ab}^{c}{}^{ij} \bar{Z}_{i}^{(ac)} \bar{Z}_{j}^{(cb)})$$

$$F_{(ab)} = \bar{m}_{i}^{ab} \zeta_{(ab)+}^{i} + n_{ab}^{i} \bar{\zeta}_{i-}^{(ab)} \equiv [\bar{M}_{\zeta}]_{(ab)} + [N_{\bar{\zeta}}]_{(ab)}$$

$$(2.12)$$

contain couplings to bilinear scalars and linear fermions respectively, parametrized by tobe-determined matrices and vectors.

The requirement for W_{fer} to be 1/2 BPS can be traded with the search for a Grassmann odd matrix G satisfying [13, 32]

$$\delta L_{\text{fer}} = \partial_{\tau} G + i[L_{\text{fer}}, G] \tag{2.13}$$

Inserting decomposition (2.11) for L_{fer} this condition splits into a set of Grassmann even and odd constraints, respectively

$$\delta B = i[F, G]$$

$$\delta F = \partial_{\tau} G + i[L_{\text{bos}} + B, G]$$
(2.14)

From the Grassmann odd one we obtain⁴

$$G = -i\bar{M}_Z\theta_+ + iN_{\bar{Z}}\bar{\theta}_-, \qquad [B, G] = 0$$
 (2.15)

where we have defined

$$[\bar{M}_Z]_{(ab)} \equiv \bar{m}_i^{ab} Z_{(ab)}^i, \qquad [N_{\bar{Z}}]_{(ab)} \equiv n_{ab}^i \bar{Z}_i^{(ab)}$$
 (2.16)

⁴For the straight line it is sufficient to focus on super–Poincaré symmetries, since once these supercharges are preserved also the superconformal ones are automatically preserved.

Using G in (2.15) in the Grassmann even part of eq. (2.14) we eventually obtain non-trivial relations among the coefficients

$$R_{abi}^{c}{}^{j} = \bar{m}_{i}^{ac} n_{cb}^{j}, \quad S_{ab}^{c}{}^{i}{}_{j} = n_{ac}^{i} \bar{m}_{j}^{cb}$$

$$R_{abij}^{c} = S_{ab}^{c}{}^{ij} = \bar{m}_{i}^{ac} \bar{m}_{j}^{cb} = n_{ac}^{i} n_{cb}^{j} = 0$$
(2.17)

In particular, they imply the following relation

$$B = \bar{M}_Z N_{\bar{Z}} + N_{\bar{Z}} \bar{M}_Z \tag{2.18}$$

It is easy to check that this expression automatically satisfies the second condition in (2.15). Solutions to constraints (2.17) exhibit several interesting features:

- Setting a = b = c in the second line of (2.17) we obtain $\bar{m}_i^{aa} = n_{aa}^i = 0$ (no summation on $a = 1, \dots, N_{aa}$). Therefore, although adjoint matter may appear in the theory, adjoint fermion fields $\zeta_{(aa)}^i$ or $\bar{\zeta}_i^{(aa)}$ can never appear in the connection. In other words, the diagonal blocks of the connection contain only bosonic couplings.
- For $a \neq b$ fixed, the $B_{(ab)}$ and $F_{(ab)}$ entries can be simultaneously non-vanishing. Therefore, in general the $L_{\rm fer}$ connection is not a superconnection, i.e. it does not give a representation of a supergroup. However, this does not contradict what has been already found for ABJM and $\mathcal{N}=4$ orbifold ABJM theories where $L_{\rm fer}$ are indeed superconnections for the $U(N_1|N_2)$ supergroup. In fact, as we show below discussing a toy model, the conditions $B_{(ab)} \neq 0$, $F_{(ab)} \neq 0$ can occur simultaneously only when the quiver diagram contains triangles.
- Finally, constraints $\bar{m}_i^{ab}\bar{m}_j^{ba}=n_{ab}^in_{ba}^j=0$ following from (2.17) imply that if a positive chirality fermion appears in $F_{(ab)}$ ($\bar{m}_i^{ab}\neq 0$), then all fermion fields of positive chirality in $F_{(ba)}$ must be absent ($\bar{m}_i^{ba}=0$). Similarly, if fermion fields of both chiralities appear in $F_{(ab)}$, then $F_{(ba)}=0$.

To summarize, for a generic $\mathcal{N}=2$ SCSM theory we have constructed bosonic and fermionic 1/2 BPS WLs (2.8) and (2.10) with connections

$$L_{\text{bos}} = \text{diag}(A_0^{(1)} - \sigma^{(1)}, A_0^{(2)} - \sigma^{(2)}, \cdots, A_0^{(n)} - \sigma^{(n)})$$

$$L_{\text{fer}} = L_{\text{bos}} + B + F, \qquad B = \bar{M}_Z N_{\bar{Z}} + N_{\bar{Z}} \bar{M}_Z, \quad F = \bar{M}_\zeta + N_{\bar{\zeta}}$$

$$[\bar{M}_Z]_{(ab)} = \bar{m}_i^{ab} Z_{(ab)}^i, \quad [N_{\bar{Z}}]_{(ab)} = n_{ab}^i \bar{Z}_i^{(ab)}$$

$$[\bar{M}_\zeta]_{(ab)} = \bar{m}_i^{ab} \zeta_{(ab)+}^i, \quad [N_{\bar{\zeta}}]_{(ab)} = n_{ab}^i \bar{\zeta}_{i-}^{(ab)}$$

$$(2.19)$$

In particular, a generic fermionic connection has the following structure

$$L_{\text{fer}} = \begin{pmatrix} A_0^{(1)} - \sigma^{(1)} + B_{(11)} & B_{(12)} + F_{(12)} & \cdots & B_{(1n)} + F_{(1n)} \\ B_{(21)} + F_{(21)} & A_0^{(2)} - \sigma^{(2)} + B_{(22)} & \cdots & B_{(2n)} + F_{(2n)} \\ \vdots & \vdots & \ddots & \vdots \\ B_{(n1)} + F_{(n1)} & B_{(n2)} + F_{(n2)} & \cdots & A_0^{(n)} - \sigma^{(n)} + B_{(nn)} \end{pmatrix}$$
(2.20)

with the caveat that some of the bosonic and fermionic couplings may be forced to be absent, as discussed above. We note that the fundamental and antifundamental fields do not appear explicitly in the connection, but the connection may depend on them through $\sigma^{(a)}$.

Similarly, we can construct bosonic and fermionic 1/2 BPS WL $\tilde{W}_{\rm bos}$, $\tilde{W}_{\rm fer}$ preserving the complementary set of supercharges $\theta_-, \bar{\theta}_+, \vartheta_-, \bar{\vartheta}_+$. The corresponding connections read

$$\tilde{L}_{\text{bos}} = \text{diag}(A_0^{(1)} + \sigma^{(1)}, A_0^{(2)} + \sigma^{(2)}, \cdots, A_0^{(n)} + \sigma^{(n)})
\tilde{L}_{\text{fer}} = \tilde{L}_{\text{bos}} + \tilde{B} + \tilde{F}, \qquad \tilde{B} = -(\bar{M}_Z N_{\bar{Z}} + N_{\bar{Z}} \bar{M}_Z), \quad \tilde{F} = \bar{M}_{\zeta} - N_{\bar{\zeta}}
[\bar{M}_Z]_{(ab)} = \bar{m}_i^{ab} Z_{(ab)}^i, \quad [N_{\bar{Z}}]_{(ab)} = n_{ab}^i \bar{Z}_i^{(ab)}
[\bar{M}_{\zeta}]_{(ab)} = \bar{m}_i^{ab} \zeta_{(ab)-}^i, \quad [N_{\bar{\zeta}}]_{(ab)} = n_{ab}^i \bar{\zeta}_{i+}^{(ab)}$$
(2.21)

where the constant parameters \bar{m}_i^{ab} , n_{ab}^i satisfy the same constraints (2.17).

Cohomological equivalence. For ABJ(M) and $\mathcal{N}=4$ SCSM theories the full parametric family of fermionic BPS WLs has been shown to be (classically) cohomologically equivalent to the bosonic WL [13–15, 17, 18]. This means that the difference between a given fermionic operator and the bosonic one can be written as \mathcal{Q} (something), with \mathcal{Q} being a suitable linear combination of conserved supercharges shared by the two operators.

We now prove that this property holds also in the $\mathcal{N}=2$ setting: The general fermionic 1/2 BPS WL with connection (2.11) is classically \mathcal{Q} -equivalent to the bosonic 1/2 BPS WL with connection (2.9).

According to the analysis of [13, 14], this is the case if we manage to find κ , Λ and \mathcal{Q} quantities that satisfy

$$\kappa \Lambda^2 = B, \quad Q\Lambda = F, \quad QL_{\text{bos}} = 0$$

$$QF = \partial_{\tau}(i\kappa\Lambda) + i[L_{\text{bos}}, i\kappa\Lambda] \tag{2.22}$$

Using (2.14), it is easy to see that in the present case a solution to the above equations is given by

$$\kappa = 1, \quad \Lambda = \bar{M}_Z + N_{\bar{Z}}, \quad \mathcal{Q} = Q_+ + \bar{Q}_-$$
(2.23)

This solution implies the classical identity

$$W_{\text{fer}} - W_{\text{bos}} = QV \tag{2.24}$$

where V is a known function of the gauge and matter fields of the theory.

A triangular quiver toy model. Aimed at highlighting novel properties of the 1/2 BPS WLs that we have constructed, we consider the simple case of a $\mathcal{N}=2$ SCSM theory associated to a triangle quiver diagram, as given in figure 1.

Specifying the general WL construction to this case, a particular solution to the BPS conditions in (2.17) takes the form

$$L_{\text{fer}} = \begin{pmatrix} \mathcal{A}^{(1)} & \bar{m}^{1}\zeta_{1+} & 0\\ 0 & \mathcal{A}^{(2)} & 0\\ n_{3}\bar{\zeta}_{-}^{3} & n_{3}\bar{m}^{1}\bar{Z}^{3}Z_{1} + \bar{m}^{2}\zeta_{2+} & \mathcal{A}^{(3)} \end{pmatrix}$$
(2.25)




Figure 1. The quiver diagram of a toy model $\mathcal{N} = 2$ SCSM theory.

where

$$\mathcal{A}^{(1)} = A_0^{(1)} + \frac{2\pi}{k_1} (-Z_1 \bar{Z}^1 - Z_3 \bar{Z}^3)$$

$$\mathcal{A}^{(2)} = A_0^{(2)} + \frac{2\pi}{k_2} (\bar{Z}^1 Z_1 + \bar{Z}^2 Z_2)$$

$$\mathcal{A}^{(3)} = A_0^{(3)} + \frac{2\pi}{k_2} (-Z_2 \bar{Z}^2 + \bar{Z}^3 Z_3)$$
(2.26)

Notably, L_{fer} has a block entry which contains a sum of both bosonic and fermionic field combinations. As a consequence, the full connection ceases to be a supermatrix.

It is easy to see that a necessary condition for this feature to appear is the presence of a triangle in the quiver diagram. In fact, non-diagonal entries in (2.20) connecting adjacent nodes may contain both bilinear scalar terms $B_{(a\,a+1)}$ and linear fermion ones $F_{(a\,a+1)}$. While the fermionic entry corresponds to a fermionic arrow connecting the two nodes, the bosonic bilinear can be formed only passing by the third vertex of the triangle. Therefore, models with suitably chosen matter content allow for the existence of matrix entries in the WL connection that exhibit mixed Grassmann parity.

Operators with this structure have no counterpart in the classification of $\mathcal{N}=4$ and $\mathcal{N}=6$ models, where the underlying dynamics of the fermionic WLs seems to be captured by a gauge supergroup (this is particularly manifest in the Higgsing derivation of the 1/2 BPS operators [32, 33]). It would be interesting to understand the implications of the existence of these WLs, especially in terms of a string dual description at strong coupling.

Lightlike WLs. We can also construct bosonic and fermionic BPS WLs along the lightlike infinite straight line $x^{\mu} = (\tau, \tau, 0)$. The corresponding connections read

$$L_{\text{bos}} = \text{diag}(A_0^{(1)} + A_1^{(1)}, A_0^{(2)} + A_1^{(2)}, \cdots, A_0^{(n)} + A_1^{(n)})$$

$$L_{\text{fer}} = L_{\text{bos}} + B + F, \qquad B = \bar{M}_Z N_{\bar{Z}} + N_{\bar{Z}} \bar{M}_Z, \quad F = \bar{M}_\zeta + N_{\bar{\zeta}}$$

$$[\bar{M}_Z]_{(ab)} = \bar{m}_i^{ab} Z_{(ab)}^i, \quad [N_{\bar{Z}}]_{(ab)} = n_{ab}^i \bar{Z}_i^{(ab)}$$

$$[\bar{M}_\zeta]_{(ab)} = \bar{m}_i^{ab} \zeta_{(ab)1}^i, \quad [N_{\bar{\zeta}}]_{(ab)} = n_{ab}^i \bar{\zeta}_{i1}^{(ab)}$$

$$(2.27)$$

where the constant parameters \bar{m}_i^{ab} , n_{ab}^i satisfy constraints (2.17). Index 1 in the last line indicates the first component of a spinor in the standard spinorial notation ψ_{α} , $\alpha = 1, 2$.

The lightlike bosonic WL is 3/4 BPS with preserved supercharges

$$\theta_2, \ \bar{\theta}_2, \ \vartheta_1, \ \bar{\vartheta}_1, \ \vartheta_2, \ \bar{\vartheta}_2$$
 (2.28)

whereas the lightlike fermionic WL is 1/2 BPS with preserved supercharges

$$\theta_2, \ \bar{\theta}_2, \ \vartheta_2, \ \bar{\vartheta}_2$$
 (2.29)

2.2 Circle BPS WLs in Euclidean space

The previous procedure can be easily generalized to construct 1/2 BPS WLs along the circle $x^{\mu} = (\cos \tau, \sin \tau, 0)$ in Euclidean space. The computational steps follow closely the ones of the Minkowskian case, thus we report only the final result.

In a generic $\mathcal{N}=2$ SCSM theory, the bosonic and fermionic WLs can be written as

$$W_{\text{bos}} = \text{Tr}\mathcal{P}e^{-i \oint d\tau L_{\text{bos}}(\tau)}, \qquad W_{\text{fer}} = \text{Tr}\mathcal{P}e^{-i \oint d\tau L_{\text{fer}}(\tau)}$$
 (2.30)

with connections

$$L_{\text{bos}} = \text{diag}(A_{\mu}^{(1)}\dot{x}^{\mu} + i\sigma^{(1)}, A_{\mu}^{(2)}\dot{x}^{\mu} + i\sigma^{(2)}, \cdots, A_{\mu}^{(n)}\dot{x}^{\mu} + i\sigma^{(n)})$$

$$L_{\text{fer}} = L_{\text{bos}} + B + F, \quad B = -i(\bar{M}_{Z}N_{\bar{Z}} + N_{\bar{Z}}\bar{M}_{Z}), \quad F = \bar{M}_{\zeta} - N_{\bar{\zeta}}$$

$$[\bar{M}_{Z}]_{(ab)} = \bar{m}_{i}^{ab}Z_{(ab)}^{i}, \quad [N_{\bar{Z}}]_{(ab)} = n_{ab}^{i}\bar{Z}_{i}^{(ab)}$$

$$[\bar{M}_{\zeta}]_{(ab)} = \bar{m}_{i}^{ab}\zeta_{(ab)+}^{i}, \quad [N_{\bar{\zeta}}]_{(ab)} = n_{ab}^{i}\bar{\zeta}_{i-}^{(ab)}$$

$$(2.31)$$

Note that $\zeta^i_{(ab)+} = \mathrm{i} u_+ \zeta^i_{(ab)}$, $\bar{\zeta}^{(ab)}_{i-} = \mathrm{i} \bar{\zeta}^{(ab)}_i u_-$, with the spinorial couplings u_\pm being defined in (A.12). They are non-trivial functions of the contour, whereas the scalar couplings are still contour independent. The supercharges preserved by W_{bos} and W_{fer} are

$$\vartheta = -i\gamma_3 \theta, \quad \bar{\vartheta} = -\bar{\theta}i\gamma_3$$
(2.32)

Similarly, we can construct 1/2 BPS bosonic and fermionic WLs \tilde{W}_{bos} , \tilde{W}_{fer} preserving complementary supercharges, $\vartheta = i\gamma_3\theta$, $\bar{\vartheta} = \bar{\theta}i\gamma_3$ and corresponding to

$$\tilde{L}_{\text{bos}} = \text{diag}(A_0^{(1)} - i\sigma^{(1)}, A_0^{(2)} - i\sigma^{(2)}, \cdots, A_0^{(n)} - i\sigma^{(n)})
\tilde{L}_{\text{fer}} = \tilde{L}_{\text{bos}} + \tilde{B} + \tilde{F}, \qquad \tilde{B} = i(\bar{M}_Z N_{\bar{Z}} + N_{\bar{Z}} \bar{M}_Z), \quad \tilde{F} = \bar{M}_\zeta + N_{\bar{\zeta}}
[\bar{M}_Z]_{(ab)} = \bar{m}_i^{ab} Z_{(ab)}^i, \quad [N_{\bar{Z}}]_{(ab)} = n_{ab}^i \bar{Z}_i^{(ab)}
[\bar{M}_\zeta]_{(ab)} = \bar{m}_i^{ab} \zeta_{(ab)-}^i, \quad [N_{\bar{\zeta}}]_{(ab)} = n_{ab}^i \bar{\zeta}_{i+}^{(ab)}$$
(2.33)

In both cases it is not difficult to prove that the classical cohomological equivalence in (2.24) still holds.

2.3 More general 1/2 BPS WLs

The previous class of 1/2 BPS WLs can be generalized to include extra operators constructed in the following way.

We arbitrarily select a subset of n' nodes of the quiver diagram and label the corresponding gauge fields as $A^{(s_{a'})}$, with $a' = 1', 2', \dots, n'$ and $s_{a'} \in \{1, 2, \dots, n\}$. Note that n' can be greater or smaller than n, and each node can be either not chosen or chosen more than once. For example, for n = 4, we may choose n' = 3 with $s_{a'} = 1, 1, 3, 4, 4$.

Along the line $x^{\mu}=(\tau,0,0)$ in Minkowski spacetime we construct the bosonic 1/2 BPS WL with connection

$$L_{\text{bos}} = \text{diag}(A_0^{(s_{1'})} - \sigma^{(s_{1'})}, A_0^{(s_{2'})} - \sigma^{(s_{2'})}, \cdots, A_0^{(s_{n'})} - \sigma^{(s_{n'})})$$
(2.34)

It is easy to prove that this operator preserves the $\theta_+, \bar{\theta}_-, \vartheta_+, \bar{\vartheta}_-$ supercharges.

More generally, starting from the most general ansatz we can construct the fermionic 1/2 BPS WL with connection

$$L_{\text{fer}} = L_{\text{bos}} + B + F$$

$$B = \bar{M}_Z N_{\bar{Z}} + N_{\bar{Z}} \bar{M}_Z, \quad F = \bar{M}_{\zeta} + N_{\bar{\zeta}}$$

$$[\bar{M}_{\zeta}]_{(a'b')} = \bar{m}_i^{(a'b')} \zeta_{(s_{a'}s_{b'})+}^i, \quad [N_{\bar{\zeta}}]_{(a'b')} = n_{(a'b')}^i \bar{\zeta}_{i-}^{(s_{a'}s_{b'})}$$

$$[\bar{M}_Z]_{(a'b')} = \bar{m}_i^{(a'b')} Z_{(s_{a'}s_{b'})}^i, \quad [N_{\bar{Z}}]_{(a'b')} = n_{(a'b')}^i \bar{Z}_i^{(s_{a'}s_{b'})}$$
(2.35)

Imposing the operator to preserve the $\theta_+, \bar{\theta}_-, \vartheta_+, \bar{\vartheta}_-$ supercharges, leads to non-trivial constraints for the constant parameters

$$\sum_{c',s_{c'}=c} \bar{m}_i^{(a'c')} \bar{m}_j^{(c'b')} = \sum_{c',s_{c'}=c} n_{(a'c')}^i n_{(c'b')}^j = 0$$
(2.36)

where we sum over c', whereas indices a', b', c, i, j are kept fixed. Again, the fundamental and antifundamental fields do not appear explicitly in the connection, but the connection may depend on them through $\sigma^{(s_{a'})}$.

2.4 The case of matter fields with non-canonical dimensions

In SCSM theories with $\mathcal{N} \geq 3$ supersymmetries, the matter fields always have canonical R-charges since the R-symmetry group $SO(\mathcal{N})$ is non-Abelian. For theories with $\mathcal{N}=2$ supersymmetry the situation is different, as the R-symmetry can mix with other U(1) flavor symmetries present in the theory. In this case the R-charges of matter fields have to be determined by F-maximization [34] and generically they turn out to be non-canonical (we informally call these matter fields non-canonical). In this section we briefly discuss how to construct BPS WLs in this case.

We consider the UV theory on S^3 . The metric on S^3 of radius r is

$$ds^{2} = \left(1 + \frac{|x|^{2}}{4r^{2}}\right)^{-2} \left[(dx^{1})^{2} + (dx^{2})^{2} + (dx^{3})^{2} \right]$$
 (2.37)

where $x^{\mu}=(x^1,x^2,x^3)$ are stereographic coordinates on S³ and $|x|^2=(x^1)^2+(x^2)^2+(x^3)^2$. Given the vielbeins $E^a_{\mu}=\left(1+\frac{|x|^2}{4r^2}\right)^{-1}e^a_{\mu}$, $e^a=(dx^1,dx^2,dx^3)$, the gamma matrices are

defined as $\Gamma_{\mu} = E_{\mu}^{a} \gamma_{a}$, $\gamma_{\mu} = e_{\mu}^{a} \gamma_{a}$, where γ^{μ} are the three–dimensional gammas in eq. (A.8).

The Killing spinors Θ , $\bar{\Theta}$ satisfy

$$D_{\mu}\Theta = \Gamma_{\mu}\tilde{\Theta}, \quad D_{\mu}\tilde{\Theta} = -\frac{1}{4r^{2}}\Gamma_{\mu}\Theta$$

$$D_{\mu}\bar{\Theta} = -\bar{\tilde{\Theta}}\Gamma_{\mu}, \quad D_{\mu}\bar{\tilde{\Theta}} = \frac{1}{4r^{2}}\bar{\Theta}\Gamma_{\mu}$$
(2.38)

Note that Θ , $\bar{\Theta}$ are independent, and are different from the ones in (2.6). The solution can be found in [1] and reads

$$\Theta = \frac{1}{\sqrt{1 + \frac{|x|^2}{4r^2}}} (\theta + x^{\mu} \gamma_{\mu} \vartheta), \quad \tilde{\Theta} = \frac{1}{\sqrt{1 + \frac{|x|^2}{4r^2}}} (\vartheta - \frac{x^{\mu} \gamma_{\mu}}{4r^2} \theta)$$

$$\bar{\Theta} = \frac{1}{\sqrt{1 + \frac{|x|^2}{4r^2}}} (\bar{\theta} - \bar{\vartheta} x^{\mu} \gamma_{\mu}), \quad \tilde{\tilde{\Theta}} = \frac{1}{\sqrt{1 + \frac{|x|^2}{4r^2}}} (\bar{\vartheta} + \bar{\theta} \frac{x^{\mu} \gamma_{\mu}}{4r^2})$$
(2.39)

When $r \to \infty$, Θ , $\bar{\Theta}$ go back to Θ , $\bar{\Theta}$ introduced in eq. (2.6), with $\tilde{\Theta}$ going to ϑ and $\bar{\bar{\Theta}}$ going to $\bar{\vartheta}$. The superconformal transformations of $A_{\mu}^{(a)}$, $\sigma^{(a)}$, $Z_{(ab)}^{i}$, $\bar{Z}_{i}^{(ab)}$ can be obtained from (2.6) by simply replacing γ_{μ} with Γ_{μ} and Θ , $\bar{\Theta}$ with (2.39), while the superconformal transformations of the fermions in the chiral multiplets are

$$\delta \zeta_{(ab)}^{i} = (-\Gamma^{\mu} D_{\mu} Z_{(ab)}^{i} - \sigma^{(a)} Z_{(ab)}^{i} + Z_{(ab)}^{i} \sigma^{(b)}) \Theta - \frac{2\Delta_{(ab)}}{3} Z_{(ab)}^{i} \Gamma^{\mu} D_{\mu} \Theta + i F_{(ab)}^{i} \bar{\Theta}$$

$$\delta \bar{\zeta}_{i}^{(ab)} = \bar{\Theta} (\Gamma^{\mu} D_{\mu} \bar{Z}_{i}^{(ab)} - \bar{Z}_{i}^{(ab)} \sigma^{(b)} + \sigma^{(a)} \bar{Z}_{i}^{(ab)}) + \frac{2\Delta_{(ba)}}{3} \bar{Z}_{i}^{(ab)} D_{\mu} \bar{\Theta} \Gamma^{\mu} - i \bar{\Theta} \bar{F}_{i}^{(ab)} (2.40)$$

where $\Delta_{(ab)}$ are the R-charges of the matter fields.

Usually, the action for the non–canonical matter is not superconformal in the UV. But one can construct actions which are invariant under supersymmetries generated by the left-invariant Killing spinors [34, 35]. In stereographic coordinates, the left-invariant Killing spinors take the form

$$D_{\mu}\Theta = \frac{\mathrm{i}}{2r}\Gamma_{\mu}\Theta \quad \Rightarrow \quad \vartheta = \frac{\mathrm{i}}{2r}\theta$$

$$D_{\mu}\bar{\Theta} = -\frac{\mathrm{i}}{2r}\bar{\Theta}\Gamma_{\mu} \quad \Rightarrow \quad \bar{\vartheta} = \frac{\mathrm{i}}{2r}\bar{\theta}$$
(2.41)

and transformations (2.40) restricted to left-invariant Killing spinors read

$$\delta \zeta_{(ab)}^{i} = (-\Gamma^{\mu} D_{\mu} Z_{(ab)}^{i} - \sigma^{(a)} Z_{(ab)}^{i} + Z_{(ab)}^{i} \sigma^{(b)}) \Theta - \frac{i \Delta_{(ab)}}{r} Z_{(ab)}^{i} \Theta + i F_{(ab)}^{i} \bar{\Theta}$$

$$\delta \bar{\zeta}_{i}^{(ab)} = \bar{\Theta} (\Gamma^{\mu} D_{\mu} \bar{Z}_{i}^{(ab)} - \bar{Z}_{i}^{(ab)} \sigma^{(b)} + \sigma^{(a)} \bar{Z}_{i}^{(ab)}) - \frac{i \Delta_{(ba)}}{r} \bar{\Theta} \bar{Z}_{i}^{(ab)} - i \bar{\Theta} \bar{F}_{i}^{(ab)}$$
(2.42)

In many cases the theory flows to an IR fixed point with superconformal symmetry. At this point the R-charges $\Delta_{(ab)}$ are determined by F-maximization [34], modulo some flat directions associated to the transformations of the R-charges which leave the partition function

in the matrix model invariant [36]. This is related to the invariance of the transformation rules in (2.42) under the shift

$$\Delta_{(ab)} \to \Delta_{(ab)} + \delta^{(a)} - \delta^{(b)}, \quad \sigma^{(a)} \to \sigma^{(a)} - i\frac{\delta^{(a)}}{r}$$
 (2.43)

Along a general contour $x^{\mu}(\tau)$ on S³, we construct the bosonic WL with connection

$$L_{\text{bos}} = \text{diag}(A_{\mu}^{(1)} \dot{x}^{\mu} + i|\dot{x}|\sigma^{(1)}, A_{\mu}^{(2)} \dot{x}^{\mu} + i|\dot{x}|\sigma^{(2)}, \cdots, A_{\mu}^{(n)} \dot{x}^{\mu} + i|\dot{x}|\sigma^{(n)})$$
(2.44)

Using the shift transformation (2.43), we can also construct fermionic WLs with non-canonical matter satisfying the condition

$$\Delta_{(ab)} = 1/2 + \delta^{(a)} - \delta^{(b)} \tag{2.45}$$

The fermionic WL has connection

$$L_{\text{fer}} = L_{\text{bos}} + B + F + C$$

$$C = -\text{diag}(\delta^{(1)}, \delta^{(2)}, \cdots, \delta^{(n)}) |\dot{x}| / r$$

$$B = -\text{i} |\dot{x}| (\bar{M}_Z N_{\bar{Z}} + N_{\bar{Z}} \bar{M}_Z), \quad F = |\dot{x}| (\bar{M}_\zeta + N_{\bar{\zeta}})$$

$$[\bar{M}_Z]_{(ab)} = \bar{m}_i^{ab} Z_{(ab)}^i, \quad [N_{\bar{Z}}]_{(ab)} = n_{ab}^i \bar{Z}_i^{(ab)}$$

$$[\bar{M}_\zeta]_{(ab)} = \bar{m}_i^{ab} u_+ \zeta_{(ab)}^i, \quad [N_{\bar{\zeta}}]_{(ab)} = n_{ab}^i \bar{\zeta}_i^{(ab)} u_-$$

$$\bar{m}_i^{ac} \bar{m}_i^{cb} = n_{ac}^i n_{cb}^j = 0$$
(2.46)

where on the x^{μ} contour the $u_{\pm}(\tau)$ spinors satisfy

$$\Gamma_{\mu}\dot{x}^{\mu}u_{\pm} = \pm |\dot{x}|u_{\pm}, \quad u_{+}u_{-} = -u_{-}u_{+} = -i, \quad u_{+}u_{+} = u_{-}u_{-} = u_{+}D_{\tau}u_{-} = u_{-}D_{\tau}u_{+} = 0$$

$$(2.47)$$

The bosonic and fermionic WLs are locally BPS with preserved supercharges

$$\Gamma_{\mu}\dot{x}^{\mu}\Theta = -|\dot{x}|\Theta, \quad \bar{\Theta}\Gamma_{\mu}\dot{x}^{\mu} = -|\dot{x}|\bar{\Theta}$$
 (2.48)

To make the WLs globally BPS, we need to choose the contour to be a great circle. Here we discuss two special cases.

In the first case, we choose the great circle $x^{\mu} = 2r(\cos \tau, \sin \tau, 0)$ in S³. Using (2.46) with $|\dot{x}| = r$, we obtain the connections of bosonic and fermionc WLs

$$L_{\text{bos}} = \text{diag}(A_{\mu}^{(1)}\dot{x}^{\mu} + \text{i}r\sigma^{(1)}, A_{\mu}^{(2)}\dot{x}^{\mu} + \text{i}r\sigma^{(2)}, \cdots, A_{\mu}^{(n)}\dot{x}^{\mu} + \text{i}r\sigma^{(n)})$$

$$L_{\text{fer}} = L_{\text{bos}} + B + F + C$$

$$C = -\text{diag}(\delta^{(1)}, \delta^{(2)}, \cdots, \delta^{(n)})$$

$$B = -\text{i}r(\bar{M}_{Z}N_{\bar{Z}} + N_{\bar{Z}}\bar{M}_{Z}), \quad F = r(\bar{M}_{\zeta} - N_{\bar{\zeta}})$$

$$[\bar{M}_{Z}]_{(ab)} = \bar{m}_{i}^{ab}Z_{(ab)}^{i}, \quad [N_{\bar{Z}}]_{(ab)} = n_{ab}^{i}\bar{Z}_{i}^{(ab)}$$

$$[\bar{M}_{\zeta}]_{(ab)} = \bar{m}_{i}^{ab}\zeta_{(ab)+}^{i}, \quad [N_{\bar{\zeta}}]_{(ab)} = n_{ab}^{i}\bar{\zeta}_{i-}^{(ab)}$$

$$\bar{m}_{i}^{ac}\bar{m}_{i}^{cb} = n_{ac}^{i}n_{cb}^{j} = 0$$
(2.49)

with u_{\pm} being the spinors defined in (A.12). The preserved supercharges are

$$\vartheta = -\frac{\mathrm{i}}{2r}\gamma_3\theta, \quad \bar{\vartheta} = -\frac{\mathrm{i}}{2r}\bar{\theta}\gamma_3 \tag{2.50}$$

Using (2.41) we can write preserved supersymmetries as $\theta_2, \bar{\theta}_1$.

As a second case, we consider the great circle obtained by the stereographic projection of the straight line $x^{\mu} = (0, 0, \tau)$ from \mathbb{R}^3 to \mathbb{S}^3 , along which we define

$$u_{+\alpha} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad u_{-\alpha} = \begin{pmatrix} 0 \\ i \end{pmatrix}$$
$$u_{+}^{\alpha} = (0, -1), \quad u_{-}^{\alpha} = (i, 0)$$
(2.51)

Here $|\dot{x}| = (1 + \frac{\tau^2}{4r^2})^{-1}$, and (2.47) are satisfied. The connections of bosonic and fermionic WLs take the form

$$L_{\text{bos}} = \operatorname{diag}(A_{\mu}^{(1)}\dot{x}^{\mu} + \mathrm{i}|\dot{x}|\sigma^{(1)}, A_{\mu}^{(2)}\dot{x}^{\mu} + \mathrm{i}|\dot{x}|\sigma^{(2)}, \cdots, A_{\mu}^{(n)}\dot{x}^{\mu} + \mathrm{i}|\dot{x}|\sigma^{(n)})$$

$$L_{\text{fer}} = L_{\text{bos}} + B + F + C$$

$$C = -\operatorname{diag}(\delta^{(1)}, \delta^{(2)}, \cdots, \delta^{(n)})|\dot{x}|/r$$

$$B = -|\dot{x}|(\bar{M}_{Z}N_{\bar{Z}} + N_{\bar{Z}}\bar{M}_{Z}), \quad F = |\dot{x}|(\bar{M}_{\zeta} + N_{\bar{\zeta}})$$

$$[\bar{M}_{Z}]_{(ab)} = \bar{m}_{i}^{ab}Z_{(ab)}^{i}, \quad [N_{\bar{Z}}]_{(ab)} = n_{ab}^{i}\bar{Z}_{i}^{(ab)}$$

$$[\bar{M}_{\zeta}]_{(ab)} = \bar{m}_{i}^{ab}\zeta_{(ab)2}^{i}, \quad [N_{\bar{\zeta}}]_{(ab)} = n_{ab}^{i}\bar{\zeta}_{i1}^{(ab)}$$

$$\bar{m}_{i}^{ac}\bar{m}_{j}^{cb} = n_{ac}^{i}n_{cb}^{j} = 0$$

$$(2.52)$$

In the limit $r \to \infty$, the corresponding WL is along an infinite straight line and the preserved supersymmetries are half of the Poincaré supersymmetries, i.e. θ_2 and $\bar{\theta}_1$. Instead, conformal supersymmetries can be preserved only if there is no non–canonical matter in the theory.

3 $\mathcal{N}=2$ orbifold ABJM theory

In this section we give full details for an explicit example of $\mathcal{N}=2$ SCSM theory that admits a gravity dual description, namely the case of the $\mathcal{N}=2$ orbifold of ABJM theory. In this case, exploiting our knowledge of the gravity duals of the 1/2 BPS WLs in ABJM theory [13, 33], after introducing the classification we identify the gravity duals of some 1/2 BPS and non-BPS WLs in the $\mathcal{N}=2$ orbifolded version.

In the rest of this section we heavily refer to appendix B where the construction of 1/2 BPS WLs and their gravity duals is reviewed for the ABJ(M) theory.

$3.1 ext{ } 1/2 ext{ BPS WLs}$

The $\mathcal{N}=2$ orbifold ABJM theory can be obtained starting from $U(rN)_k \times U(rN)_{-k}$ ABJM model and performing the Z_r quotient as in [22] ⁵. In the usual notations of ABJM theory,

⁵For ABJM two different orbifold projections can be applied, which lead to $\mathcal{N}=2$ and $\mathcal{N}=4$ quotients. Here we focus on the $\mathcal{N}=2$ orbifold, whereas the $\mathcal{N}=4$ case is reviewed in appendix C.

which we summarize in appendix B, the fields are decomposed as

$$A_{\mu} = \operatorname{diag}(A_{\mu}^{(1)}, A_{\mu}^{(2)}, \cdots, A_{\mu}^{(r)}), \quad B_{\mu} = \operatorname{diag}(B_{\mu}^{(1)}, B_{\mu}^{(2)}, \cdots, B_{\mu}^{(r)})$$

$$\phi_{1,2} = \operatorname{diag}(\phi_{1,2}^{(1)}, \phi_{1,2}^{(2)}, \cdots, \phi_{1,2}^{(r)}), \quad \psi^{1,2} = \operatorname{diag}(\psi_{(1)}^{1,2}, \psi_{(2)}^{1,2}, \cdots, \psi_{(r)}^{1,2})$$

$$\phi_{3} = \begin{pmatrix} 0 & \phi_{3}^{(r)} \\ \phi_{3}^{(1)} & 0 & \\ \phi_{3}^{(2)} & \ddots & \\ & \ddots & 0 \\ & & \phi_{3}^{(r-1)} & 0 \end{pmatrix}, \quad \phi_{4} = \begin{pmatrix} 0 & \phi_{4}^{(1)} \\ & 0 & \phi_{4}^{(2)} \\ & & \ddots & \ddots \\ & & 0 & \phi_{4}^{(r-1)} \\ \phi_{4}^{(r)} & & 0 \end{pmatrix}$$

$$\psi^{3} = \begin{pmatrix} 0 & \psi_{(1)}^{3} \\ & 0 & \psi_{(2)}^{3} \\ & & \ddots & \ddots \\ & & 0 & \psi_{(r-1)}^{4} \end{pmatrix}, \quad \psi^{4} = \begin{pmatrix} 0 & \psi_{(1)}^{4} & 0 \\ & \psi_{(1)}^{4} & 0 \\ & & \ddots & 0 \\ & & & \ddots & 0 \\ & & & & \ddots & 0 \\ & & & & & & \ddots & 0 \end{pmatrix}$$

$$\psi_{(r)}^{3} & 0 & 0 & 0 \end{pmatrix}, \quad \psi^{4} = \begin{pmatrix} 0 & \psi_{(1)}^{4} & 0 & 0 \\ & \psi_{(1)}^{4} & 0 & 0 \\ & & & & \ddots & \ddots \\ & & & & \ddots & 0 \\ & & & & & \ddots & 0 \\ & & & & & & & \ddots & 0 \end{pmatrix}$$

$$\psi_{(r)}^{3} & 0 & 0 & 0 & 0 \end{pmatrix}$$

$$(3.1)$$

where A_{μ} and B_{μ} are the gauge connections associated to $U(rN)_k$ and $U(rN)_{-k}$ respectively, whereas in SU(4) R–symmetry notations $\phi_{I=1,\dots,4}$ and the corresponding $\bar{\psi}_I$ fermions build up the matter multiplets in the bifundamental representation of the gauge group.

Orbifold decomposition (3.1) corresponds to choosing the unbroken supercharges as $\theta^{12} = \bar{\theta}_{34} = \theta$, $\theta^{34} = \bar{\theta}_{12} = \bar{\theta}$, $\vartheta^{12} = \bar{\vartheta}_{34} = \vartheta$, $\vartheta^{34} = \bar{\vartheta}_{12} = \bar{\vartheta}$. The resulting $\mathcal{N} = 2$ theory is described by the quiver diagram in figure 2(a).

In order to make contact with the classification of the previous section we have to temporarily use the alternative notation defined in the quiver diagram of figure 2(b). In equations, the two sets of conventions are related as

$$A_{\mu}^{(\ell)} = A_{\mu}^{(\ell)}, \quad B_{\mu}^{(\ell)} = A_{\mu}^{(r+\ell)}$$

$$\phi_{I}^{(\ell)} = (Z_{1}^{(\ell)}, Z_{2}^{(\ell)}, \bar{Z}_{(\ell)}^{3}, \bar{Z}_{(\ell)}^{4})$$

$$\psi_{(\ell)}^{I} = (-\zeta_{2}^{(\ell)}, \zeta_{1}^{(\ell)}, -\bar{\zeta}_{(\ell)}^{4}, \bar{\zeta}_{(\ell)}^{3})$$
(3.2)

A similar change of notations for the ABJ(M) case is detailed in appendix B. We will refer to the notations of figure 2(a) and 2(b) as to the ABJM and $\mathcal{N}=2$ notation, respectively.

The use of $\mathcal{N}=2$ notation allows to easily exploit the results of section 2 for writing down the connection of 1/2 BPS fermionic WLs. They are given by the general recipe (2.19) with

$$\sigma^{(\ell)} = \frac{2\pi}{k} (Z_1^{(\ell)} \bar{Z}_{(\ell)}^1 + Z_2^{(\ell)} \bar{Z}_{(\ell)}^2 - \bar{Z}_{(\ell-1)}^3 Z_3^{(\ell-1)} - \bar{Z}_{(\ell)}^4 Z_4^{(\ell)})$$

$$\sigma^{(r+\ell)} = \frac{2\pi}{k} (\bar{Z}_{(\ell)}^1 Z_1^{(\ell)} + \bar{Z}_{(\ell)}^2 Z_2^{(\ell)} - Z_3^{(\ell)} \bar{Z}_{(\ell)}^3 - Z_4^{(\ell-1)} \bar{Z}_{(\ell-1)}^4)$$
(3.3)




Figure 2. The quiver diagram of $\mathcal{N}=2$ orbifold ABJM theory in (a) ABJM notation and (b) $\mathcal{N}=2$ notation. We have omitted the complex conjugates of the matter fields.

and non–vanishing blocks of $\bar{M}_Z,\,N_{\bar{Z}},\,\bar{M}_\zeta,\,N_{\bar{\zeta}}$ matrices given by

$$[\bar{M}_{Z}]_{(\ell,r+\ell)} = \bar{m}_{(\ell)}^{1} Z_{1}^{(\ell)} + \bar{m}_{(\ell)}^{2} Z_{2}^{(\ell)}, \quad [\bar{M}_{Z}]_{(r+\ell,\ell+1)} = \bar{m}_{(\ell)}^{3} Z_{3}^{(\ell)}, \quad [\bar{M}_{Z}]_{(r+\ell+1,\ell)} = \bar{m}_{(\ell)}^{4} Z_{4}^{(\ell)}$$

$$[N_{\bar{Z}}]_{(\ell,r+\ell+1)} = n_{4}^{(\ell)} \bar{Z}_{(\ell)}^{4}, \quad [N_{\bar{Z}}]_{(\ell+1,r+\ell)} = n_{3}^{(\ell)} \bar{Z}_{(\ell)}^{3}, \quad [N_{\bar{Z}}]_{(r+\ell,\ell)} = n_{1}^{(\ell)} \bar{Z}_{(\ell)}^{1} + n_{2}^{(\ell)} \bar{Z}_{(\ell)}^{2}$$

$$[\bar{M}_{\zeta}]_{(\ell,r+\ell)} = \bar{m}_{(\ell)}^{1} \zeta_{1+}^{(\ell)} + \bar{m}_{(\ell)}^{2} \zeta_{2+}^{(\ell)}, \quad [\bar{M}_{\zeta}]_{(r+\ell,\ell+1)} = \bar{m}_{3}^{3} \zeta_{3+}^{(\ell)}, \quad [\bar{M}_{\zeta}]_{(r+\ell+1,\ell)} = \bar{m}_{(\ell)}^{4} \zeta_{4+}^{4}$$

$$[N_{\bar{\zeta}}]_{(\ell,r+\ell+1)} = n_{4}^{(\ell)} \bar{\zeta}_{(\ell)-}^{4}, \quad [N_{\bar{\zeta}}]_{(\ell+1,r+\ell)} = n_{3}^{(\ell)} \bar{\zeta}_{(\ell)-}^{3}, \quad [N_{\bar{\zeta}}]_{(r+\ell,\ell)} = n_{1}^{(\ell)} \bar{\zeta}_{(\ell)-}^{1} + n_{2}^{(\ell)} \bar{\zeta}_{(\ell)-}^{2}$$

$$(3.4)$$

The parameters are subject to the following constraints

$$\bar{m}_{(\ell)}^{1,2}\bar{m}_{(\ell-1)}^{3,4} = \bar{m}_{(\ell)}^{1,2}\bar{m}_{(\ell)}^{3,4} = n_{1,2}^{(\ell)}n_{3,4}^{(\ell-1)} = n_{1,2}^{(\ell)}n_{3,4}^{(\ell)} = 0 \tag{3.5}$$

Having in mind to find out the gravity duals of these WLs we now translate the connections back to ABJM notation. Mimicking what is done in appendix B for the ABJ(M) case (see eq. (B.12)), we first redefine the parameters as

$$\bar{m}_{(\ell)}^{1} = \sqrt{\frac{4\pi}{k}} \bar{\alpha}_{2}^{(\ell)}, \quad \bar{m}_{(\ell)}^{2} = -\sqrt{\frac{4\pi}{k}} \bar{\alpha}_{1}^{(\ell)}, \quad \bar{m}_{(\ell)}^{3} = -\sqrt{\frac{4\pi}{k}} \delta_{(\ell)}^{4}, \quad \bar{m}_{(\ell)}^{4} = \sqrt{\frac{4\pi}{k}} \delta_{(\ell)}^{3},$$

$$n_{1}^{(\ell)} = \sqrt{\frac{4\pi}{k}} \beta_{(\ell)}^{2}, \quad n_{2}^{(\ell)} = -\sqrt{\frac{4\pi}{k}} \beta_{(\ell)}^{1}, \quad n_{3}^{(\ell)} = \sqrt{\frac{4\pi}{k}} \bar{\gamma}_{4}^{(\ell)}, \quad n_{4}^{(\ell)} = -\sqrt{\frac{4\pi}{k}} \bar{\gamma}_{3}^{(\ell)}$$

$$(3.6)$$

Constraints (3.5) now read

$$\bar{\alpha}_{1,2}^{(\ell)}\delta_{(\ell-1)}^{3,4} = \bar{\alpha}_{1,2}^{(\ell)}\delta_{(\ell)}^{3,4} = \bar{\gamma}_{3,4}^{(\ell)}\beta_{(\ell)}^{1,2} = \bar{\gamma}_{3,4}^{(\ell)}\beta_{(\ell+1)}^{1,2} = 0 \tag{3.7}$$

Then, expressing the 1/2 BPS operator W_{fer} in terms of $\mathcal{N}=2$ orbifold ABJM fields (3.2), we can rewrite the matrix connection as

$$L_{\text{fer}} = \begin{pmatrix} \mathcal{A} & f_1 \\ f_2 & \mathcal{B} \end{pmatrix} \tag{3.8}$$

where for $r \geq 5$ the explicit expressions of the matrix blocks read

$$\mathcal{A} = \begin{pmatrix}
\mathcal{A}^{(1)} & 0 & h_{1}^{(1)} & h_{3}^{(r-1)} & 0 \\
0 & \mathcal{A}^{(2)} & 0 & \ddots & h_{3}^{(r)} \\
h_{3}^{(1)} & 0 & \mathcal{A}^{(3)} & \ddots & h_{1}^{(r-3)} \\
& \ddots & \ddots & \ddots & 0 & h_{1}^{(r-2)} \\
h_{1}^{(r-1)} & h_{3}^{(r-3)} & 0 & \mathcal{A}^{(r-1)} & 0 \\
0 & h_{1}^{(r)} & h_{3}^{(r-2)} & 0 & \mathcal{A}^{(r)}
\end{pmatrix}$$

$$\mathcal{B} = \begin{pmatrix}
\mathcal{B}^{(1)} & 0 & h_{4}^{(1)} & h_{2}^{(r-2)} & 0 \\
0 & \mathcal{B}^{(2)} & 0 & \ddots & h_{2}^{(r)} \\
h_{2}^{(1)} & 0 & \mathcal{B}^{(3)} & \ddots & h_{4}^{(r-3)} \\
& \ddots & \ddots & \ddots & 0 & h_{4}^{(r-2)} \\
h_{4}^{(r-1)} & h_{2}^{(r-3)} & 0 & \mathcal{B}^{(r-1)} & 0 \\
0 & h_{4}^{(r)} & h_{2}^{(r-2)} & 0 & \mathcal{B}^{(r)}
\end{pmatrix}$$

$$f_{1} = \begin{pmatrix}
f_{1}^{(1)} f_{3}^{(1)} & f_{5}^{(r)} \\
f_{5}^{(1)} f_{1}^{(2)} & \ddots & \\
& \ddots & \ddots & f_{3}^{(r-1)} \\
f_{2}^{(r)} & f_{5}^{(r-1)} f_{1}^{(r)}
\end{pmatrix}, \quad f_{2} = \begin{pmatrix}
f_{2}^{(1)} f_{6}^{(1)} & f_{4}^{(r)} \\
f_{4}^{(1)} f_{2}^{(2)} & \ddots & \\
& \ddots & \ddots & f_{6}^{(r-1)} \\
f_{6}^{(r)} & f_{4}^{(r-1)} f_{2}^{(r)}
\end{pmatrix}$$

with definitions

$$\begin{split} \mathcal{A}^{(\ell)} &= A_0^{(\ell)} + \frac{2\pi}{k} \left[(-1 + 2\beta_{(\ell)}^2 \bar{\alpha}_2^{(\ell)}) \phi_1^{(\ell)} \bar{\phi}_{(\ell)}^1 + (-1 + 2\beta_{(\ell)}^1 \bar{\alpha}_1^{(\ell)}) \phi_2^{(\ell)} \bar{\phi}_{(\ell)}^2 \right. \\ &\quad - 2\beta_{(\ell)}^1 \bar{\alpha}_2^{(\ell)} \phi_1^{(\ell)} \bar{\phi}_{(\ell)}^2 - 2\beta_{(\ell)}^2 \bar{\alpha}_1^{(\ell)} \phi_2^{(\ell)} \bar{\phi}_{(\ell)}^1 \\ &\quad + (1 - 2\delta_{(\ell-1)}^4 \bar{\gamma}_4^{(\ell-1)}) \phi_3^{(\ell-1)} \bar{\phi}_{(\ell-1)}^3 + (1 - 2\delta_{(\ell)}^3 \bar{\gamma}_3^{(\ell)}) \phi_4^{(\ell)} \bar{\phi}_4^{(\ell)} \right] \\ \mathcal{B}^{(\ell)} &= B_0^{(\ell)} + \frac{2\pi}{k} \left[(-1 + 2\beta_{(\ell)}^2 \bar{\alpha}_2^{(\ell)}) \bar{\phi}_{(\ell)}^1 \phi_1^{(\ell)} + (-1 + 2\beta_{(\ell)}^1 \bar{\alpha}_1^{(\ell)}) \bar{\phi}_{(\ell)}^2 \phi_2^{(\ell)} \right. \\ &\quad - 2\beta_{(\ell)}^1 \bar{\alpha}_2^{(\ell)} \bar{\phi}_{(\ell)}^2 \phi_1^{(\ell)} - 2\beta_{(\ell)}^2 \bar{\alpha}_1^{(\ell)} \bar{\phi}_{(\ell)}^1 \phi_2^{(\ell)} \\ &\quad + (1 - 2\delta_{(\ell)}^4 \bar{\gamma}_4^{(\ell)}) \bar{\phi}_3^3 \phi_3^{(\ell)} + (1 - 2\delta_{(\ell-1)}^3 \bar{\gamma}_3^{(\ell-1)}) \bar{\phi}_{(\ell-1)}^4 \phi_4^{(\ell-1)} \right] \\ h_1^{(\ell)} &= \frac{4\pi}{k} \delta_{(\ell+1)}^4 \bar{\gamma}_3^{(\ell)} \phi_4^{(\ell)} \bar{\phi}_3^3 + (1 - 2\delta_{(\ell-1)}^3 \bar{\gamma}_3^{(\ell-1)}) \bar{\phi}_4^4 \phi_4^4 \right. \\ h_2^{(\ell)} &= \frac{4\pi}{k} \delta_{(\ell+1)}^3 \bar{\gamma}_3^{(\ell)} \phi_4^{(\ell)} \bar{\phi}_3^3 + h_4^{(\ell)} = \frac{4\pi}{k} \delta_{(\ell)}^4 \bar{\gamma}_3^{(\ell)} \bar{\phi}_4^{(\ell+1)} \bar{\phi}_4^3 + \delta_{(\ell)}^4 \bar{\phi}_4^4 \right. \\ h_2^{(\ell)} &= \frac{4\pi}{k} \delta_{(\ell+1)}^3 \bar{\gamma}_4^{(\ell)} \bar{\phi}_4^4 + \bar{\phi}_3^{(\ell)} , \quad h_4^{(\ell)} &= \frac{4\pi}{k} \delta_{(\ell)}^4 \bar{\gamma}_3^{(\ell)} \phi_3^4 + \bar{\phi}_4^4 \right. \\ h_2^{(\ell)} &= \sqrt{\frac{4\pi}{k}} (\bar{\alpha}_1^{(\ell)} \psi_{(\ell)+}^1 + \bar{\alpha}_2^{(\ell)} \psi_{(\ell)+}^2), \quad f_3^{(\ell)} &= \sqrt{\frac{4\pi}{k}} \bar{\gamma}_3^{(\ell)} \psi_{(\ell)-}^3, \quad f_5^{(\ell)} &= \sqrt{\frac{4\pi}{k}} \bar{\gamma}_4^{(\ell)} \psi_{(\ell)-}^4 \\ f_2^{(\ell)} &= \sqrt{\frac{4\pi}{k}} (\bar{\psi}_{1-}^{(\ell)} \beta_{(\ell)}^1 + \bar{\psi}_{2-}^{(\ell)} \beta_{(\ell)}^2), \quad f_4^{(\ell)} &= -\sqrt{\frac{4\pi}{k}} \bar{\psi}_3^{(\ell)} \delta_3^4, \quad f_6^{(\ell)} &= -\sqrt{\frac{4\pi}{k}} \bar{\psi}_4^{(\ell)} \delta_4^4 \right. \\ h_2^{(\ell)} &= \sqrt{\frac{4\pi}{k}} (\bar{\psi}_{1-}^2 \beta_{(\ell)}^1 + \bar{\psi}_{2-}^2 \beta_{(\ell)}^2), \quad f_4^{(\ell)} &= -\sqrt{\frac{4\pi}{k}} \bar{\psi}_3^{(\ell)} \delta_3^4, \quad f_6^{(\ell)} &= -\sqrt{\frac{4\pi}{k}} \bar{\psi}_4^{(\ell)} \delta_4^4 \right)$$

In the case of shorter orbifold quivers (r = 2, 3, 4), the connections get slightly modified by boundary effects. We report their explicit expressions in appendix D.

We note that in this case, compared with the general $\mathcal{N}=2$ quiver models, the full connection (3.8) is still given by a proper supermatrix.

It is interesting to observe that the subset of 1/2 BPS operators (3.8)–(3.10) identified by the condition

$$\bar{\alpha}_{1,2}^{(\ell)} = \bar{\alpha}_{1,2}, \quad \beta_{(\ell)}^{1,2} = \beta^{1,2}, \quad \bar{\gamma}_{3,4}^{(\ell)} = \bar{\gamma}_{3,4}, \quad \delta_{(\ell)}^{3,4} = \delta^{3,4}$$
 (3.11)

correspond to direct orbifold projections of the 1/6 BPS $W_{\rm fer}$ of the ABJ(M) theory defined in (B.3). Among them, it is possible to select the ones which arise from the orbifold projection of 1/2 BPS WLs $W_{1/2}$ in ABJ(M) defined in (B.13), after setting $\beta^{1,2} = \alpha^{1,2}/|\alpha|^2$, $\bar{\gamma}_{3,4} = \delta^{3,4} = 0$.

Similarly, from results in section 2 we obtain the 1/2 BPS WLs \tilde{W}_{fer} that preserve supercharges complementary to the ones of W_{fer} , and among them we can select the WLs corresponding to the projections of the \tilde{W}_{fer} and $\tilde{W}_{1/2}$ operators of ABJ(M) theory defined in (B.6) and (B.13).

3.2 Gravity duals

Given the known gravity dual configurations of 1/2 BPS WLs $W_{1/2}$ and $W_{1/2}$ in ABJM theory, we can perform their orbifold projection and identify the (anti–)M2-brane solutions dual to WLs in $\mathcal{N}=2$ orbifold ABJM theory. In general, this operation may break SUSY. Nonetheless, as we now show some BPS configurations survive, which are dual to particular 1/2 BPS WLs constructed in the previous section.

The $\mathcal{N}=2$ orbifold ABJM theory is dual to M-theory in $AdS_4 \times S^7/(Z_{rk} \times Z_r)$ background with metric

$$ds^{2} = R^{2} \left(\frac{1}{4} ds_{AdS_{4}}^{2} + ds_{S^{7}/(Z_{rk} \times Z_{r})}^{2} \right)$$
(3.12)

where the AdS_4 metric is given in (B.17) and the S^7 one in (B.19), respectively.

In this case the $Z_{rk} \times Z_r$ quotient is realized by imposing the following identification on the C^4 coordinates

$$(z_1, z_2, z_3, z_4) \sim e^{\frac{2\pi i}{rk}} (z_1, z_2, z_3, z_4), \quad (z_1, z_2, z_3, z_4) \sim (e^{\frac{2\pi i}{r}} z_1, e^{-\frac{2\pi i}{r}} z_2, z_3, z_4)$$
 (3.13)

This is equivalent to requiring (see eq. (B.18))

$$\zeta \sim \zeta - \frac{8\pi}{rk}, \quad \phi_1 \sim \phi_1 - \frac{4\pi}{r}$$
(3.14)

As reviewed in appendix B.2, the general solution to the Killing spinor equations in M-theory on $AdS_4 \times S^7$ background reads [10, 33]

$$\epsilon = u^{\frac{1}{2}}h(\epsilon_1 + x^{\mu}\gamma_{\mu}\epsilon_2) - u^{-\frac{1}{2}}\gamma_3h\epsilon_2 \qquad \mu = 0, 1, 2$$
(3.15)

where ϵ_1 , ϵ_2 are two constant Majorana spinors satisfying $\gamma^{012}\epsilon_i = \epsilon_i$, i = 1, 2, and h is given in eq. (B.22).

We decompose the Killing spinors on the basis of gamma matrices eigenstates as in (B.27, B.25) where θ^i are associated to the Poincaré supercharges, while ϑ^i are the superconformal supercharges. Imposing the orbifold projection in (3.14) leads to the constraints

$$\mathcal{L}_{\partial_{\zeta}} \epsilon = \mathcal{L}_{\partial_{\phi_1}} \epsilon = 0 \tag{3.16}$$

where the definition of spinorial Lie derivative with respect to a Killing vector K is [37]

$$\mathcal{L}_{K}\epsilon = K^{\mu}\nabla_{\mu}\epsilon + \frac{1}{4}\nabla_{\mu}K_{\nu}\Gamma^{\mu\nu}\epsilon \tag{3.17}$$

Here we convert gamma matrices with tangent space indices to the ones with curved space indices using the vielbein of S^7 . Eq. (3.16) leads then to the constraints

$$(\gamma_{3b} + \gamma_{58} + \gamma_{47} + \gamma_{69})\epsilon_i = (\gamma_{3b} - \gamma_{58})\epsilon_i = 0, \qquad i = 1, 2$$
(3.18)

which on the η_i spinors defined in (B.26) translates into

$$t_1 + t_2 + t_3 + t_4 = 0, t_1 - t_2 = 0 (3.19)$$

Therefore, the only surviving eigenstates correspond to

$$(t_1, t_2, t_3, t_4) = (+ + --), (- - ++)$$
(3.20)

and we are led to $\epsilon_1 = \theta^2 \otimes \eta_2 + \theta^7 \otimes \eta_7$ and $\epsilon_2 = \vartheta^2 \otimes \eta_2 + \vartheta^7 \otimes \eta_7$.

Redefining

$$\theta^2 \equiv \theta, \quad \theta^7 \equiv \bar{\theta}, \quad \vartheta^2 \equiv \vartheta, \quad \vartheta^7 \equiv \bar{\vartheta}, \quad \eta_2 \equiv \eta, \quad \eta_7 \equiv \bar{\eta}$$
 (3.21)

we finally obtain the two Killing spinors surviving the orbifold projection

$$\epsilon_1 = \theta \otimes \eta + \bar{\theta} \otimes \bar{\eta}, \qquad \epsilon_2 = \vartheta \otimes \eta + \bar{\vartheta} \otimes \bar{\eta}$$
 (3.22)

The corresponding field theory is in fact $\mathcal{N}=2$ superconformal invariant.

Comparing (3.21) with (B.32), it turns out that the relation of SUSY parameters in ABJM and $\mathcal{N}=2$ orbifold ABJM theories is the following

$$\theta^{12} = \bar{\theta}_{34} = \theta, \quad \theta^{34} = \bar{\theta}_{12} = \bar{\theta}, \quad \vartheta^{12} = \bar{\vartheta}_{34} = \vartheta, \quad \vartheta^{34} = \bar{\vartheta}_{12} = \bar{\vartheta}$$
 (3.23)

As reviewed in appendix B, in ABJM theory the 1/2 BPS operator $W_{1/2}[\bar{\alpha}_I]$ is dual to an M2-brane wrapping the cycle in S^7/Z_k specified by $\bar{\alpha}_I$ given in eq. (B.34). Similarly, $\tilde{W}_{1/2}[\bar{\alpha}_I]$ is dual to an anti-M2-brane wrapping the same cycle [33]. On the other hand, as described in the previous section, orbifolding $W_{1/2}[\bar{\alpha}_I]$ and $\tilde{W}_{1/2}[\bar{\alpha}_I]$ leads to WLs in $\mathcal{N}=2$ orbifold ABJM theory. Therefore, performing the same orbifold projection on the corresponding gravity dual solutions, we obtain the (anti)-M2-brane configurations dual to a particular subset of WLs of the $\mathcal{N}=2$ theory.

In general, when the $\bar{\alpha}_I$ parameters (B.14) appearing in the $L_{1/2}[\bar{\alpha}_I]$ connection (B.13) satisfy $\bar{\alpha}_1\alpha^1 + \bar{\alpha}_2\alpha^2 \neq 0$, $\bar{\alpha}_3\alpha^3 + \bar{\alpha}_4\alpha^4 \neq 0$, the resulting operators and their dual configurations are non–BPS. However, for special choices of the parameters this may happen. Precisely,

- When $\bar{\alpha}_{3,4} = 0$, the WL in $\mathcal{N} = 2$ orbifold ABJM theory and its dual M2-brane in $\mathrm{AdS}_4 \times \mathrm{S}^7/(\mathrm{Z}_{rk} \times \mathrm{Z}_r)$ are 1/2 BPS. The WL preserves supercharges $\theta_+, \bar{\theta}_-, \vartheta_+, \bar{\vartheta}_-,$ and is just a special case of the 1/2 BPS operator W_{fer} of the previous subsection.
- When $\bar{\alpha}_{1,2} = 0$, the WL and its dual M2-brane are 1/2 BPS. The WL preserves supercharges $\theta_-, \bar{\theta}_+, \vartheta_-, \bar{\vartheta}_+$, and is a special case of the 1/2 BPS WL \tilde{W}_{fer} of the previous subsection.

A similar investigation can be performed for $\tilde{W}_{1/2}[\bar{\alpha}_I]$ with $|\alpha|^2 \neq 0$, which leads to the same conclusions. For $\bar{\alpha}_{3,4} = 0$ or $\bar{\alpha}_{1,2} = 0$ the resulting WLs are BPS and preserve sets of supercharges that are complementary to the ones already listed.

4 BPS WLs at quantum level

We now promote WLs to quantum operators and consider the problem of evaluating their vacuum expectation values (vev's). Similarly to what happens in the ABJ(M) theory, vev's of BPS WLs along a timelike straight line introduced in section 2.1 are constants that can be trivially normalized to one. Instead, non-trivial vev's can be obtained for circular BPS operators in Euclidean space defined in section 2.2. These can be evaluated by using localization techniques and the output turns out to be a non-trivial function of the couplings that

interpolates between the weak coupling result obtained via ordinary perturbation theory and the strong coupling result possibly obtained by holographic methods.

In this context it becomes particularly important to understand whether and how the classical cohomological \mathcal{Q} -equivalence between $W_{\rm bos}$ and $W_{\rm fer}$ discussed in subsection 2.1 gets promoted at quantum level. In fact, if we manage to prove that under suitable conditions the equivalence survives quantum corrections, then using the \mathcal{Q} charge to localize the functional integral we can conclude that the expectation value of the general parametric–dependent $W_{\rm fer}$ is identical to the one of $W_{\rm bos}$, making them effectively quantum equivalent and independent of the choice of the parameters.

In the case of ABJ(M) theory this problem has been extensively analysed [25–27] and it has been shown that its solution is deeply interconnected with the choice of a framing regularization for the perturbative definition of the WL. Precisely, the cohomological equivalence at quantum level leads to [13]

$$\langle W_{\text{bos}} \rangle_{1}^{(\text{ABJ(M)})} = \langle W_{\text{fer}} \rangle_{1}^{(\text{ABJ(M)})}$$
 (4.1)

where the subscript indicates that the identity holds only at framing one [2].

With a direct perturbative computation we are now going to show that, up to two loops, the problem of studying the quantum cohomological equivalence between circular W_{fer} and W_{bos} in generic $\mathcal{N}=2$ SCSM models can be mapped to the parallel problem in ABJ(M) theory. Using the speculations in [30], we can thus conclude that the classical cohomological equivalence can be promoted at quantum level, at least at two loops, if we work at framing one. Indeed, the two-loop result supports a stronger version of the \mathcal{Q} -equivalence, which we probe at three loops in subsection 4.2.

4.1 The perturbative analysis

In order to test the cohomological equivalence at quantum level, it is convenient to consider the difference between the expectation values of fermionic and bosonic WLs, order by order in perturbation theory. At classical level, the cohomological equivalence

$$W_{\text{fer}} - W_{\text{bos}} = QV \tag{4.2}$$

can be expanded as [13]

$$\sum_{n=1}^{\infty} \operatorname{Tr} \mathcal{P}(e^{-i \oint d\tau L_{\text{bos}}(\tau)} W_n) = \mathcal{Q}V$$
(4.3)

where the W_n expressions arise from the expansion of $e^{-i\oint (L_{\text{fer}}-L_{\text{bos}})} = e^{-i\oint (B+F)}$ inside the WL (see eq. (2.11)). The label n indicates the total number of scalar and fermion fields in the product. As follows from the explicit expressions of B and F in (2.31), at a given order n the W_n function is built up by p powers of B, which is quadratic in the scalar fields, and q powers of F, such that 2p + q = n (in particular, when n is odd the corresponding W_n contains an odd number of spinors F). As an example, we report the first few even

terms in (4.3)

$$W_{2} = -i \oint d\tau B(\tau) - \oint d\tau_{1>2} F(\tau_{1}) F(\tau_{2})$$

$$W_{4} = -\oint d\tau_{1>2} B(\tau_{1}) B(\tau_{2}) + i \oint d\tau_{1>2>3} \left[B(\tau_{1}) F(\tau_{2}) F(\tau_{3}) + F(\tau_{1}) B(\tau_{2}) F(\tau_{3}) + F(\tau_{1}) F(\tau_{2}) B(\tau_{3}) \right] + \oint d\tau_{1>2>3>4} F(\tau_{1}) F(\tau_{2}) F(\tau_{3}) F(\tau_{4})$$

$$W_{6} = i \oint d\tau_{1>2>3} B(\tau_{1}) B(\tau_{2}) B(\tau_{3}) + \oint d\tau_{1>2>3>4} \left[B(\tau_{1}) B(\tau_{2}) F(\tau_{3}) F(\tau_{4}) + B(\tau_{1}) F(\tau_{2}) B(\tau_{3}) F(\tau_{4}) + F(\tau_{1}) B(\tau_{2}) B(\tau_{3}) F(\tau_{4}) + F(\tau_{1}) B(\tau_{2}) F(\tau_{3}) B(\tau_{4}) + F(\tau_{1}) B(\tau_{2}) F(\tau_{3}) B(\tau_{4}) + F(\tau_{1}) F(\tau_{2}) B(\tau_{3}) F(\tau_{4}) \right] - i \oint d\tau_{1>2>3>4>5} \left[B(\tau_{1}) F(\tau_{2}) F(\tau_{3}) F(\tau_{4}) F(\tau_{5}) + F(\tau_{1}) B(\tau_{2}) F(\tau_{3}) F(\tau_{4}) F(\tau_{5}) + F(\tau_{1}) F(\tau_{2}) B(\tau_{3}) F(\tau_{4}) F(\tau_{5}) + F(\tau_{1}) F(\tau_{2}) F(\tau_{3}) F(\tau_{4}) B(\tau_{5}) \right] - \oint d\tau_{1>2>3>4>5} F(\tau_{1}) F(\tau_{2}) F(\tau_{3}) F(\tau_{4}) F(\tau_{5}) F(\tau_{6})$$

$$(4.4)$$

The structure of higher order terms clearly follows. Here $\oint d\tau_{1>2>\cdots>j}$ means integrals over the contour parameters τ_1,\ldots,τ_j satisfying $\tau_1>\tau_2>\cdots>\tau_j$.

It has been shown, for the first few orders in [13] and proved at all orders in [14], that at classical level the \mathcal{Q} -equivalence in (4.2) follows from a stronger set of identities that read

$$\operatorname{Tr} \mathcal{P}(e^{-i \oint d\tau L_{\text{bos}}(\tau)} W_n) = \mathcal{Q} V_n, \qquad n = 1, 2, \dots$$
(4.5)

that is, every single term in the summation (4.3) is cohomologically trivial for a suitable choice of the V_n function (see Appendix D in [14] for their explicit forms).

As already mentioned, in the ABJ(M) case when we move to quantum level and compute expectation values at framing one, no anomalies arise and relation (4.2) implies

$$\langle W_{\text{fer}} \rangle_1 = \langle W_{\text{bos}} \rangle_1$$
 (4.6)

which in turns can be written as

$$\sum_{n=1}^{\infty} \langle \operatorname{Tr} \mathcal{P}(e^{-i \oint d\tau L_{\text{bos}}(\tau)} W_{2n}) \rangle_{1} = 0$$
(4.7)

Here we have already neglected W_n for n odd, since strings of an odd number of F would have vanishing expectation values.

An interesting question is whether at quantum level and framing one, identities (4.5) remain true separately, as this would imply the non-trivial results

$$\langle \operatorname{Tr} \mathcal{P}(e^{-i \oint d\tau L_{\text{bos}}(\tau)} W_{2n}) \rangle_1 = 0, \qquad n = 1, 2, \dots$$
 (4.8)

We devote the rest of this section to the two-loop evaluation of (4.7) and the discussion of identities (4.8).

To this aim we assume CS levels k_a to be of the same order k, and all the WL parameters \bar{m}_i^{ab} and n_{ab}^i to be of order $1/\sqrt{k}$. In particular, this implies that a given W_{2m} term in (4.7) starts contributing at order $O(1/k^m)$. Therefore, in order to obtain the full result up to order $1/k^2$ we need to evaluate the terms in (4.7) involving correlators with W_2 and W_4 up to two loops and show that at framing one either their sum vanishes or they vanish separately. As a by–product, we also compute the two–loop expectation values of bosonic and general fermionic WLs at framing zero.

We focus on euclidean circle WLs (2.30) with connections (2.31). Moreover, for simplicity we restrict to quiver theories without adjoint or (anti-)fundamental matter. The more general case can be similarly worked out, but the results are more involved.

We organize our calculation as follows. We first compute W_{bos} at one and two loops, with relevant diagrams shown in figures 3 and 4. Then we consider equation (4.7) and evaluate correlators involving W_2 , W_4 as defined in (4.4). We compute the correlator with a W_2 insertion at one and two loops (figures 5 and 6) whereas the term with W_4 starts directly at two loops (figure 7). Writing the bosonic connection as $L_{\text{bos}} = A + S$ (and consequently $L_{\text{fer}} = A + S + B + F$), in these figures A insertions on the contour represent gauge connections, B and F insertions indicate quantities defined in (2.31), whereas S stays for the scalar bilinears which arise from $\sigma^{(a)}$, $a = 1, 2, \dots, n$ in L_{bos} when we use equations of motion (2.5).



Figure 3. One-loop Feynman diagram for $\langle W_{\rm bos} \rangle_f$. Wavy lines represent gauge fields.




Figure 4. Two-loop Feynman diagrams for $\langle W_{\text{bos}} \rangle_f$. Here S stays for the scalar bilinears which arise from $\sigma^{(a)}$, $a = 1, 2, \dots, n$ in L_{bos} when we use equations of motion (2.5).

The interesting observation is that a full fledged computation of all these Feynman diagrams is not required, as we can heavily rely on the results already present in the literature for a WL in pure CS theory [38] and 1/2 and 1/6 WLs in ABJ(M) theory [10–12, 25–27, 29, 30]. In fact, at the order we are working the internal vertices involving superpotential interactions do not play any role, and we have to deal only with pure gauge vertices and minimal couplings of matter fields to gauge vectors. Therefore, the difference



Figure 5. One-loop Feynman diagram for $\langle \text{Tr} \mathcal{P}(e^{-i \oint d\tau L_{\text{bos}}(\tau)} W_2) \rangle_f$. F indicates the fermionic quantities defined in (2.31).




Figure 6. Two-loop Feynman diagrams for $\langle \text{Tr} \mathcal{P}(e^{-i \oint d\tau L_{\text{bos}}(\tau)} W_2) \rangle_f$. Here B indicates the bosonic quantities defined in (2.31).




Figure 7. Two-loop Feynman diagrams for $\langle \text{Tr} \mathcal{P}(e^{-i \oint d\tau L_{\text{bos}}(\tau)} W_4) \rangle_f$.

between diagrams in ABJ(M) theory and any $\mathcal{N}=2$ SCSM model is only due to the different matter content, that is the different number of matter multiplets linking pairs of quiver nodes. It follows that at this order diagrams in the two theories differ only by the overall combinatorial factor, whereas the corresponding integrals are the same. Clearly, if we were to consider higher order corrections to W_n correlators the specific structure of the superpotential would generally kick in and different models might display different behaviours.

A long but straightforward evaluation of the diagrams leads to the results that can be found in appendix F. For each diagram we extract the color factor and the parametric dependence coming from the WL expansion, and indicate with italic capital letters the corresponding integrals that include both combinatorics and any other factors coming from the Feynman rules listed in appendix E. We stress that some of the fermionic diagrams can give rise to more than one integral, which differ by the order of the fermions along the contour. We present the results at framing f, meaning that f can be either zero or one.

Summing over all the contributions in each figure and using constraints (2.17), we obtain

for $W_{\rm bos}$

$$\frac{3}{a} = \sum_{a} \frac{N_a^2}{k_a} \mathcal{I}_3^{(f)}$$

$$\frac{4}{a} = \sum_{a,b} \frac{(N_{ab} + N_{ba})N_a^2 N_b}{k_a^2} (\mathcal{I}_{4(a)}^{(f)} + \mathcal{I}_{4(d)}^{(f)}) + \sum_{a} \frac{N_a^3}{k_a^2} (\mathcal{I}_{4(b)}^{(f)} + \mathcal{I}_{4(c)}^{(f)})$$

$$+ \sum_{a} \frac{N_a}{k_a^2} (-\mathcal{I}_{4(b)}^{(f)} + \mathcal{J}_{4(c)}^{(f)}) \tag{4.9}$$

for W_2

$$\underbrace{5} = \sum_{a,b} \bar{m}_{i}^{ab} n_{ba}^{i} N_{a} N_{b} \mathcal{I}_{5}^{(f)}$$

$$\underbrace{6} = \sum_{a,b} [\bar{m}_{i}^{ab} n_{ba}^{i} (\mathcal{I}_{6(a)}^{(f)} + \mathcal{I}_{6(b)}^{(f)} + \mathcal{I}_{6(c)}^{(f)} + \mathcal{I}_{6(d)}^{(f)})$$

$$+ \bar{m}_{i}^{ba} n_{ab}^{i} (-\mathcal{I}_{6(a)}^{(f)} + \mathcal{I}_{6(b)}^{(f)} + \mathcal{J}_{6(c)}^{(f)} + \mathcal{J}_{6(d)}^{(f)})] \frac{N_{a}^{2} N_{b}}{k_{c}} \tag{4.10}$$

for W_4

$$\begin{array}{l}
\boxed{7} = \sum_{a,b,c} \left[\bar{m}_{i}^{ab} n_{ba}^{i} \bar{m}_{j}^{ac} n_{ca}^{j} \left(\mathcal{I}_{7(\mathbf{a})}^{(f)} + \mathcal{I}_{7(\mathbf{b})}^{(f)} \right) \\
+ \bar{m}_{i}^{ba} n_{ab}^{i} \bar{m}_{j}^{ca} n_{ac}^{j} \left(\mathcal{I}_{7(\mathbf{a})}^{(f)} + \mathcal{J}_{7(\mathbf{b})}^{(f)} \right) \right] N_{a} N_{b} N_{c}
\end{array} (4.11)$$

The exact expression of the \mathcal{I} , \mathcal{J} integrals can be found in appendix F. In particular, from their explicit expression it follows that

$$\mathcal{I}_{6(a)}^{(f)} = -\frac{1}{\pi} \mathcal{I}_{4(d)}^{(f)} \qquad \mathcal{I}_{7(a)}^{(f)} = \frac{1}{4\pi^2} \mathcal{I}_{4(d)}^{(f)}$$
(4.12)

We stress that these results are common to any $N \geq 2$ SCSM theory, ABJ(M) models included. Therefore, the explicit values of the integrals can be extracted from the literature on the pure Chern–Simons theory [38] and ABJ(M) theory [10–12].

At generic framing it is known that [30]

$$\mathcal{I}_{3}^{(f)} = -\pi i f, \quad \mathcal{I}_{4(a)}^{(f)} + \mathcal{I}_{4(d)}^{(f)} = \frac{\pi^{2}}{4}, \quad \mathcal{I}_{4(b)}^{(f)} = -\frac{\pi^{2}}{6}, \quad \mathcal{I}_{4(c)}^{(f)} = -\frac{\pi^{2}}{2} f^{2}, \quad \mathcal{J}_{4(c)}^{(f)} = 0 \quad (4.13)$$

whereas the ones appearing in (4.10, 4.11) are explicitly known only at framing zero [25–27, 30]

$$\mathcal{I}_{5}^{(0)} = \mathcal{I}_{6(a)}^{(0)} = \mathcal{I}_{6(b)}^{(0)} = \mathcal{I}_{6(d)}^{(0)} = \mathcal{I}_{6(d)}^{(0)} = 0, \quad \mathcal{I}_{6(c)}^{(0)} = -\mathcal{I}_{6(c)}^{(0)} = -\frac{\pi}{2}, \quad \mathcal{I}_{7(b)}^{(0)} = \mathcal{I}_{7(b)}^{(0)} = \frac{3}{32} \tag{4.14}$$

However, it can be checked numerically that both $\mathcal{I}_{4(\mathbf{a})}^{(f)}$ and $\mathcal{I}_{4(\mathbf{d})}^{(f)}$ are framing independent. Therefore, since at framing zero $\mathcal{I}_{4(\mathbf{d})}^{(0)} = \mathcal{I}_{6(\mathbf{a})}^{(0)} = \mathcal{I}_{7(\mathbf{a})}^{(0)} = 0$, at generic framing we obtain

$$\mathcal{I}_{4(a)}^{(f)} = \frac{\pi^2}{4}, \quad \mathcal{I}_{4(d)}^{(f)} = \mathcal{I}_{6(a)}^{(f)} = \mathcal{I}_{7(a)}^{(f)} = 0$$
 (4.15)

Using redefinitions (B.12) for the parameters and choosing $\beta^I = \alpha^I/|\alpha|^2$, $\bar{\gamma}_I = \delta^I = 0$, expressions (4.10–4.11) reduce to W_2, W_4 terms for the expansion of $\langle W_{1/2} - W_{\rm bos} \rangle_f$ in ABJ(M) theory. Therefore, using the arguments and the well–based speculations of [30] for 1/2 BPS WLs in ABJ(M) theory, we obtain that at framing one

$$\mathcal{I}_{5}^{(1)} = \mathcal{I}_{6(b)}^{(1)} + \mathcal{I}_{6(c)}^{(1)} = \mathcal{I}_{6(b)}^{(1)} + \mathcal{J}_{6(c)}^{(1)} = \mathcal{I}_{6(d)}^{(1)} = \mathcal{J}_{6(d)}^{(1)} = \mathcal{I}_{7(b)}^{(1)} = \mathcal{J}_{7(b)}^{(1)} = 0$$
(4.16)

Therefore, we can conclude that at framing one diagrams 5–7 are identically vanishing

$$5 = 6 = 7 = 0 \quad \text{for } f = 1$$
 (4.17)

and the cohomological equivalence between W_{bos} and W_{fer} holds for any $\mathcal{N} \geq 2$ SCSM theory, i.e.

$$\langle W_{\text{fer}} \rangle_1 - \langle W_{\text{bos}} \rangle_1 = 0$$
 up to two loops (4.18)

Actually, we find a much stronger result. In fact, results (4.17) imply that

$$\langle \operatorname{Tr} \mathcal{P}(e^{-i \oint d\tau L_{\text{bos}}(\tau)} W_2) \rangle_1 = \langle \operatorname{Tr} \mathcal{P}(e^{-i \oint d\tau L_{\text{bos}}(\tau)} W_4) \rangle_1 = 0 \tag{4.19}$$

separately. This is nothing but eq. (4.8) that we expect to hold quantum mechanically as a consequence of the classical relations in (4.5).

We stress that the cohomological equivalence up to two loops and framing one and its stronger version are still valid when there are adjoint and fundamental matter fields in the theory.

Specializing these findings to the ABJ(M) theory, we find that $W_{\rm bos}$ is cohomological equivalent not only to the 1/2 BPS fermionic operator [13], but also to generic 1/6 BPS fermionic WLs introduced in [16, 17], at least at the order we are working. Similarly, in $\mathcal{N}=4$ circular quiver SCSM theory with alternating levels, the fermionic 1/4 BPS WLs introduced in [18] are also quantum mechanically cohomological equivalent to the bosonic 1/4 BPS WL up to two loops.

Exploiting results (4.13) and summing up all the contributions in (4.9) we obtain the general result for W_{bos} in $\mathcal{N}=2$ SCSM models up to two loops and generic framing

$$\langle W_{\text{bos}} \rangle_f = \sum_a \left\{ N_a - \frac{\pi i f N_a^2}{k_a} + \frac{\pi^2 [-(3f^2 + 1)N_a^3 + N_a]}{6k_a^2} \right\}$$

$$+ \sum_{a,b} \frac{\pi^2 (N_{ab} + N_{ba})N_a^2 N_b}{4k_a^2} + O\left(\frac{1}{k^3}\right)$$
(4.20)

Similarly, we can now use results (4.14) in eq. (4.10-4.11) and obtain the two-loop expression

$$\langle W_{\text{fer}} - W_{\text{bos}} \rangle_{0} = -\sum_{a,b} (\bar{m}_{i}^{ab} n_{ba}^{i} - \bar{m}_{i}^{ba} n_{ab}^{i}) \frac{\pi N_{a}^{2} N_{b}}{2k_{a}}$$

$$+ \sum_{a,b,c} (\bar{m}_{i}^{ab} n_{ba}^{i} \bar{m}_{j}^{ac} n_{ca}^{j} + \bar{m}_{i}^{ba} n_{ab}^{i} \bar{m}_{j}^{ca} n_{ac}^{j}) \frac{3N_{a} N_{b} N_{c}}{32} + O\left(\frac{1}{k^{3}}\right)$$

$$(4.21)$$

In particular, this shows that at framing zero cohomological equivalence is in general broken.

By combining (4.21) with (4.20) evaluated at f = 0 we also obtain the framing zero result for W_{fer} up to two loops

$$\langle W_{\text{fer}} \rangle_{0} = \sum_{a} \left[N_{a} + \frac{\pi^{2}(-N_{a}^{3} + N_{a})}{6k_{a}^{2}} \right]$$

$$+ \sum_{a,b} \left[\frac{\pi^{2}(N_{ab} + N_{ba})N_{a}^{2}N_{b}}{4k_{a}^{2}} - (\bar{m}_{i}^{ab}n_{ba}^{i} - \bar{m}_{i}^{ba}n_{ab}^{i}) \frac{\pi N_{a}^{2}N_{b}}{2k_{a}} \right]$$

$$+ \sum_{a,b,c} (\bar{m}_{i}^{ab}n_{ba}^{i}\bar{m}_{j}^{ac}n_{ca}^{j} + \bar{m}_{i}^{ba}n_{ab}^{i}\bar{m}_{j}^{ca}n_{ac}^{j}) \frac{3N_{a}N_{b}N_{c}}{32} + O\left(\frac{1}{k^{3}}\right)$$

$$(4.22)$$

The result displays a non-trivial parametric dependence on the fermion couplings, and thus it is different for different WLs.

More interestingly, at framing one, independently of the choice of the parametric couplings we find

$$\langle W_{\text{bos}} \rangle_{1} = \langle W_{\text{fer}} \rangle_{1} = \sum_{a} \left\{ N_{a} - \frac{\pi i N_{a}^{2}}{k_{a}} + \frac{\pi^{2} [-4N_{a}^{3} + N_{a}]}{6k_{a}^{2}} \right\} + \sum_{a,b} \frac{\pi^{2} (N_{ab} + N_{ba}) N_{a}^{2} N_{b}}{4k_{a}^{2}} + O\left(\frac{1}{k^{3}}\right)$$
(4.23)

In particular, both $\langle W_{\text{bos}} \rangle_1$ and $\langle W_{\text{fer}} \rangle_1$ can in principle be evaluated by using localization techniques [2]. The result, expanded at second order, should match (4.23).

Specializing our results to the ABJ(M) theory and normalizing the operators properly, we find

$$\frac{\langle W_{\text{bos}} \rangle_f}{N_1 + N_2} = 1 - \frac{\pi i f(N_1 - N_2)}{k} + \frac{\pi^2}{6k^2} \Big[- (3f^2 + 1)(N_1^2 + N_2^2) \\
+ (3f^2 + 7)N_1 N_2 + 1 \Big] + O\Big(\frac{1}{k^3}\Big) \\
\frac{\langle W_{\text{fer}} - W_{\text{bos}} \rangle_0}{N_1 + N_2} = \frac{\pi^2 N_1 N_2}{2k^2} \Big\{ 3[(\bar{\alpha}_I \beta^I)^2 + (\bar{\gamma}_I \delta^I)^2] - 4(\bar{\alpha}_I \beta^I + \bar{\gamma}_I \delta^I) \Big\} + O\Big(\frac{1}{k^3}\Big) \\
\frac{\langle W_{\text{fer}} \rangle_0}{N_1 + N_2} = 1 + \frac{\pi^2}{6k^2} \Big\{ -N_1^2 - N_2^2 + \Big\{ 9[(\bar{\alpha}_I \beta^I)^2 + (\bar{\gamma}_I \delta^I)^2] \\
- 12(\bar{\alpha}_I \beta^I + \bar{\gamma}_I \delta^I) + 7 \Big\} N_1 N_2 + 1 \Big\} + O\Big(\frac{1}{k^3}\Big) \tag{4.24}$$

4.2 A conjecture

As already stressed, the two-loop computation of the previous section displays an important feature which could lead to far reaching consequences, if confirmed at higher orders. In fact, using in (4.7) the shorthand $\langle \text{Tr} \mathcal{P}(e^{-i\oint d\tau L_{\text{bos}}(\tau)}W_{2n})\rangle_1 \equiv \langle W_{2n}\rangle_1$, up to order $1/k^2$ we find not only $\langle W_2\rangle_1 + \langle W_4\rangle_1 = 0$, but also $\langle W_2\rangle_1 = 0$ and $\langle W_4\rangle_1 = 0$, separately. At order 1/k this is trivially realized, since only the W_2 correlator can be constructed, but at order $1/k^2$ this is a non-trivial statement.

Repeating the analysis of [30], this property can be understood as follows. If we temporarily consider WL operators with non-trivial winding m, as a rule of thumb it can be argued that, independently of framing, a generic perturbative diagram has a polynomial dependence on m, whose leading power is $m^{2[r/2]}$ where r is the number of contour insertions, that is the number of contour integrations. Up to two loops this property has been tested explicitly in [30]. Focusing on the two-loop diagrams in figure 6 and 7 contributing to the W_2 and W_4 correlators, at framing one the only potentially non-vanishing contributions come from diagrams 6(b), 6(c) and 7(b). These diagrams produce contributions to W_2 and W_4 correlators that display different leading powers in the winding number. Precisely, contributions to $\langle W_2 \rangle$ go as m^2 , while the ones contributing to $\langle W_4 \rangle$ go as m^4 . Thus, at non-trivial winding, cohomological equivalence in ABJ(M) forces the two correlators to vanish separately. We note that at this order what plays a crucial role in assigning different powers of m to W_2 and W_4 correlators is the fact that scalar eye-like diagrams 6(a) and 7(a)vanish. If this were not the case, since they appear in both correlators the winding argument would get spoiled. However, the eye-like diagram is vanishing at framing zero by analytical continuation in dimensional regularization, and is shown to be framing-independent numerically.

Reinforced by this preliminary result, we can then reasonably believe that a similar pattern may survive at higher orders, so allowing to conjecture that $\langle W_{2n} \rangle_1 = 0$ at any order in perturbation theory, with the definition of W_{2n} in (4.3).

The validity of these identities implies strong constraints on the (unknown) integrals at a given perturbative order. As a non-trivial example, we consider (4.8) at three loops. At this order we should prove that $\langle W_2 \rangle_1$, $\langle W_4 \rangle_1$ and $\langle W_6 \rangle_1$ vanish separately.

In particular, focusing on $\langle W_6 \rangle_1$ the corresponding contributions arise from diagrams in figure 8. We note that these diagrams do not contain superpotential vertices, and are then common to all $\mathcal{N} \geq 2$ SCSM theories, ABJ(M) theory included.

The corresponding analytic expressions read

$$\begin{aligned}
& = \sum_{a,b,c,d} \left[\bar{m}_{i}^{ab} n_{ba}^{i} \bar{m}_{j}^{ac} n_{ca}^{j} \bar{m}_{k}^{ad} n_{da}^{k} (\mathcal{I}_{8(a)}^{(f)} + \mathcal{I}_{8(c)}^{(f)}) \right. \\
& + \bar{m}_{i}^{ba} n_{ab}^{i} \bar{m}_{j}^{ca} n_{ac}^{j} \bar{m}_{k}^{da} n_{ad}^{k} (\mathcal{I}_{8(a)}^{(f)} + \mathcal{J}_{8(c)}^{(f)}) \\
& + \bar{m}_{i}^{ab} n_{ba}^{i} \bar{m}_{j}^{ac} n_{ca}^{j} \bar{m}_{k}^{db} n_{bd}^{k} (\mathcal{I}_{8(b)}^{(f)} + \mathcal{K}_{8(c)}^{(f)}) \right] N_{a} N_{b} N_{c} N_{d} \\
& + \sum_{a,b} (\bar{m}_{i}^{ab} n_{ba}^{i})^{3} N_{a} N_{b} (2 \mathcal{I}_{8(a)}^{(f)} + \mathcal{J}_{8(b)}^{(f)} + \mathcal{L}_{8(c)}^{(f)})
\end{aligned} \tag{4.25}$$




Figure 8. Three-loop Feynman diagrams for $\langle \text{Tr} \mathcal{P}(e^{-i \oint d\tau L_{\text{bos}}(\tau)} W_6) \rangle_f$.

where the explicit expression of the integrals can be found in appendix F (see eq. (F.12)). It follows that conjecture (4.8) for n=3 forces the framing-one integrals to satisfy the non-trivial identities

$$\mathcal{I}_{8(a)}^{(1)} + \mathcal{I}_{8(c)}^{(1)} = \mathcal{I}_{8(a)}^{(1)} + \mathcal{J}_{8(c)}^{(1)} = \mathcal{I}_{8(b)}^{(1)} + \mathcal{K}_{8(c)}^{(1)} = 2\mathcal{I}_{8(a)}^{(1)} + \mathcal{J}_{8(b)}^{(1)} + \mathcal{L}_{8(c)}^{(1)} = 0$$
 (4.26)

Again we expect to be able to refine this set of constraints by a direct analysis of the explicit expression of the integrals. Moreover, it would be important to check these relations by evaluating explicitly all the integrals. This is a quite hard task which goes beyond the scopes of the present paper.

We conclude with an important observation. If rigorously proved, identities (4.8) would have strong implications for the defect CFT defined on the bosonic Wilson contour. In fact, for any local or nonlocal operator \mathcal{O} localized on the contour of the WL, the expression

$$\langle\!\langle \mathcal{O} \rangle\!\rangle_f \equiv \frac{\langle \text{Tr} \mathcal{P}(e^{-i \oint d\tau L_{\text{bos}}(\tau)} \mathcal{O}) \rangle_f}{\langle \text{Tr} \mathcal{P} e^{-i \oint d\tau L_{\text{bos}}(\tau)} \rangle_f}$$
(4.27)

defines correlations functions in the one-dimensional defect CFT on the circle. Interestingly, the defect CFT would depend on the framing we choose. In this language identities (4.8) would read

$$\langle \langle W_{2n} \rangle \rangle_1 = 0, \quad n = 1, 2, \cdots$$
 (4.28)

These equations would then impose strong constraints on the defect CFT with interesting implications on the corresponding boostrap program.

5 Conclusions

In this paper we have investigated the class of 1/2 BPS WLs in $\mathcal{N}=2$ SCSM theories featured by constant parametric couplings to matter scalars and fermions. Beyond the bosonic

WL that contains only couplings to scalars, we have found an infinite family of fermionic WLs. In general, the corresponding connections cannot be decomposed as double—node connections and cannot be interpreted as superconnections of a supergroup. Nonetheless, the new fermionic 1/2 BPS WLs are classically cohomologically equivalent to the bosonic 1/2 BPS WLs.

In order to exemplify the general results, we have revisited the case of ABJ(M) theory and $\mathcal{N}=4$ orbifold ABJ(M) in $\mathcal{N}=2$ language, and studied in details the $\mathcal{N}=2$ orbifold ABJM theory. In $\mathcal{N}=4$ and $\mathcal{N}=2$ orbifold ABJM theories, some of the newly found BPS WLs can be obtained from the orbifold decomposition of the 1/2 BPS WLs in ABJM theory. For these operators we have identified the corresponding gravity duals by direct orbifolding the brane configurations dual to 1/2 BPS WLs in ABJM theory. Whether gravity duals of more general BPS WLs can be identified is an important open question that requires further investigation.

We have discussed the cohomological equivalence of the fermionic and bosonic BPS WLs at quantum level by studying their expectation values up to two loops. In fact, since at this order the superpotential couplings do not enter, it happens that the arguments and well-based speculations of [30] lead to cohomological equivalence in any $\mathcal{N}=2$ CS-matter models as in ABJ(M) theory. We have further conjectured that in general the cohomological equivalence may occur in the stronger version of eq. (4.8). Since this condition would have far–reaching implications for the defect CFT defined on the bosonic WL contour, we plan to further investigate it in the future.

For the ordinary WLs along closed curves in gauge theories, we can take the trace in any representation of the gauge group. In four dimensional $\mathcal{N}=4$ super Yang-Mills theory, the BPS WLs in higher dimensional representations have elegant holographic description in terms of D-branes [39–43] or bubbling geometries [44–46]. For 1/2 BPS fermionic WLs in ABJ(M) theory, the trace can also be taken in higher dimensional representations of the supergroup $U(N_1|N_2)$ [47]. As we already stressed, in general the connection L_{fer} constructed here is not a superconnection with respect to a supergroup. This raises the question whether, for these WLs along closed curves, we can take the trace in some analog of higher dimensional representations mentioned above. We would like to leave this interesting question for further work.

Acknowledgments

We would like to thank Jin-Peng An, Nadav Drukker, Si Li, Hong Lü, Zhao-Long Wang for helpful discussions. H.O. gratefully acknowledges the kind hospitality of Tianjin University. J.Z. would like to thank Peking University and Tianjin University for hospitality, where part of the work has been done. The work of A.M., S.P., and J.Z. has been supported in part by Italian Ministero dell'Istruzione, Università e Ricerca (MIUR), and Istituto Nazionale di Fisica Nucleare (INFN) through the "Gauge Theories, Strings, Supergravity" (GSS) research projects. The work of H.O. and J.–B.W. has been supported in part by the National Natural Science Foundation of China, Grant No. 11575202. The work of J.Z. has been supported by Fondazione Cariplo and Regione Lombardia, Grant No. 2015-1253.

A Spinor conventions in three dimensions

In this Appendix we collect our spinor conventions, both in Minkowski and Euclidean signatures.

A.1 Minkowski spacetime

In three–dimensional Minkowski spacetime we follow the convention in [33, 48], where the reader can find more details. We use coordinates $x^{\mu} = (x^0, x^1, x^2)$ and metric $\eta_{\mu\nu} = \text{diag}(-++)$. We choose gamma matrices

$$\gamma^{\mu}_{\alpha}{}^{\beta} = (i\sigma^2, \sigma^1, \sigma^3) \tag{A.1}$$

where $\sigma^{1,2,3}$ are the Pauli matrices. They satisfy $\gamma^{\mu}\gamma^{\nu} = \eta^{\mu\nu} + \varepsilon^{\mu\nu\rho}\gamma_{\rho}$ with $\varepsilon^{012} = 1$. Note the spinor index $\alpha = 1, 2$.

The charge conjugate of spinors is defined as

$$\bar{\theta}_{\alpha} = \theta_{\alpha}^{*}, \quad \bar{\theta}_{\alpha}^{*} = \theta_{\alpha}$$
 (A.2)

The spinor indices are raised and lowered as

$$\theta^{\alpha} = \varepsilon^{\alpha\beta}\theta_{\beta}$$
 , $\theta_{\alpha} = \varepsilon_{\alpha\beta}\theta^{\beta}$ (A.3)

where $\varepsilon^{12} = -\varepsilon_{12} = 1$. We also define the shorthand notation

$$\theta\psi = \theta^{\alpha}\psi_{\alpha}, \quad (\gamma^{\mu}\theta)_{\alpha} = \gamma^{\mu}_{\alpha}{}^{\beta}\theta_{\beta}, \quad (\theta\gamma^{\mu})^{\alpha} = \theta^{\beta}\gamma^{\mu}_{\beta}{}^{\alpha}, \quad \theta\gamma^{\mu}\psi = \theta^{\alpha}\gamma^{\mu}_{\alpha}{}^{\beta}\psi_{\beta}$$
 (A.4)

On the straight line $x^{\mu} = (\tau, 0, 0)$, we introduce the bosonic spinors

$$u_{\pm \alpha} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1\\ \mp i \end{pmatrix}, \quad u_{\pm}^{\alpha} = \frac{1}{\sqrt{2}} (\mp i, -1)$$
 (A.5)

and decompose a generic spinor as

$$\theta_{\alpha} = u_{+\alpha}\theta_{-} + u_{-\alpha}\theta_{+} \tag{A.6}$$

where θ_{\pm} are one–component Grassmann numbers. The product of two spinors now reads

$$\theta \psi = i \left(\theta_+ \psi_- - \theta_- \psi_+ \right) \tag{A.7}$$

A.2 Euclidean space

In Euclidean space we follow the spinor conventions of [48]. We use coordinates $x^{\mu} = (x^1, x^2, x^3)$ and metric $\delta_{\mu\nu} = \text{diag}(+++)$. We choose gamma matrices

$$\gamma^{\mu}_{\alpha}{}^{\beta} = (-\sigma^2, \sigma^1, \sigma^3) \tag{A.8}$$

that satisfy $\gamma^{\mu}\gamma^{\nu} = \delta^{\mu\nu} + i\varepsilon^{\mu\nu\rho}\gamma_{\rho}$ with $\varepsilon^{123} = 1$.

The spinor indices are raised and lowered as

$$\theta^{\alpha} = \varepsilon^{\alpha\beta}\theta_{\beta}, \quad \theta_{\alpha} = \varepsilon_{\alpha\beta}\theta^{\beta} \tag{A.9}$$

where $\varepsilon^{12} = -\varepsilon_{12} = 1$.

In Euclidean space $\bar{\theta}$ and θ are independent spinors. For a general spinor θ one can define θ^{\dagger} that satisfies

$$\theta_{\alpha}^* = \theta^{\dagger \alpha}, \quad \theta^{\alpha *} = -\theta_{\alpha}^{\dagger}, \quad \theta^{\dagger \alpha *} = \theta_{\alpha}, \quad \theta_{\alpha}^{\dagger *} = -\theta^{\alpha}$$
 (A.10)

Formally one has $\theta^{\dagger\dagger} = -\theta$.

We use the shorthand notation

$$\theta\psi = \theta^{\alpha}\psi_{\alpha}, \quad (\gamma^{\mu}\theta)_{\alpha} = \gamma^{\mu\beta}_{\alpha}\theta_{\beta}, \quad (\theta\gamma^{\mu})^{\alpha} = \theta^{\beta}\gamma^{\mu\alpha}_{\beta}, \quad \theta\gamma^{\mu}\psi = \theta^{\alpha}\gamma^{\mu\beta}_{\alpha}\psi_{\beta}$$
 (A.11)

For the circle $x^{\mu} = (\cos \tau, \sin \tau, 0)$, we choose the $u_{\pm \alpha}$ spinors as

$$u_{+\alpha} = \frac{1}{\sqrt{2}} \begin{pmatrix} e^{-\frac{i\tau}{2}} \\ e^{\frac{i\tau}{2}} \end{pmatrix}, \quad u_{-\alpha} = \frac{i}{\sqrt{2}} \begin{pmatrix} -e^{-\frac{i\tau}{2}} \\ e^{\frac{i\tau}{2}} \end{pmatrix}$$
$$u_{+}^{\alpha} = \frac{1}{\sqrt{2}} \left(e^{\frac{i\tau}{2}}, -e^{-\frac{i\tau}{2}} \right), \quad u_{-}^{\alpha} = \frac{i}{\sqrt{2}} \left(e^{\frac{i\tau}{2}}, e^{-\frac{i\tau}{2}} \right)$$
(A.12)

We decompose spinors in Euclidean space formally in the same way as in Minkowski spacetime, see eq. (A.6). The product of two spinors is still given in (A.7). However, the u_{\pm} spinors are defined differently, and in particular in Euclidean space they are not constant.

B BPS WLs in ABJ(M) theory

In this appendix, we review 1/6 and 1/2 BPS WLs in ABJ(M) theory, including both line WLs in Minkowski spacetime and circle WLs in Euclidean space [10–13, 16, 17]. We also reproduce these known WLs using the general construction of sections 2.1 and 2.2 valid for generic $\mathcal{N}=2$ SCSM theories. This requires a notational translation from what we call ABJ(M) notations to $\mathcal{N}=2$ notations, which we describe in details. For 1/2 BPS line WLs in Minkowski spacetime we also review the construction of (anti–)M2–brane duals in M–theory.

B.1 1/6 BPS line WLs in Minkowski spacetime

The $U(N_1)_k \times U(N_2)_{-k}$ ABJ(M) theory [6, 7, 49] is usually written in manifest SU(4) R-symmetry notations. Gauge fields A_{μ} , B_{μ} corresponding to the two nodes of the quiver diagram in figure 9(a) are linked by ϕ_I , ψ^I , I=1,2,3,4 bosonic and fermionic matter fields in the bifundamental representation of the gauge group and in the fundamental of the R-symmetry group. The SUSY parameters are θ^{IJ} , $\bar{\theta}_{IJ}$, $\bar{\vartheta}_{IJ}$ with

$$\theta^{IJ} = -\theta^{JI}, \quad (\theta^{IJ})^* = \bar{\theta}_{IJ}, \quad \bar{\theta}_{IJ} = \frac{1}{2} \epsilon_{IJKL} \theta^{KL}$$

$$\vartheta^{IJ} = -\vartheta^{JI}, \quad (\vartheta^{IJ})^* = \bar{\vartheta}_{IJ}, \quad \bar{\vartheta}_{IJ} = \frac{1}{2} \epsilon_{IJKL} \vartheta^{KL}$$
(B.1)




Figure 9. The quiver diagram of ABJ(M) theory in (a) ABJM notations and (b) $\mathcal{N}=2$ notations.

Here θ^{IJ} , $\bar{\theta}_{IJ}$ are related to Poincaré supercharges, and ϑ^{IJ} , $\bar{\vartheta}_{IJ}$ are related to superconformal charges.

In Minkowski spacetime, the bosonic 1/6 BPS WL along the line $x^{\mu} = (\tau, 0, 0)$ is defined as in (2.8) with connection matrix [10–12]

$$L_{\text{bos}} = \text{diag}(A_0 + \frac{2\pi}{k} R^I{}_J \phi_I \bar{\phi}^J, B_0 + \frac{2\pi}{k} R^I{}_J \bar{\phi}^J \phi_I)$$

$$R^I{}_J = \text{diag}(-1, -1, 1, 1)$$
(B.2)

Fermionic 1/6 BPS WL W_{fer} can also be constructed as in (2.10), which correspond to the superconnection [16, 17]

$$L_{\text{fer}} = \begin{pmatrix} A_0 + \frac{2\pi}{k} U^I_{J} \phi_I \bar{\phi}^J & \sqrt{\frac{4\pi}{k}} (\bar{\alpha}_I \psi_+^I + \bar{\gamma}_I \psi_-^I) \\ \sqrt{\frac{4\pi}{k}} (\bar{\psi}_{I-} \beta^I - \bar{\psi}_{I+} \delta^I) & B_0 + \frac{2\pi}{k} U^I_{J} \bar{\phi}^J \phi_I \end{pmatrix}$$
(B.3)

where the couplings to matter are featured by constant parameters

$$U^{I}_{J} = \begin{pmatrix} -1 + 2\beta^{2}\bar{\alpha}_{2} & -2\beta^{1}\bar{\alpha}_{2} \\ -2\beta^{2}\bar{\alpha}_{1} & -1 + 2\beta^{1}\bar{\alpha}_{1} \\ & 1 - 2\delta^{4}\bar{\gamma}_{4} & 2\delta^{3}\bar{\gamma}_{4} \\ & 2\delta^{4}\bar{\gamma}_{3} & 1 - 2\delta^{3}\bar{\gamma}_{3} \end{pmatrix}$$

$$\bar{\alpha}_{I} = (\bar{\alpha}_{1}, \bar{\alpha}_{2}, 0, 0), \quad \beta^{I} = (\beta^{1}, \beta^{2}, 0, 0), \quad \bar{\gamma}_{I} = (0, 0, \bar{\gamma}_{3}, \bar{\gamma}_{4}), \quad \delta^{I} = (0, 0, \delta^{3}, \delta^{4})$$
(B.4)

satisfying the BPS constraints

$$\bar{\alpha}_{1,2}\delta^{3,4} = \bar{\gamma}_{3,4}\beta^{1,2} = 0 \tag{B.5}$$

The corresponding preserved supercharges are $\theta_+^{12},~\theta_-^{34},~\vartheta_+^{12},~\vartheta_-^{34}$.

Similarly, along the line $x^{\mu}=(\tau,0,0)$ we can define $\tilde{W}_{\rm bos},\,\tilde{W}_{\rm fer}$ operators with connections

$$\tilde{L}_{\text{bos}} = \text{diag} \left(A_0 - \frac{2\pi}{k} R^I{}_J \phi_I \bar{\phi}^J, B_0 - \frac{2\pi}{k} R^I{}_J \bar{\phi}^J \phi_I \right)
\tilde{L}_{\text{fer}} = \begin{pmatrix} A_0 - \frac{2\pi}{k} U^I{}_J \phi_I \bar{\phi}^J & \sqrt{\frac{4\pi}{k}} (\bar{\alpha}_I \psi_-^I + \bar{\gamma}_I \psi_+^I) \\ \sqrt{\frac{4\pi}{k}} (-\bar{\psi}_{I+} \beta^I + \bar{\psi}_{I-} \delta^I) & B_0 - \frac{2\pi}{k} U^I{}_J \bar{\phi}^J \phi_I \end{pmatrix}$$
(B.6)

where the constant parameters are the same as in (B.2), (B.4) and satisfy the same constraints (B.5). The preserved supercharges θ_{-}^{12} , θ_{+}^{34} , ϑ_{-}^{12} , ϑ_{+}^{34} are complementary to the ones preserved by W_{bos} , W_{fer} .

 W_{bos} and W_{fer} operators are cohomologically equivalent, and that is their difference is $\mathcal{Q}(\text{something})$, where \mathcal{Q} is a supercharge preserved by both WLs. Similarly, one can prove that \tilde{W}_{bos} , \tilde{W}_{fer} are cohomologically equivalent.

In order to make contact with the general WL construction presented in the main text, we rewrite the ABJ(M) theory in $\mathcal{N}=2$ superspace formalism. This is obtained by identifying $\mathcal{N}=6$ and $\mathcal{N}=2$ SUSY parameters as

$$\theta^{12} = \bar{\theta}_{34} = \theta$$
 $\theta^{34} = \bar{\theta}_{12} = \bar{\theta}$ $\theta^{12} = \bar{\theta}_{34} = \theta$ $\theta^{34} = \bar{\theta}_{12} = \bar{\theta}$ (B.7)

The ABJ(M) theory in $\mathcal{N}=2$ superspace formalism has only SU(2) R–symmetry invariance manifest and corresponds to the quiver diagram in figure 9(b). The two sets of fields are related by

$$A_{\mu} = A_{\mu}^{(1)}, \quad B_{\mu} = A_{\mu}^{(2)}, \qquad \phi_{I} = (Z_{1}, Z_{2}, \bar{Z}^{3}, \bar{Z}^{4}), \quad \psi^{I} = (-\zeta_{2}, \zeta_{1}, -\bar{\zeta}^{4}, \bar{\zeta}^{3})$$
 (B.8)

Using the general construction of WL operators in section 2 and adapting it to the ABJ(M) case, we find that the $W_{\rm bos}$ and $W_{\rm fer}$ operators along the line $x^{\mu}=(\tau,0,0)$ and preserving supercharges θ_+ , $\bar{\theta}_-$, ϑ_+ , $\bar{\vartheta}_-$, have connections (2.19), with the gauge auxiliary fields given by

$$\sigma^{(1)} = \frac{2\pi}{k} (Z_1 \bar{Z}^1 + Z_2 \bar{Z}^2 - \bar{Z}^3 Z_3 - \bar{Z}^4 Z_4)$$

$$\sigma^{(2)} = \frac{2\pi}{k} (\bar{Z}^1 Z_1 + \bar{Z}^2 Z_2 - Z_3 \bar{Z}^3 - Z_4 \bar{Z}^4)$$
(B.9)

and matrix couplings

$$\bar{M}_{Z} = \begin{pmatrix} \bar{m}^{1}Z_{1} + \bar{m}^{2}Z_{2} \\ \bar{m}^{3}Z_{3} + \bar{m}^{4}Z_{4} \end{pmatrix}$$

$$N_{\bar{Z}} = \begin{pmatrix} n_{3}\bar{Z}^{3} + n_{4}\bar{Z}^{4} \\ n_{1}\bar{Z}^{1} + n_{2}\bar{Z}^{2} \end{pmatrix}$$

$$\bar{M}_{\zeta} = \begin{pmatrix} \bar{m}^{1}\zeta_{1+} + \bar{m}^{2}\zeta_{2+} \\ \bar{m}^{3}\zeta_{3+} + \bar{m}^{4}\zeta_{4+} \end{pmatrix}$$

$$N_{\bar{\zeta}} = \begin{pmatrix} n_{3}\bar{\zeta}_{3}^{3} + n_{4}\bar{\zeta}_{4}^{4} \\ n_{1}\bar{\zeta}_{1}^{1} + n_{2}\bar{\zeta}_{2}^{2} \end{pmatrix}$$
(B.10)

The coupling parameters satisfy the BPS constraints

$$\bar{m}^{1,2}\bar{m}^{3,4} = n_{1,2}n_{3,4} = 0 \tag{B.11}$$

It is now easy to verify that redefining the parameters as

$$\bar{m}^{1} = \sqrt{\frac{4\pi}{k}}\bar{\alpha}_{2}, \quad \bar{m}^{2} = -\sqrt{\frac{4\pi}{k}}\bar{\alpha}_{1}, \quad \bar{m}^{3} = -\sqrt{\frac{4\pi}{k}}\delta^{4}, \quad \bar{m}^{4} = \sqrt{\frac{4\pi}{k}}\delta^{3}$$

$$n_{1} = \sqrt{\frac{4\pi}{k}}\beta^{2}, \quad n_{2} = -\sqrt{\frac{4\pi}{k}}\beta^{1}, \quad n_{3} = \sqrt{\frac{4\pi}{k}}\bar{\gamma}_{4}, \quad n_{4} = -\sqrt{\frac{4\pi}{k}}\bar{\gamma}_{3}$$
 (B.12)

and using relations (B.8) between the two sets of conventions, we reproduce exactly the ABJ(M) WLs in (B.2), (B.3). Similarly, from the general construction in section 2 for BPS WLs $\tilde{W}_{\rm bos}$, $\tilde{W}_{\rm fer}$ and applying the same notational translation, we reproduce the ABJ(M) WLs with connections (B.6). This is a consistency check of our general construction.

B.2 1/2 BPS line WLs and their gravity duals

For special values of the parameters in (B.4) the number of supercharges preserved by W_{fer} , \tilde{W}_{fer} can enhance. For instance, it has been proved [13] that connections

$$L_{1/2}[\bar{\alpha}_{I}] = \begin{pmatrix} A_{0} + \frac{2\pi}{k} \left(\delta_{J}^{I} - \frac{2\alpha^{I}\bar{\alpha}_{J}}{|\alpha|^{2}}\right) \phi_{I}\bar{\phi}^{J} & \sqrt{\frac{4\pi}{k}}\bar{\alpha}_{I}\psi_{+}^{I} \\ \sqrt{\frac{4\pi}{k}}\bar{\psi}_{I-\frac{\alpha^{I}}{|\alpha|^{2}}} & B_{0} + \frac{2\pi}{k} \left(\delta_{J}^{I} - \frac{2\alpha^{I}\bar{\alpha}_{J}}{|\alpha|^{2}}\right)\bar{\phi}^{J}\phi_{I} \end{pmatrix}$$

$$\tilde{L}_{1/2}[\bar{\alpha}_{I}] = \begin{pmatrix} A_{0} - \frac{2\pi}{k} \left(\delta_{J}^{I} - \frac{2\alpha^{I}\bar{\alpha}_{J}}{|\alpha|^{2}}\right)\phi_{I}\bar{\phi}^{J} & \sqrt{\frac{4\pi}{k}}\bar{\alpha}_{I}\psi_{-}^{I} \\ -\sqrt{\frac{4\pi}{k}}\bar{\psi}_{I+\frac{\alpha^{I}}{|\alpha|^{2}}} & B_{0} - \frac{2\pi}{k} \left(\delta_{J}^{I} - \frac{2\alpha^{I}\bar{\alpha}_{J}}{|\alpha|^{2}}\right)\bar{\phi}^{J}\phi_{I} \end{pmatrix}$$
(B.13)

with constant parameters

$$\bar{\alpha}_I = (\bar{\alpha}_1, \bar{\alpha}_2, \bar{\alpha}_3, \bar{\alpha}_4), \quad \alpha^I \equiv (\bar{\alpha}_I)^*, \quad |\alpha|^2 \equiv \bar{\alpha}_I \alpha^I \neq 0$$
 (B.14)

give rise to 1/2 BPS fermionic operators $W_{1/2}[\bar{\alpha}_I]$, $\tilde{W}_{1/2}[\bar{\alpha}_I]$ that preserve complementary supercharges

$$\bar{\alpha}_{I}\theta_{+}^{IJ}, \quad \epsilon_{IJKL}\alpha^{J}\theta_{-}^{KL}, \quad \bar{\alpha}_{I}\vartheta_{+}^{IJ}, \quad \epsilon_{IJKL}\alpha^{J}\vartheta_{-}^{KL}$$

$$\bar{\alpha}_{I}\theta_{-}^{IJ}, \quad \epsilon_{IJKL}\alpha^{J}\theta_{+}^{KL}, \quad \bar{\alpha}_{I}\vartheta_{-}^{IJ}, \quad \epsilon_{IJKL}\alpha^{J}\vartheta_{+}^{KL}$$
(B.15)

These operators can be obtained from (B.3) and (B.6) with constant parameters (B.4) by setting $\beta^{1,2} = \frac{\alpha^{1,2}}{|\alpha|^2}$, $\bar{\gamma}_{3,4} = \delta^{3,4} = 0$ and performing a R-symmetry rotation $(\bar{\alpha}_1, \bar{\alpha}_2, 0, 0) \rightarrow (\bar{\alpha}_1, \bar{\alpha}_2, \bar{\alpha}_3, \bar{\alpha}_4)$.

For these 1/2 BPS operators gravity duals have been found [13, 33], which we now review.

The ABJM theory is dual to M-theory in $AdS_4 \times S^7/Z_k$ background

$$ds^{2} = R^{2} \left(\frac{1}{4} ds_{AdS_{4}}^{2} + ds_{S^{7}/Z_{k}}^{2} \right)$$
 (B.16)

where the metric of AdS_4 is

$$ds_{AdS_4}^2 = u^2(-dt^2 + dx_1^2 + dx_2^2) + \frac{du^2}{u^2}$$
(B.17)

and embedding S^7 in C^4 as [10]

$$z_{1} = \cos \frac{\beta}{2} \cos \frac{\theta_{1}}{2} e^{i\xi_{1}}, \quad \xi_{1} = -\frac{1}{4} (2\phi_{1} + \chi + \zeta)$$

$$z_{2} = \cos \frac{\beta}{2} \sin \frac{\theta_{1}}{2} e^{i\xi_{2}}, \quad \xi_{2} = -\frac{1}{4} (-2\phi_{1} + \chi + \zeta)$$

$$z_{3} = \sin \frac{\beta}{2} \cos \frac{\theta_{2}}{2} e^{i\xi_{3}}, \quad \xi_{3} = -\frac{1}{4} (2\phi_{2} - \chi + \zeta)$$

$$z_{4} = \sin \frac{\beta}{2} \sin \frac{\theta_{2}}{2} e^{i\xi_{4}}, \quad \xi_{4} = -\frac{1}{4} (-2\phi_{2} - \chi + \zeta)$$
(B.18)

we can write

$$ds_{S7}^{2} = \frac{1}{4} \left[d\beta^{2} + \cos^{2} \frac{\beta}{2} \left(d\theta_{1}^{2} + \sin^{2} \theta_{1} d\varphi_{1}^{2} \right) + \sin^{2} \frac{\beta}{2} \left(d\theta_{2}^{2} + \sin^{2} \theta_{2} d\varphi_{2}^{2} \right) \right.$$

$$\left. + \sin^{2} \frac{\beta}{2} \cos^{2} \frac{\beta}{2} \left(d\chi + \cos \theta_{1} d\varphi_{1} - \cos \theta_{2} d\varphi_{2} \right)^{2} \right.$$

$$\left. + \left(\frac{1}{2} d\zeta + \cos^{2} \frac{\beta}{2} \cos \theta_{1} d\varphi_{1} + \sin^{2} \frac{\beta}{2} \cos \theta_{2} d\varphi_{2} + \frac{1}{2} \cos \beta d\chi \right)^{2} \right]$$
(B.19)

Here $\beta, \theta_{1,2} \in [0, \pi]$, $\xi_{1,2,3,4} \in [0, 2\pi]$, so that $\phi_{1,2} \in [0, 2\pi]$, $\chi \in [0, 4\pi]$, $\zeta \in [0, 8\pi]$. The M-theory cycle is taken along the ζ direction.

The orbifold projection is realized by the \mathbf{Z}_k identification

$$(z_1, z_2, z_3, z_4) \sim e^{\frac{2\pi i}{k}}(z_1, z_2, z_3, z_4)$$
 (B.20)

or equivalently $\zeta \sim \zeta - \frac{8\pi}{k}$.

The general solution to the Killing spinor equations in M-theory on $AdS_4 \times S^7$ background reads [10, 33]

$$\epsilon = u^{\frac{1}{2}}h(\epsilon_1 + x^{\mu}\gamma_{\mu}\epsilon_2) - u^{-\frac{1}{2}}\gamma_3h\epsilon_2$$
 $\mu = 0, 1, 2$ (B.21)

where ϵ_1 , ϵ_2 are two constant Majorana spinors satisfying $\gamma^{012}\epsilon_i = \epsilon_i$, i = 1, 2, and

$$h = e^{\frac{\beta}{4}(\gamma_{34} - \gamma_{7\natural})} e^{\frac{\theta_1}{4}(\gamma_{35} - \gamma_{8\natural})} e^{\frac{\theta_2}{4}(\gamma_{46} + \gamma_{79})} e^{\frac{\xi_1}{2}\gamma_{3\natural}} e^{\frac{\xi_2}{2}\gamma_{58}} e^{\frac{\xi_3}{2}\gamma_{47}} e^{\frac{\xi_4}{2}\gamma_{69}}$$
(B.22)

Here γ_A , $A=0,\cdots,9, \natural$ are the eleven dimensional gamma matrices, and $\gamma_{0123456789 \natural}=1$. We decompose γ_A in terms of gamma matrices $\gamma_{a=0,1,2}$ in $\mathbf{R}^{1,2}$ and $\Gamma_{p=3,\cdots,9,\natural}$ in $\mathbf{C}^4\cong\mathbf{R}^8$ as

$$\gamma_a = -\gamma_a \otimes \Gamma, \quad a = 0, 1, 2$$

$$\gamma_p = \mathbf{1} \otimes \Gamma_p, \quad p = 3, \dots, 9, \natural \tag{B.23}$$

Correspondingly, ϵ_1, ϵ_2 get decomposed into direct products of Grassmann odd spinors θ , ϑ in $\mathbb{R}^{1,2}$ and Grassmann even spinors η in $\mathbb{C}^4 \cong \mathbb{R}^8$

$$\epsilon_1 \sim \theta \otimes \eta \qquad \epsilon_2 \sim \vartheta \otimes \eta$$
 (B.24)

The ϵ_1 decomposition is related to Poincaré supercharges θ and the ϵ_2 one is related to superconformal charges ϑ .

We write the η spinors in terms of gamma matrix eigenstates (a similar procedure applies also to ϑ)

$$\Gamma_{3b}\eta = it_1\eta, \quad \Gamma_{58}\eta = it_2\eta, \quad \Gamma_{47}\eta = it_3\eta, \quad \Gamma_{69}\eta = it_4\eta$$
 (B.25)

with $t_I = \pm$ for I = 1, 2, 3, 4. They satisfy $t_1 t_2 t_3 t_4 = 1$ as a consequence of the constraint $\gamma^{012} \epsilon_1 = \epsilon_1$. The ϵ_i Killing spinors can then be expressed as a linear combination of eight eigenstates

$$(t_1, t_2, t_3, t_4) = (+ + ++), (+ + --), (+ -+-), (+ --+), (- + +-), (- + +-), (- - ++), (- ---)$$
(B.26)

where each of them corresponds to one real degree of freedom. In the order of (B.26), we then write

$$\epsilon_1 = \sum_{i=1}^8 \theta^i \otimes \eta_i \quad , \quad \epsilon_2 = \sum_{i=1}^8 \vartheta^i \otimes \eta_i$$
(B.27)

For the Killing spinor (B.21), the quotient (B.20) leads to the constraint

$$\mathcal{L}_{\partial_{\mathcal{L}}} \epsilon = 0 \tag{B.28}$$

which gives

$$(\gamma_{3b} + \gamma_{58} + \gamma_{47} + \gamma_{69})\epsilon_i = 0, \quad i = 1, 2$$
 (B.29)

which on the decomposition (B.25) translates into

$$t_1 + t_2 + t_3 + t_4 = 0 (B.30)$$

The surviving states in (B.26) are then

$$(t_1, t_2, t_3, t_4) = (+ + --), (+ - +-), (+ - -+),$$

$$(- + +-), (- + -+), (- - ++)$$
(B.31)

Defining

$$\theta^{2} = \theta^{12} = -\theta^{21}, \quad \theta^{3} = \theta^{13} = -\theta^{31}, \quad \theta^{4} = \theta^{14} = -\theta^{41}$$

$$\theta^{5} = \theta^{23} = -\theta^{32}, \quad \theta^{6} = -\theta^{24} = \theta^{42}, \quad \theta^{7} = \theta^{34} = -\theta^{43}$$

$$\vartheta^{2} = \vartheta^{12} = -\vartheta^{21}, \quad \vartheta^{3} = \vartheta^{13} = -\vartheta^{31}, \quad \vartheta^{4} = \vartheta^{14} = -\vartheta^{41}$$

$$\vartheta^{5} = \vartheta^{23} = -\vartheta^{32}, \quad \vartheta^{6} = -\vartheta^{24} = \vartheta^{42}, \quad \vartheta^{7} = \vartheta^{34} = -\vartheta^{43}$$

$$\eta_{2} = \eta_{12} = -\eta_{21}, \quad \eta_{3} = \eta_{13} = -\eta_{31}, \quad \eta_{4} = \eta_{14} = -\eta_{41}$$

$$\eta_{5} = \eta_{23} = -\eta_{32}, \quad \eta_{6} = -\eta_{24} = \eta_{42}, \quad \eta_{7} = \eta_{34} = -\eta_{43}$$
(B.32)

we obtain

$$\epsilon_1 = \frac{1}{2}\theta^I \otimes \eta_I \quad , \quad \epsilon_2 = \frac{1}{2}\vartheta^I \otimes \eta_I$$
(B.33)

where I runs from 2 to 7.

The 1/2 BPS operator $W_{1/2}[\bar{\alpha}_I]$ along the line $x^{\mu}=(\tau,0,0)$ and corresponding to the superconnection $L_{1/2}[\bar{\alpha}_I]$ in (B.13) is dual to an M2-brane embedded as $t=\sigma^0, x_1=x_2=0, u=\sigma^1, \zeta=\sigma^2$ and localized at a point specified by the complex vector [33]

$$\frac{\alpha^{I}}{|\alpha|} = \left(\cos\frac{\beta}{2}\cos\frac{\theta_{1}}{2}e^{-\frac{i}{4}(2\phi_{1}+\chi+\zeta)}, \cos\frac{\beta}{2}\sin\frac{\theta_{1}}{2}e^{-\frac{i}{4}(-2\phi_{1}+\chi+\zeta)}, \sin\frac{\beta}{2}\cos\frac{\theta_{2}}{2}e^{-\frac{i}{4}(2\phi_{2}-\chi+\zeta)}, \sin\frac{\beta}{2}\sin\frac{\theta_{2}}{2}e^{-\frac{i}{4}(-2\phi_{2}-\chi+\zeta)}\right)$$
(B.34)

The 1/2 BPS WL $\tilde{W}_{1/2}[\bar{\alpha}_I]$ is dual to an anti–M2–brane at the same position that is specified by $\bar{\alpha}_I$.

The gravity duals of the 1/2 BPS WLs in ABJ(M) theory are helpful for identifying the gravity duals of some BPS WLs in $\mathcal{N}=4$ orbifold ABJM theory [18, 33], as we will review in the next appendix, and for identifying the gravity duals of some WLs in $\mathcal{N}=2$ orbifold ABJM theory as described in section 3.

B.3 Circle WLs in Euclidean space

In the Euclidean version of the ABJ(M) theory we can define BPS WLs along the circle

$$x^{\mu} = (\cos \tau, \sin \tau, 0) \tag{B.35}$$

In ABJ(M) notations, 1/6 BPS operators $W_{\rm bos},\,W_{\rm fer},\,\tilde{W}_{\rm bos},\,\tilde{W}_{\rm fer}$ correspond to superconnections

$$L_{\text{bos}} = \operatorname{diag}(A_{\mu}\dot{x}^{\mu} - \frac{2\pi i}{k}R^{I}{}_{J}\phi_{I}\bar{\phi}^{J}, B_{\mu}\dot{x}^{\mu} - \frac{2\pi i}{k}R^{I}{}_{J}\bar{\phi}^{J}\phi_{I})$$

$$L_{\text{fer}} = \begin{pmatrix} A_{\mu}\dot{x}^{\mu} - \frac{2\pi i}{k}U^{I}{}_{J}\phi_{I}\bar{\phi}^{J} & \sqrt{\frac{4\pi}{k}}(\bar{\alpha}_{I}\psi_{+}^{I} + \bar{\gamma}_{I}\psi_{-}^{I}) \\ \sqrt{\frac{4\pi}{k}}(-\bar{\psi}_{I-}\beta^{I} + \bar{\psi}_{I+}\delta^{I}) & B_{\mu}\dot{x}^{\mu} - \frac{2\pi i}{k}U^{I}{}_{J}\bar{\phi}^{J}\phi_{I} \end{pmatrix}$$

$$\tilde{L}_{\text{bos}} = \operatorname{diag}(A_{\mu}\dot{x}^{\mu} + \frac{2\pi i}{k}R^{I}{}_{J}\phi_{I}\bar{\phi}^{J}, B_{\mu}\dot{x}^{\mu} + \frac{2\pi i}{k}R^{I}{}_{J}\bar{\phi}^{J}\phi_{I})$$

$$\tilde{L}_{\text{fer}} = \begin{pmatrix} A_{\mu}\dot{x}^{\mu} + \frac{2\pi i}{k}U^{I}{}_{J}\phi_{I}\bar{\phi}^{J} & \sqrt{\frac{4\pi}{k}}(\bar{\alpha}_{I}\psi_{-}^{I} + \bar{\gamma}_{I}\psi_{+}^{I}) \\ \sqrt{\frac{4\pi}{k}}(\bar{\psi}_{I+}\beta^{I} - \bar{\psi}_{I-}\delta^{I}) & B_{\mu}\dot{x}^{\mu} + \frac{2\pi i}{k}U^{I}{}_{J}\bar{\phi}^{J}\phi_{I} \end{pmatrix}$$
(B.36)

with the same constant parameters $R^I{}_J, U^I{}_J, \bar{\alpha}_I, \beta^I, \bar{\gamma}_I, \delta^I$ in (B.2), (B.4). $W_{\rm bos}, W_{\rm fer}$ operators preserve supercharges

$$\vartheta^{12} = -i\gamma_3 \theta^{12}, \quad \vartheta^{34} = i\gamma_3 \theta^{34}$$
 (B.37)

whereas \tilde{W}_{bos} , \tilde{W}_{fer} preserve the complementary set

$$\vartheta^{12} = i\gamma_3\theta^{12}, \quad \vartheta^{34} = -i\gamma_3\theta^{34} \tag{B.38}$$

 $W_{\rm bos}$, $\tilde{W}_{\rm bos}$ are related by a R-symmetry rotation $I=1,2\leftrightarrow I=3,4$, whereas the $W_{\rm fer}$, $\tilde{W}_{\rm fer}$ operators are related by a R-symmetry rotation $I=1,2\leftrightarrow I=3,4$ plus a parameter redefinition $\bar{\alpha}_I\leftrightarrow\bar{\gamma}_I$, $\beta^I\leftrightarrow\delta^I$.

1/2 BPS operators $W_{1/2}[\bar{\alpha}_I]$, $\tilde{W}_{1/2}[\bar{\alpha}_I]$ can also be defined in Euclidean signature. They correspond to connections

$$L_{1/2}[\bar{\alpha}_{I}] = \begin{pmatrix} A_{\mu}\dot{x}^{\mu} - \frac{2\pi i}{k} \left(\delta_{J}^{I} - \frac{2\alpha^{I}\bar{\alpha}_{J}}{|\alpha|^{2}}\right)\phi_{I}\bar{\phi}^{J} & \sqrt{\frac{4\pi}{k}}\bar{\alpha}_{I}\psi_{+}^{I} \\ -\sqrt{\frac{4\pi}{k}}\bar{\psi}_{I-\frac{\alpha^{I}}{|\alpha|^{2}}} & B_{\mu}\dot{x}^{\mu} - \frac{2\pi i}{k} \left(\delta_{J}^{I} - \frac{2\alpha^{I}\bar{\alpha}_{J}}{|\alpha|^{2}}\right)\bar{\phi}^{J}\phi_{I} \end{pmatrix}$$

$$\tilde{L}_{1/2}[\bar{\alpha}_{I}] = \begin{pmatrix} A_{\mu}\dot{x}^{\mu} + \frac{2\pi i}{k} \left(\delta_{J}^{I} - \frac{2\alpha^{I}\bar{\alpha}_{J}}{|\alpha|^{2}}\right)\phi_{I}\bar{\phi}^{J} & \sqrt{\frac{4\pi}{k}}\bar{\alpha}_{I}\psi_{-}^{I} \\ \sqrt{\frac{4\pi}{k}}\bar{\psi}_{I+\frac{\alpha^{I}}{|\alpha|^{2}}} & B_{\mu}\dot{x}^{\mu} + \frac{2\pi i}{k} \left(\delta_{J}^{I} - \frac{2\alpha^{I}\bar{\alpha}_{J}}{|\alpha|^{2}}\right)\bar{\phi}^{J}\phi_{I} \end{pmatrix}$$
(B.39)

The $W_{1/2}[\bar{\alpha}_I]$ operator preserves supercharges

$$\bar{\alpha}_I \vartheta^{IJ} = -i\bar{\alpha}_I \gamma_3 \theta^{IJ}, \quad \epsilon_{IJKL} \alpha^J \vartheta^{KL} = i\gamma_3 \epsilon_{IJKL} \alpha^J \theta^{KL}$$
 (B.40)

while $\tilde{W}_{1/2}[\bar{\alpha}_I]$ preserves complementary supercharges

$$\bar{\alpha}_I \vartheta^{IJ} = i\bar{\alpha}_I \gamma_3 \theta^{IJ}, \quad \epsilon_{IJKL} \alpha^J \vartheta^{KL} = -i\gamma_3 \epsilon_{IJKL} \alpha^J \theta^{KL}$$
 (B.41)

C BPS WLs in $\mathcal{N} = 4$ orbifold ABJM theory

As a further check of our general construction we now apply it to the case of $\mathcal{N}=4$ orbifold ABJM theory and show that, under a suitable change of notations, it reproduces the known 1/4 BPS WLs found in [18].

General circular quiver $\mathcal{N}=4$ SCSM theories were constructed in [50, 51], while the special case of $\mathcal{N}=4$ orbifold ABJM theory was introduced in [22]. For gauge group $[U(N)_k \times U(N)_{-k}]^r$ it can be obtained by applying a Z_r quotient to the $U(rN)_k \times U(rN)_{-k}$ ABJM theory. The SU(4) R-symmetry is broken to $SU(2) \times SU(2)$, and we decompose the R-symmetry index as

$$I = 1, 2, 4, 3 \rightarrow i = 1, 2, \ \hat{\imath} = \hat{1}, \hat{2}$$
 (C.1)

We can write the theory in ABJM notations as in figure 10(a), or in $\mathcal{N}=2$ notations, as in figure 10(b), under the supercharge identifications $\theta^{1\hat{1}}=\bar{\theta}_{2\hat{2}}=\theta$, $\theta^{2\hat{2}}=\bar{\theta}_{1\hat{1}}=\bar{\theta}$, $\vartheta^{1\hat{1}}=\bar{\vartheta}_{2\hat{2}}=\vartheta$. The two notations are related by

$$A_{\mu}^{(2\ell-1)} = A_{\mu}^{(2\ell-1)}, \quad B_{\mu}^{(2\ell)} = A_{\mu}^{(2\ell)}$$

$$\phi_{i}^{(2\ell)} = (Z_{1}^{(2\ell)}, \bar{Z}_{(2\ell)}^{2}), \quad \phi_{\hat{i}}^{(2\ell-1)} = (Z_{4}^{(2\ell-1)}, \bar{Z}_{(2\ell-1)}^{3})$$

$$\psi_{(2\ell)}^{\hat{i}} = (\zeta_{1}^{(2\ell)}, \bar{\zeta}_{(2\ell)}^{2}), \quad \psi_{(2\ell-1)}^{i} = (-\zeta_{4}^{(2\ell-1)}, -\bar{\zeta}_{(2\ell-1)}^{3})$$
(C.2)

From the results in section 2 we obtain the 1/4 BPS WL W_{bos} , W_{fer} in $\mathcal{N}=2$ notations with connections

$$L_{\text{bos}} = \text{diag}(A_{\mu}^{(1)} - \sigma^{(1)}, A^{(2)} - \sigma^{(2)}, \cdots, A^{(2r)} - \sigma^{(2r)})$$

$$L_{\text{fer}} = L_{\text{bos}} + B + F, \quad B = \bar{M}_Z N_{\bar{Z}} + N_{\bar{Z}} \bar{M}_Z, \quad F = \bar{M}_{\zeta} + N_{\bar{\zeta}}$$
(C.3)




Figure 10. The quiver diagram of $\mathcal{N}=4$ orbifold ABJM theory in (a) ABJM notations and (b) $\mathcal{N}=2$ notations.

with

$$\sigma^{(2\ell-1)} = \frac{2\pi}{k} \left(Z_1^{(2\ell-2)} \bar{Z}_{(2\ell-2)}^1 + Z_4^{(2\ell-1)} \bar{Z}_{(2\ell-1)}^4 - \bar{Z}_{(2\ell-2)}^2 Z_2^{(2\ell-2)} - \bar{Z}_{(2\ell-1)}^3 Z_3^{(2\ell-1)} \right)$$

$$\sigma^{(2\ell)} = \frac{2\pi}{k} \left(\bar{Z}_{(2\ell)}^1 Z_1^{(2\ell)} + \bar{Z}_{(2\ell-1)}^4 Z_4^{(2\ell-1)} - Z_2^{(2\ell)} \bar{Z}_{(2\ell)}^2 - Z_3^{(2\ell-1)} \bar{Z}_{(2\ell-1)}^3 \right)$$
(C.4)

The nonvanishing blocks of the matrices $\bar{M}_Z, N_{\bar{Z}}, \bar{M}_{\zeta}, N_{\bar{\zeta}}$ are

$$\begin{split} [\bar{M}_Z]_{(2\ell-1,2\ell)} &= \bar{m}_{(2\ell-1)}^4 Z_4^{(2\ell-1)}, \quad [\bar{M}_Z]_{(2\ell,2\ell+1)} = \bar{m}_{(2\ell)}^2 Z_2^{(2\ell)} \\ [\bar{M}_Z]_{(2\ell,2\ell-1)} &= \bar{m}_{(2\ell-1)}^3 Z_3^{(2\ell-1)}, \quad [\bar{M}_Z]_{(2\ell+1,2\ell)} = \bar{m}_{(2\ell)}^1 Z_1^{(2\ell)} \\ [N_{\bar{Z}}]_{(2\ell-1,2\ell)} &= n_3^{(2\ell-1)} \bar{Z}_{(2\ell-1)}^3, \quad [N_{\bar{Z}}]_{(2\ell,2\ell+1)} = n_1^{(2\ell)} \bar{Z}_{(2\ell)}^1 \\ [N_{\bar{Z}}]_{(2\ell,2\ell-1)} &= n_4^{(2\ell-1)} \bar{Z}_{(2\ell-1)}^4, \quad [N_{\bar{Z}}]_{(2\ell+1,2\ell)} = n_2^{(2\ell)} \bar{Z}_{(2\ell)}^2 \\ [\bar{M}_\zeta]_{(2\ell-1,2\ell)} &= \bar{m}_{(2\ell-1)}^4 \zeta_{4+}^{(2\ell-1)}, \quad [\bar{M}_\zeta]_{(2\ell,2\ell+1)} = \bar{m}_{(2\ell)}^2 \zeta_{2+}^{(2\ell)} \\ [\bar{M}_\zeta]_{(2\ell,2\ell-1)} &= \bar{m}_{(2\ell-1)}^3 \zeta_{3+}^{(2\ell-1)}, \quad [\bar{M}_\zeta]_{(2\ell+1,2\ell)} = \bar{m}_{(2\ell)}^1 \zeta_{1+}^{(2\ell)} \\ [N_{\bar{\zeta}}]_{(2\ell-1,2\ell)} &= n_3^{(2\ell-1)} \bar{\zeta}_{3+}^3, \quad [N_{\bar{\zeta}}]_{(2\ell,2\ell+1)} = n_1^{(2\ell)} \bar{\zeta}_{(2\ell)-}^1 \\ [N_{\bar{\zeta}}]_{(2\ell,2\ell-1)} &= n_4^{(2\ell-1)} \bar{\zeta}_{(2\ell-1)-}^4, \quad [N_{\bar{\zeta}}]_{(2\ell+1,2\ell)} = n_2^{(2\ell)} \bar{\zeta}_{(2\ell)-}^2 \end{split} \tag{C.5}$$

with constraints on the parameters

$$\begin{split} \bar{m}_{(2\ell-1)}^4 \bar{m}_{(2\ell-1)}^3 &= \bar{m}_{(2\ell-1)}^4 \bar{m}_{(2\ell-2)}^2 = \bar{m}_{(2\ell-1)}^4 \bar{m}_{(2\ell)}^2 = 0 \\ \bar{m}_{(2\ell)}^1 \bar{m}_{(2\ell)}^2 &= \bar{m}_{(2\ell)}^1 \bar{m}_{(2\ell-1)}^3 = \bar{m}_{(2\ell)}^1 \bar{m}_{(2\ell+1)}^3 = 0 \\ n_3^{(2\ell-1)} n_4^{(2\ell-1)} &= n_3^{(2\ell-1)} n_1^{(2\ell-2)} = n_3^{(2\ell-1)} n_1^{(2\ell)} = 0 \\ n_2^{(2\ell)} n_1^{(2\ell)} &= n_2^{(2\ell)} n_4^{(2\ell-1)} = n_2^{(2\ell)} n_4^{(2\ell+1)} = 0 \end{split} \tag{C.6}$$

By redefining the parameters as

$$\begin{split} \bar{m}_{(2\ell)}^1 &= \sqrt{\frac{4\pi}{k}} \bar{\alpha}_{\hat{1}}^{(2\ell)}, \quad \bar{m}_{(2\ell)}^2 = -\sqrt{\frac{4\pi}{k}} \delta_{(2\ell)}^{\hat{2}} \\ \bar{m}_{(2\ell-1)}^3 &= \sqrt{\frac{4\pi}{k}} \delta_{(2\ell-1)}^2, \quad \bar{m}_{(2\ell-1)}^4 = -\sqrt{\frac{4\pi}{k}} \bar{\alpha}_{\hat{1}}^{(2\ell-1)} \\ n_1^{(2\ell)} &= \sqrt{\frac{4\pi}{k}} \beta_{(2\ell)}^{\hat{1}}, \quad n_2^{(2\ell)} = \sqrt{\frac{4\pi}{k}} \bar{\gamma}_{\hat{2}}^{(2\ell)} \\ n_3^{(2\ell-1)} &= -\sqrt{\frac{4\pi}{k}} \bar{\gamma}_2^{(2\ell-1)}, \quad n_4^{(2\ell-1)} = -\sqrt{\frac{4\pi}{k}} \beta_{(2\ell-1)}^1 \end{split} \tag{C.7}$$

and taking into account relations (C.2), we can write the 1/4 BPS WL along the line $x^{\mu} = (\tau, 0, 0)$ in ABJM notations.

The connection of the bosonic 1/4 BPS WL $W_{\rm bos}$ reads

$$L_{\text{bos}} = \text{diag}(\mathcal{A}^{(1)}, \mathcal{B}^{(2)}, \cdots, \mathcal{A}^{(2r-1)}, \mathcal{B}^{(2r)})$$
(C.8)

whereas, for $r \geq 3$ the connection of the fermionic 1/4 BPS WL W_{fer} is

$$L_{\text{fer}} = \begin{pmatrix} \mathcal{A}^{(1)} & f_1^{(1)} & h_1^{(1)} & h_2^{(2r-1)} & f_2^{(2r)} \\ f_2^{(1)} & \mathcal{B}^{(2)} & f_1^{(2)} & h_1^{(2)} & h_2^{(2r)} \\ h_2^{(1)} & f_2^{(2)} & \mathcal{A}^{(3)} & f_1^{(3)} & \ddots \\ & & & & & & & \\ & & h_2^{(2)} & f_2^{(3)} & \mathcal{B}^{(4)} & \ddots & \ddots \\ & & & \ddots & \ddots & \ddots & h_1^{(2r-3)} \\ & & & \ddots & \ddots & \ddots & h_1^{(2r-3)} \\ & & & & \ddots & \ddots & \ddots & f_1^{(2r-2)} & h_1^{(2r-2)} \\ h_1^{(2r-1)} & & & & h_2^{(2r-3)} & f_2^{(2r-2)} & \mathcal{A}^{(2r-1)} & f_1^{(2r-1)} \\ f_1^{(2r)} & h_1^{(2r)} & & & h_2^{(2r-2)} & f_2^{(2r-1)} & \mathcal{B}^{(2r)} \end{pmatrix}$$

$$(C.9)$$

and for r=2 it is

$$L_{\text{fer}} = \begin{pmatrix} \mathcal{A}^{(1)} & f_1^{(1)} & h_1^{(1)} + h_2^{(3)} & f_2^{(4)} \\ f_2^{(1)} & \mathcal{B}^{(2)} & f_1^{(2)} & h_1^{(2)} + h_2^{(4)} \\ h_1^{(3)} + h_2^{(1)} & f_2^{(2)} & \mathcal{A}^{(3)} & f_1^{(3)} \\ f_1^{(4)} & h_1^{(4)} + h_2^{(2)} & f_2^{(3)} & \mathcal{B}^{(4)} \end{pmatrix}$$
(C.10)

Here we have defined

$$\begin{split} \mathcal{A}^{(2\ell-1)} &= A_0^{(2\ell-1)} - \frac{2\pi}{k} \Big(U_{(2\ell-1)}{}^i{}_j \phi_i^{(2\ell-2)} \bar{\phi}_{(2\ell-2)}^j + U_{(2\ell-1)}{}^i{}_j \phi_i^{(2\ell-1)} \bar{\phi}_{(2\ell-1)}^j \Big) \\ \mathcal{B}^{(2\ell)} &= B_0^{(2\ell)} - \frac{2\pi}{k} \Big(U_{(2\ell)}{}^i{}_j \bar{\phi}_{(2\ell)}^j \phi_i^{(2\ell)} + U_{(2\ell)}{}^i{}_j \bar{\phi}_{(2\ell-1)}^j \phi_i^{(2\ell-1)} \Big) \\ f_1^{(2\ell-1)} &= \sqrt{\frac{4\pi}{k}} \Big(\bar{\alpha}_1^{(2\ell-1)} \psi_{(2\ell-1)+}^1 + \bar{\gamma}_2^{(2\ell-1)} \psi_{(2\ell-1)-}^2 \Big) \\ f_1^{(2\ell)} &= \sqrt{\frac{4\pi}{k}} \Big(\bar{\psi}_{1-}^{(2\ell)} \beta_{(2\ell)}^i - \bar{\psi}_{2+}^{(2\ell)} \delta_{(2\ell)}^2 \Big) \\ f_2^{(2\ell-1)} &= \sqrt{\frac{4\pi}{k}} \Big(\bar{\psi}_{1-}^{(2\ell-1)} \beta_{(2\ell-1)}^1 - \bar{\psi}_{2+}^{(2\ell-1)} \delta_{(2\ell-1)}^2 \Big) \\ f_2^{(2\ell)} &= \sqrt{\frac{4\pi}{k}} \Big(\bar{\alpha}_{\hat{1}}^{(2\ell)} \psi_{(2\ell)+}^{\hat{1}} + \bar{\gamma}_{\hat{2}}^{(2\ell)} \psi_{(2\ell)-}^{\hat{2}} \Big) \\ h_1^{(2\ell-1)} &= -\frac{2\pi}{k} U_{(2\ell-1)}{}^i{}_j \phi_i^{(2\ell-1)} \bar{\phi}_{(2\ell)}^j, \quad h_1^{(2\ell)} &= -\frac{2\pi}{k} U_{(2\ell)}{}^i{}_j \bar{\phi}_{(2\ell)}^j \phi_i^{(2\ell+1)} \\ h_2^{(2\ell-1)} &= -\frac{2\pi}{k} U_{(2\ell-1)}{}^i{}_j \phi_i^{(2\ell)} \bar{\phi}_{(2\ell-1)}^j, \quad h_2^{(2\ell)} &= -\frac{2\pi}{k} U_{(2\ell)}{}^i{}_j \bar{\phi}_{(2\ell+1)}^j \phi_i^{(2\ell)} \end{aligned} \tag{C.11}$$

with constant parameters

$$\begin{split} &U_{(2\ell-1)}{}^{i}{}_{j} = \mathrm{diag} \big(1 - 2\bar{\alpha}_{\hat{1}}^{(2\ell-2)} \beta_{(2\ell-2)}^{\hat{1}}, -1 + 2\bar{\gamma}_{\hat{2}}^{(2\ell-2)} \delta_{(2\ell-2)}^{\hat{2}} \big) \\ &U_{(2\ell-1)}{}^{i}{}_{\hat{j}} = \mathrm{diag} \big(1 - 2\bar{\alpha}_{1}^{(2\ell-1)} \beta_{(2\ell-1)}^{1}, -1 + 2\bar{\gamma}_{2}^{(2\ell-1)} \delta_{(2\ell-1)}^{2} \big) \\ &U_{(2\ell)}{}^{i}{}_{\hat{j}} = \mathrm{diag} \big(1 - 2\bar{\alpha}_{\hat{1}}^{(2\ell)} \beta_{(2\ell)}^{\hat{1}}, -1 + 2\bar{\gamma}_{\hat{2}}^{(2\ell)} \delta_{(2\ell)}^{\hat{2}} \big) \\ &U_{(2\ell)}{}^{i}{}_{\hat{j}} = \mathrm{diag} \big(1 - 2\bar{\alpha}_{1}^{(2\ell-1)} \beta_{(2\ell-1)}^{1}, -1 + 2\bar{\gamma}_{2}^{(2\ell-1)} \delta_{(2\ell-1)}^{2} \big) \\ &U_{(2\ell-1)}{}^{i}{}_{\hat{j}} = \mathrm{diag} \big(2\bar{\alpha}_{1}^{(2\ell-1)} \beta_{(2\ell)}^{\hat{1}}, -2\bar{\gamma}_{2}^{(2\ell-1)} \delta_{(2\ell)}^{\hat{2}} \big) \\ &U_{(2\ell-1)}{}^{i}{}_{\hat{j}} = \mathrm{diag} \big(2\bar{\alpha}_{\hat{1}}^{(2\ell)} \beta_{(2\ell-1)}^{1}, -2\bar{\gamma}_{\hat{2}}^{(2\ell)} \delta_{(2\ell-1)}^{2} \big) \\ &U_{(2\ell)}{}^{i}{}_{\hat{j}} = \mathrm{diag} \big(2\bar{\alpha}_{1}^{(2\ell+1)} \beta_{(2\ell)}^{\hat{1}}, -2\bar{\gamma}_{2}^{(2\ell+1)} \delta_{(2\ell)}^{\hat{2}} \big) \\ &U_{(2\ell)}{}^{i}{}_{\hat{j}} = \mathrm{diag} \big(2\bar{\alpha}_{\hat{1}}^{(2\ell)} \beta_{(2\ell+1)}^{1}, -2\bar{\gamma}_{\hat{2}}^{(2\ell)} \delta_{(2\ell+1)}^{2} \big) \end{split} \tag{C.12}$$

The parameters are subject to the constraints

$$\begin{split} \bar{\alpha}_{1}^{(2\ell-1)} \delta_{(2\ell-1)}^{2} &= \bar{\alpha}_{1}^{(2\ell-1)} \delta_{(2\ell-2)}^{\hat{2}} = \bar{\alpha}_{1}^{(2\ell-1)} \delta_{(2\ell)}^{\hat{2}} = 0 \\ \bar{\alpha}_{\hat{1}}^{(2\ell)} \delta_{(2\ell)}^{2} &= \bar{\alpha}_{\hat{1}}^{(2\ell)} \delta_{(2\ell-1)}^{\hat{2}} = \bar{\alpha}_{\hat{1}}^{(2\ell)} \delta_{(2\ell+1)}^{\hat{2}} = 0 \\ \bar{\gamma}_{2}^{(2\ell-1)} \beta_{(2\ell-1)}^{1} &= \bar{\gamma}_{2}^{(2\ell-1)} \beta_{(2\ell-2)}^{\hat{1}} = \bar{\gamma}_{2}^{(2\ell-1)} \beta_{(2\ell)}^{\hat{1}} = 0 \\ \bar{\gamma}_{\hat{2}}^{(2\ell)} \beta_{(2\ell)}^{1} &= \bar{\gamma}_{\hat{2}}^{(2\ell)} \beta_{(2\ell-1)}^{\hat{1}} = \bar{\gamma}_{\hat{2}}^{(2\ell)} \beta_{(2\ell+1)}^{\hat{1}} = 0 \end{split} \tag{C.13}$$

We have exactly reproduced the fermionic 1/4 BPS WL in [18] preserving supercharges

$$\theta_{+}^{1\hat{1}}, \quad \theta_{-}^{2\hat{2}}, \quad \vartheta_{+}^{1\hat{1}}, \quad \vartheta_{-}^{2\hat{2}}$$
 (C.14)

Similarly, from the 1/2 BPS WL \tilde{W}_{fer} in section 2, we can construct a 1/4 BPS WL in $\mathcal{N}=4$ orbifold ABJM theory that preserves supercharges $\theta_{-}^{1\hat{1}}$, $\theta_{+}^{2\hat{2}}$, $\vartheta_{-}^{1\hat{1}}$, $\vartheta_{+}^{2\hat{2}}$.

In general, we do not know how to construct the gravity duals of BPS WLs in $\mathcal{N}=4$ orbifold ABJM, directly. However, for WLs that can be obtained by an orbifold quotient of the 1/2 BPS operators of the ABJM theory, we can exploit their known gravity duals [18, 33] and obtain the corresponding ones in $\mathcal{N}=4$ orbifold ABJM theory by taking their orbifold quotient.

The $\mathcal{N}=4$ orbifold ABJM theory is dual to M-theory in $AdS_4 \times S^7/(Z_{rk} \times Z_r)$ space-time [22, 52, 53]

$$ds^{2} = R^{2} \left(\frac{1}{4} ds_{AdS_{4}}^{2} + ds_{S^{7}/(Z_{rk} \times Z_{r})}^{2} \right)$$
 (C.15)

where the metric of AdS₄ is given in (B.17), the metric of S⁷ in (B.19), and the $Z_{rk} \times Z_r$ quotient is generated by

$$(z_1, z_2, z_3, z_4) \sim e^{\frac{2\pi i}{rk}}(z_1, z_2, z_3, z_4), \quad (z_1, z_2, z_3, z_4) \sim (e^{\frac{2\pi i}{r}}z_1, e^{\frac{2\pi i}{r}}z_2, z_3, z_4)$$
 (C.16)

which is equivalent to

$$\zeta \sim \zeta - \frac{8\pi}{rk}, \quad \chi \sim \chi - \frac{4\pi}{r}, \quad \zeta \sim \zeta - \frac{4\pi}{r}$$
 (C.17)

Performing the quotient of the 1/2 BPS operators $W_{1/2}[\bar{\alpha}_I]$ and $\tilde{W}_{1/2}[\bar{\alpha}_I]$ in ABJM theory corresponding to connections (B.13), we obtain 1/2 or 1/4 BPS WL in $\mathcal{N}=4$ ABJM theory, depending on the value of $\bar{\alpha}_I$. The operator coming from $W_{1/2}[\bar{\alpha}_I]$ is dual to an M2-brane that wraps a cycle in the internal space specified by $\bar{\alpha}_I$, whereas the operator from $\tilde{W}_{1/2}[\bar{\alpha}_I]$ is dual to an anti-M2-brane that wraps the same interval cycle.

D Connections for 1/2 BPS WL in $\mathcal{N} = 2$ orbifold ABJM

In this appendix we collect the connections for 1/2 BPS WL in $\mathcal{N}=2$ orbifold ABJM when r=4,3,2. With definitions (3.10), we have for r=4

$$\mathcal{A} = \begin{pmatrix}
\mathcal{A}^{(1)} & 0 & h_1^{(1)} + h_3^{(3)} & 0 \\
0 & \mathcal{A}^{(2)} & 0 & h_1^{(2)} + h_3^{(4)} \\
h_1^{(3)} + h_3^{(1)} & 0 & \mathcal{A}^{(3)} & 0 \\
0 & h_1^{(4)} + h_3^{(2)} & 0 & \mathcal{A}^{(4)}
\end{pmatrix}$$

$$\mathcal{B} = \begin{pmatrix}
\mathcal{B}^{(1)} & 0 & h_4^{(1)} + h_2^{(3)} & 0 \\
0 & \mathcal{B}^{(2)} & 0 & h_4^{(2)} + h_2^{(4)} \\
h_4^{(3)} + h_2^{(1)} & 0 & \mathcal{B}^{(3)} & 0 \\
0 & h_4^{(4)} + h_2^{(2)} & 0 & \mathcal{B}^{(4)}
\end{pmatrix}$$

$$f_1 = \begin{pmatrix}
f_1^{(1)} f_3^{(1)} & 0 & f_5^{(4)} \\
f_5^{(1)} f_1^{(2)} f_3^{(2)} & 0 \\
0 & f_5^{(2)} f_1^{(3)} f_3^{(3)} \\
f_3^{(4)} & 0 & f_5^{(3)} f_1^{(4)}
\end{pmatrix}, \quad f_2 = \begin{pmatrix}
f_2^{(1)} f_6^{(1)} & 0 & f_4^{(4)} \\
f_4^{(1)} f_2^{(2)} f_6^{(2)} & 0 \\
0 & f_4^{(2)} f_2^{(3)} f_6^{(3)} \\
f_6^{(4)} & 0 & f_4^{(4)} f_2^{(4)}
\end{pmatrix}$$
(D.1)

for r = 3

$$\mathcal{A} = \begin{pmatrix}
\mathcal{A}^{(1)} & h_3^{(2)} & h_1^{(1)} \\
h_1^{(2)} & \mathcal{A}^{(2)} & h_3^{(3)} \\
h_3^{(1)} & h_1^{(3)} & \mathcal{A}^{(3)}
\end{pmatrix}, \quad \mathcal{B} = \begin{pmatrix}
\mathcal{B}^{(1)} & h_2^{(2)} & h_4^{(1)} \\
h_2^{(2)} & \mathcal{B}^{(2)} & h_2^{(3)} \\
h_2^{(1)} & h_4^{(3)} & \mathcal{B}^{(3)}
\end{pmatrix}$$

$$f_1 = \begin{pmatrix}
f_1^{(1)} & f_3^{(1)} & f_5^{(3)} \\
f_5^{(1)} & f_1^{(2)} & f_3^{(2)} \\
f_5^{(2)} & f_1^{(2)} & f_3^{(2)}
\end{pmatrix}, \quad f_2 = \begin{pmatrix}
f_2^{(1)} & f_6^{(1)} & f_4^{(3)} \\
f_4^{(1)} & f_2^{(2)} & f_6^{(2)} \\
f_4^{(3)} & f_2^{(2)} & f_2^{(2)} \\
f_6^{(3)} & f_4^{(2)} & f_2^{(2)}
\end{pmatrix} \tag{D.2}$$

and for r=2

$$\mathcal{A} = \begin{pmatrix} \mathcal{A}^{(1)} + h_1^{(1)} + h_3^{(1)} & 0 \\ 0 & \mathcal{A}^{(2)} + h_1^{(2)} + h_3^{(2)} \end{pmatrix}
\mathcal{B} = \begin{pmatrix} \mathcal{B}^{(1)} + h_2^{(1)} + h_4^{(1)} & 0 \\ 0 & \mathcal{B}^{(2)} + h_2^{(2)} + h_4^{(2)} \end{pmatrix}
f_1 = \begin{pmatrix} f_1^{(1)} & f_3^{(1)} + f_5^{(2)} \\ f_3^{(2)} + f_5^{(1)} & f_1^{(2)} \end{pmatrix}
f_2 = \begin{pmatrix} f_2^{(1)} & f_6^{(1)} + f_4^{(2)} \\ f_6^{(2)} + f_4^{(1)} & f_2^{(2)} \end{pmatrix} \tag{D.3}$$

E Lagrangian and Feynman rules

In Minkowski spacetime, from the superspace lagrangian (2.4) we obtain the relevant terms in components

$$\mathcal{L}_{CS} = \sum_{a} \frac{k_{a}}{4\pi} \varepsilon^{\mu\nu\rho} \text{Tr} \left(A_{\mu}^{(a)} \partial_{\nu} A_{\rho}^{(a)} + \frac{2i}{3} A_{\mu}^{(a)} A_{\nu}^{(a)} A_{\rho}^{(a)} \right)$$

$$\mathcal{L}_{k} = \sum_{ab} \text{Tr} \left(-D_{\mu} \bar{Z}_{i}^{(ba)} D^{\mu} Z_{(ab)}^{i} + i \bar{\zeta}_{i}^{(ba)} \gamma^{\mu} D_{\mu} \zeta_{(ab)}^{i} \right)$$
(E.1)

By standard Wick rotation, the lagrangian in Euclidean space is given by

$$\mathcal{L}_{CS} = -\sum_{a} \frac{ik_{a}}{4\pi} \varepsilon^{\mu\nu\rho} \text{Tr} \left(A_{\mu}^{(a)} \partial_{\nu} A_{\rho}^{(a)} + \frac{2i}{3} A_{\mu}^{(a)} A_{\nu}^{(a)} A_{\rho}^{(a)} \right)$$

$$\mathcal{L}_{k} = \sum_{a,b} \text{Tr} \left(D_{\mu} \bar{Z}_{i}^{(ba)} D^{\mu} Z_{(ab)}^{i} - i \bar{\zeta}_{i}^{(ba)} \gamma^{\mu} D_{\mu} \zeta_{(ab)}^{i} \right)$$
(E.2)

Here the definitions of covariant derivatives are

$$D_{\mu}Z_{(ab)}^{i} = \partial_{\mu}Z_{(ab)}^{i} + iA_{\mu}^{(a)}Z_{(ab)}^{i} - iZ_{(ab)}^{i}A_{\mu}^{(b)}, \tag{E.3}$$

$$D_{\mu}\zeta_{(ab)}^{i} = \partial_{\mu}\zeta_{(ab)}^{i} + iA_{\mu}^{(a)}\zeta_{(ab)}^{i} - i\zeta_{(ab)}^{i}A_{\mu}^{(b)}$$
(E.4)

We work in Landau gauge for vector fields. Tree and one-loop propagators are drawn in figure 11. In dimensional regularization, $d = 3 - 2\epsilon$, their explicit expressions at tree level are

$$\langle A_{\mu}^{(a)}{}_{p}{}^{q}(x) A_{\nu}^{(b)}{}_{r}{}^{s}(y) \rangle^{(0)} = \delta^{ab} \delta_{p}^{s} \delta_{r}^{q} \frac{\mathrm{i}}{k_{a}} \frac{\Gamma(\frac{3}{2} - \epsilon)}{\pi^{\frac{1}{2} - \epsilon}} \frac{\varepsilon_{\mu\nu\rho}(x - y)^{\rho}}{|x - y|^{3 - 2\epsilon}}$$

$$\langle Z_{(ab)p}^{i}{}_{q}(x) \bar{Z}_{j}^{(cd)}{}_{r}{}^{s}(y) \rangle^{(0)} = \delta_{a}^{d} \delta_{b}^{c} \delta_{j}^{i} \delta_{p}^{s} \delta_{r}^{q} \frac{\Gamma(\frac{1}{2} - \epsilon)}{4\pi^{\frac{3}{2} - \epsilon}} \frac{1}{|x - y|^{1 - 2\epsilon}}$$

$$\langle \zeta_{(ab)p}^{i}{}_{\alpha}(x) \bar{\zeta}_{j}^{(cd)}{}_{r}{}^{s\beta}(y) \rangle^{(0)} = \mathrm{i} \delta_{a}^{d} \delta_{b}^{c} \delta_{j}^{i} \delta_{p}^{s} \delta_{r}^{q} \frac{\Gamma(\frac{3}{2} - \epsilon)}{2\pi^{\frac{3}{2} - \epsilon}} \frac{\gamma_{\mu\alpha}{}^{\beta}(x - y)^{\mu}}{|x - y|^{3 - 2\epsilon}}$$

$$(E.5)$$

whereas their one-loop corrections read

$$\langle A_{\mu}^{(a)}{}_{p}{}^{q}(x)A_{\nu}^{(a)}{}_{r}{}^{s}(y)\rangle^{(1)} = \delta_{p}^{s}\delta_{r}^{q}\sum_{b}\frac{(N_{ab}+N_{ba})N_{b}}{k_{a}}\frac{\Gamma^{2}(\frac{1}{2}-\epsilon)}{4\pi^{1-2\epsilon}}\left(\frac{\delta_{\mu\nu}}{|x-y|^{2-4\epsilon}} - \frac{\partial_{\mu}\partial_{\nu}|x-y|^{4\epsilon}}{4\epsilon(1+2\epsilon)}\right)$$

$$\langle \zeta_{(ab)p}^{i}{}_{\alpha}(x)\bar{\zeta}_{j}^{(cd)}{}_{r}{}^{s\beta}(y)\rangle^{(1)} = -\mathrm{i}\delta_{a}^{d}\delta_{b}^{c}\delta_{j}^{i}\delta_{p}^{s}\delta_{r}^{q}\delta_{a}^{\beta}\left(\frac{N_{a}}{k_{a}} + \frac{N_{b}}{k_{b}}\right)\frac{\Gamma^{2}(\frac{1}{2}-\epsilon)}{8\pi^{2-2\epsilon}}\frac{1}{|x-y|^{2-4\epsilon}} \tag{E.6}$$

Here the latic indices p, q, r, s are color indices.




Figure 11. Above: The tree propagators of gauge, scalar and fermionic fields. Below: The one–loop propagators of gauge and fermionic fields, respectively.

From the lagrangians in (E.2) the cubic vertices of figure 12 are given by

$$-\sum_{a} \frac{k_{a}}{6\pi} \int d^{3}x \varepsilon^{\mu\nu\rho} A_{\mu}^{(a)}{}_{p}{}^{q}(x) A_{\nu}^{(a)}{}_{q}{}^{r}(x) A_{\rho}^{(a)}{}_{r}{}^{p}(x)$$

$$-\sum_{a,b} \int d^{3}x \bar{\zeta}_{i}^{(ba)}{}_{p}{}^{q}(x) \gamma^{\mu} [A_{\mu}^{(a)}{}_{q}{}^{r}(x) \zeta_{(ab)}^{i}{}_{r}{}^{p}(x) - \zeta_{(ab)q}^{i}{}_{q}{}^{r}(x) A_{\mu}^{(b)}{}_{r}{}^{p}(x)]$$
(E.7)




Figure 12. The pure gauge vertex, and the mixed gauge–fermion vertex coming from the minimal coupling lagrangian.

F Details on the perturbative computation

In this appendix we give the explicit expressions of the intergrals corresponding to diagrams in figures 3–8. The intergrals are defined as

$$(5.1)$$

$$= \sum_{a} \frac{N_a^2}{k_a} \mathcal{I}_3^{(f)}$$

$$\underbrace{4(\mathbf{a})} = \sum_{a,b} \frac{(N_{ab} + N_{ba})N_a^2 N_b}{k_a^2} \mathcal{I}_{4(\mathbf{a})}^{(f)}, \quad \underbrace{4(\mathbf{b})} = \sum_a \frac{N_a^3 - N_a}{k_a^2} \mathcal{I}_{4(\mathbf{b})}^{(f)}$$

$$\underbrace{4(\mathbf{c})} = \sum_{a} \frac{N_a^3}{k_a^2} \mathcal{I}_{4(\mathbf{c})}^{(f)} + \sum_{a} \frac{N_a}{k_a^2} \mathcal{J}_{4(\mathbf{c})}^{(f)}, \quad \underbrace{4(\mathbf{d})} = \sum_{a,b} \frac{(N_{ab} + N_{ba})N_a^2 N_b}{k_a^2} \mathcal{I}_{4(\mathbf{d})}^{(f)} \qquad (F.2)$$

$$\underbrace{5} = \sum_{a,b} \bar{m}_i^{ab} n_{ba}^i N_a N_b \mathcal{I}_5^{(f)}$$
(F.3)

$$\widehat{\text{(6(a)}} = \sum_{a,b} (\bar{m}_i^{ab} n_{ba}^i - \bar{m}_i^{ba} n_{ab}^i) \frac{N_a^2 N_b}{k_a} \mathcal{I}_{\text{6(a)}}^{(f)}$$

$$(6(b)) = \sum_{a,b} (\bar{m}_i^{ab} n_{ba}^i + \bar{m}_i^{ba} n_{ab}^i) \frac{N_a^2 N_b}{k_a} \mathcal{I}_{6(b)}^{(f)}$$

$$(6(c)) = \sum_{a.b} (\bar{m}_i^{ab} n_{ba}^i \mathcal{I}_{6(c)}^{(f)} + \bar{m}_i^{ba} n_{ab}^i \mathcal{J}_{6(c)}^{(f)}) \frac{N_a^2 N_b}{k_a}$$

$$(6(d)) = \sum_{a,b} (\bar{m}_i^{ab} n_{ba}^i \mathcal{I}_{6(d)}^{(f)} + \bar{m}_i^{ba} n_{ab}^i \mathcal{J}_{6(d)}^{(f)}) \frac{N_a^2 N_b}{k_a}$$
 (F.4)

$$(7(a)) = \sum_{a,b,c} (\bar{m}_i^{ab} n_{ba}^i \bar{m}_j^{ac} n_{ca}^j + \bar{m}_i^{ba} n_{ab}^i \bar{m}_j^{ca} n_{ac}^j) N_a N_b N_c \mathcal{I}_{7(a)}^{(f)}$$

$$\underbrace{7(\mathbf{b})}_{a,b,c} = \sum_{a,b,c} (\bar{m}_i^{ab} n_{ba}^i \bar{m}_j^{ac} n_{ca}^j \mathcal{I}_{7(\mathbf{b})}^{(f)} + \bar{m}_i^{ba} n_{ab}^i \bar{m}_j^{ca} n_{ac}^j \mathcal{J}_{7(\mathbf{b})}^{(f)}) N_a N_b N_c$$
(F.5)

$$\underbrace{8 \text{(a)}} = \Big[\sum_{a,b} 2 (\bar{m}_i^{ab} n_{ba}^i)^3 N_a N_b + \sum_{a,b,c,d} (\bar{m}_i^{ab} n_{ba}^i \bar{m}_j^{ac} n_{ca}^j \bar{m}_k^{ad} n_{da}^k \Big]$$

$$+ \bar{m}_{i}^{ba} n_{ab}^{i} \bar{m}_{j}^{ca} n_{ac}^{j} \bar{m}_{k}^{da} n_{ad}^{k}) N_{a} N_{b} N_{c} N_{d} \Big] \mathcal{I}_{8(\mathbf{a})}^{(f)}$$

$$(8(b)) = + \sum_{a,b,c,d} \bar{m}_{i}^{ab} n_{ba}^{i} \bar{m}_{j}^{ac} n_{ca}^{j} \bar{m}_{k}^{db} n_{bd}^{k} N_{a} N_{b} N_{c} N_{d} \mathcal{I}_{8(b)}^{(f)} + \sum_{a,b} (\bar{m}_{i}^{ab} n_{ba}^{i})^{3} N_{a} N_{b} \mathcal{J}_{8(b)}^{(f)}$$

$$\underbrace{8(\mathbf{c})} = \sum_{a,b,c,d} [\bar{m}_{i}^{ab} n_{ba}^{i} \bar{m}_{j}^{ac} n_{ca}^{j} \bar{m}_{k}^{ad} n_{da}^{k} \mathcal{I}_{8(\mathbf{c})}^{(f)} + \bar{m}_{i}^{ba} n_{ab}^{i} \bar{m}_{j}^{ca} n_{ac}^{j} \bar{m}_{k}^{da} n_{ad}^{k} \mathcal{J}_{8(\mathbf{c})}^{(f)}$$

$$+\bar{m}_{i}^{ab}n_{ba}^{i}\bar{m}_{j}^{ac}n_{ca}^{j}\bar{m}_{k}^{db}n_{bd}^{k}\mathcal{K}_{8(c)}^{(f)}]N_{a}N_{b}N_{c}N_{d} + \sum_{a,b}(\bar{m}_{i}^{ab}n_{ba}^{i})^{3}N_{a}N_{b}\mathcal{L}_{8(c)}^{(f)}$$
(F.6)

Using the Feynman rules in appendix E, the integral from figure 3 is given by

$$\mathcal{I}_{3}^{(f)} = -i \frac{\Gamma(\frac{3}{2} - \epsilon)}{\pi^{\frac{1}{2} - \epsilon}} \oint d\tau_{1>2} \frac{\varepsilon_{\mu\nu\rho} \dot{x}_{1}^{\mu} \dot{x}_{2}^{\nu} x_{12}^{\rho}}{|x_{12}|^{3-2\epsilon}}$$
 (F.7)

where we have defined $x_i \equiv x(\tau_i)$, $x_{ij} \equiv x_i - x_j$ and $\oint d\tau_{1>2}$ means a double contour integral over $\tau_1 > \tau_2$.

With similar notations, from figure 4 we obtain

$$\mathcal{I}_{4(a)}^{(f)} = -\frac{\Gamma^{2}(\frac{1}{2} - \epsilon)}{4\pi^{1-2\epsilon}} \oint d\tau_{1>2} \frac{\dot{x}_{1} \cdot \dot{x}_{2}}{|x_{12}|^{2-4\epsilon}} \\
\mathcal{I}_{4(b)}^{(f)} = -\frac{\Gamma^{3}(\frac{3}{2} - \epsilon)}{2\pi^{\frac{5}{2} - 3\epsilon}} \oint d\tau_{1>2>3} \int d^{3}x \frac{\dot{x}_{1}^{\mu} \dot{x}_{2}^{\nu} \dot{x}_{3}^{\rho} \varepsilon^{\alpha\beta\gamma} \varepsilon_{\mu\alpha\sigma} \varepsilon_{\nu\beta\lambda} \varepsilon_{\rho\gamma\eta} (x - x_{1})^{\sigma} (x - x_{2})^{\lambda} (x - x_{3})^{\eta}}{|x - x_{1}|^{3-2\epsilon} |x - x_{2}|^{3-2\epsilon} |x - x_{3}|^{3-2\epsilon}} \\
\mathcal{I}_{4(c)}^{(f)} = -\frac{\Gamma^{2}(\frac{3}{2} - \epsilon)}{\pi^{1-2\epsilon}} \oint d\tau_{1>2>3>4} \left(\frac{\varepsilon_{\mu\nu\lambda} \dot{x}_{1}^{\mu} \dot{x}_{2}^{\nu} \dot{x}_{12}^{\lambda} \varepsilon_{\rho\sigma\eta} \dot{x}_{3}^{\rho} \dot{x}_{4}^{\sigma} \dot{x}_{34}^{\eta}}{|x_{12}|^{3-2\epsilon} |x_{34}|^{3-2\epsilon}} + (1432) \right) \\
\mathcal{J}_{4(c)}^{(f)} = -\frac{\Gamma^{2}(\frac{3}{2} - \epsilon)}{\pi^{1-2\epsilon}} \oint d\tau_{1>2>3>4} \frac{\varepsilon_{\mu\nu\lambda} \dot{x}_{1}^{\mu} \dot{x}_{3}^{\nu} \dot{x}_{13}^{\lambda} \varepsilon_{\rho\sigma\eta} \dot{x}_{2}^{\rho} \dot{x}_{4}^{\sigma} \dot{x}_{24}^{\eta}}{|x_{13}|^{3-2\epsilon} |x_{24}|^{3-2\epsilon}} \\
\mathcal{I}_{4(d)}^{(f)} = \frac{\Gamma^{2}(\frac{1}{2} - \epsilon)}{4\pi^{1-2\epsilon}} \oint d\tau_{1>2} \frac{|\dot{x}_{1}||\dot{x}_{2}|}{|x_{12}|^{2-4\epsilon}} \tag{F.8}$$

with the symbol (1423) in $\mathcal{I}_{4(c)}^{(f)}$ indicating the term obtained from the first one by permuting $\tau_{1,2,3,4} \to \tau_{1,4,2,3}$.

Similarly, from figures 5-8 we obtain

$$\mathcal{I}_{5}^{(f)} = -i \frac{\Gamma(\frac{3}{2} - \epsilon)}{2\pi^{\frac{3}{2} - \epsilon}} \oint d\tau_{1>2} \left(\frac{|\dot{x}_{1}||\dot{x}_{2}|u_{+}(\tau_{1})\gamma_{\mu}u_{-}(\tau_{2})x_{12}^{\mu}}{|x_{12}|^{3-2\epsilon}} - (21) \right)$$
 (F.9)

$$\begin{split} &\mathcal{I}_{6(\mathbf{a})}^{(f)} = -\frac{\Gamma^2(\frac{1}{2} - \epsilon)}{4\pi^2 - 2\epsilon} \oint d\tau_{1>2} \frac{|\dot{x}_1||\dot{x}_2|}{|x_{12}|^{2-4\epsilon}} \\ &\mathcal{I}_{6(\mathbf{b})}^{(f)} = \mathrm{i} \frac{\Gamma^2(\frac{1}{2} - \epsilon)}{8\pi^2 - 2\epsilon} \oint d\tau_{1>2} \left(\frac{|\dot{x}_1||\dot{x}_2|u_+(\tau_1)u_-(\tau_2)}{|x_{12}|^{2-4\epsilon}} - (21) \right) \\ &\mathcal{I}_{6(\mathbf{c})}^{(f)} = -\frac{\Gamma^3(\frac{3}{2} - \epsilon)}{4\pi^{\frac{7}{2} - 3\epsilon}} \oint d\tau_{1>2>3} \int d^3x \left[\frac{\dot{x}_1^{\mu} \varepsilon_{\mu\nu\rho}(x - x_1)^{\rho}}{|x - x_1|^{3-2\epsilon}} \right. \\ &\qquad \times \frac{|\dot{x}_2||\dot{x}_3|u_+(\tau_2)\gamma_\sigma\gamma^{\nu}\gamma_\lambda u_-(\tau_3)(x - x_2)^{\sigma}(x - x_3)^{\lambda}}{|x - x_2|^{3-2\epsilon}|x - x_3|^{3-2\epsilon}} - (231) + (312) \right] \\ &\mathcal{J}_{6(\mathbf{c})}^{(f)} = -\frac{\Gamma^3(\frac{3}{2} - \epsilon)}{4\pi^{\frac{5}{2} - 3\epsilon}} \oint d\tau_{1>2>3} \int d^3x \left[\frac{\dot{x}_1^{\mu} \varepsilon_{\mu\nu\rho}(x - x_1)^{\rho}}{|x - x_1|^{3-2\epsilon}} \right. \\ &\qquad \times \frac{|\dot{x}_3||\dot{x}_2|u_+(\tau_3)\gamma_\sigma\gamma^{\nu}\gamma_\lambda u_-(\tau_2)(x - x_3)^{\sigma}(x - x_2)^{\lambda}}{|x - x_3|^{3-2\epsilon}|x - x_2|^{3-2\epsilon}} - (231) + (312) \right] \\ &\mathcal{I}_{6(\mathbf{d})}^{(f)} = -\frac{\Gamma^2(\frac{3}{2} - \epsilon)}{2\pi^{2-2\epsilon}} \oint d\tau_{1>2>3>4} \left(\frac{\dot{x}_1^{\mu} \dot{x}_2^{\nu}|\dot{x}_3||\dot{x}_4|\varepsilon_{\mu\nu\rho}x_{12}^{\rho}u_+(\tau_3)\gamma_\sigma u_-(\tau_4)x_{34}^{\sigma}}{|x_{12}|^{3-2\epsilon}|x_{34}|^{3-2\epsilon}} \right. \\ &\qquad + (3412) + (4123) - (2341) \right) \\ &\mathcal{I}_{7(\mathbf{a})}^{(f)} = -\frac{\Gamma^2(\frac{3}{2} - \epsilon)}{2\pi^{2-2\epsilon}} \oint d\tau_{1>2>3>4} \left(\frac{\dot{x}_1^{\mu} \dot{x}_2^{\nu}|\dot{x}_4||\dot{x}_3|\varepsilon_{\mu\nu\rho}x_{12}^{\rho}u_+(\tau_4)\gamma_\sigma u_-(\tau_3)x_{43}^{\sigma}}{|x_{12}|^{3-2\epsilon}|x_{34}|^{3-2\epsilon}} \right. \\ &\qquad + (3412) + (4123) - (2341) \right) \end{aligned} \tag{F.10} \\ &\mathcal{I}_{7(\mathbf{b})}^{(f)} = -\frac{\Gamma^2(\frac{3}{2} - \epsilon)}{16\pi^{3-2\epsilon}} \oint d\tau_{1>2>3>4} \left(\frac{|\dot{x}_1||\dot{x}_2|}{|x_{12}|^{3-2\epsilon}|x_{34}|^{3-2\epsilon}} \right. \\ &\qquad \times \frac{|\dot{x}_3||\dot{x}_4|u_+(\tau_3)\gamma_\nu u_-(\tau_4)x_{34}^{\sigma}}{|x_{34}|^{3-2\epsilon}} - (2341) \right) \\ &\qquad \mathcal{I}_{7(\mathbf{b})}^{(f)} = -\frac{\Gamma^2(\frac{3}{2} - \epsilon)}{4\pi^{3-2\epsilon}} \oint d\tau_{1>2>3>4} \left(\frac{|\dot{x}_1||\dot{x}_2|u_+(\tau_1)\gamma_\mu u_-(\tau_2)x_{12}^{\mu}}{|x_{12}|^{3-2\epsilon}} \right. \\ &\qquad \times \frac{|\dot{x}_3||\dot{x}_4|u_+(\tau_3)\gamma_\nu u_-(\tau_4)x_{34}^{\sigma}}{|x_{34}|^{3-2\epsilon}} - (2341) \right) \\ &\qquad \mathcal{I}_{7(\mathbf{b})}^{(f)} = -\frac{\Gamma^2(\frac{3}{2} - \epsilon)}{4\pi^{3-2\epsilon}} \oint d\tau_{1>2>3>4} \left(\frac{|\dot{x}_1||\dot{x}_2|u_+(\tau_1)\gamma_\mu u_-(\tau_1)x_{12}^{\mu}}{|x_{12}|^{3-2\epsilon}} \right. \\ &\qquad \times \frac{|\dot{x}_3||\dot{x}_3|u_+(\tau_4)\gamma_\nu u_-(\tau_3)x_{34}^{\sigma}}{|x_{12}|^{3-2\epsilon}} - (2341) \right) \end{aligned}$$

$$\begin{split} \mathcal{I}_{8(a)}^{(f)} &= -\frac{\Gamma^{3}(\frac{1}{2}-\epsilon)}{64\pi^{\frac{9}{2}-3\epsilon}} \oint d\tau_{1>2>3} \frac{|\dot{x}_{1}||\dot{x}_{2}||\dot{x}_{3}|}{|x_{12}|^{1-2\epsilon}|x_{13}|^{1-2\epsilon}|x_{23}|^{1-2\epsilon}} \\ \mathcal{I}_{8(b)}^{(f)} &= -\mathrm{i} \frac{\Gamma^{2}(\frac{1}{2}-\epsilon)\Gamma(\frac{3}{2}-\epsilon)}{32\pi^{\frac{9}{2}-3\epsilon}} \oint d\tau_{1>2>3>4} \left(\frac{|\dot{x}_{1}||\dot{x}_{2}||\dot{x}_{3}||\dot{x}_{4}|u_{+}(\tau_{3})\gamma_{\mu}u_{-}(\tau_{4})x_{34}^{\mu}}{|x_{12}|^{2-4\epsilon}|x_{34}|^{3-2\epsilon}} \right. \\ &\quad + (3412) - (4123) - (2341) - (2143) - (4321) - (1432) + (3214) \right) \\ \mathcal{J}_{8(b)}^{(f)} &= -\mathrm{i} \frac{\Gamma^{2}(\frac{1}{2}-\epsilon)\Gamma(\frac{3}{2}-\epsilon)}{32\pi^{\frac{9}{2}-3\epsilon}} \oint d\tau_{1>2>3>4} \left(\frac{|\dot{x}_{1}||\dot{x}_{2}||\dot{x}_{4}|u_{+}(\tau_{2})\gamma_{\mu}u_{-}(\tau_{4})x_{24}^{\mu}}{|x_{13}|^{2-4\epsilon}|x_{24}|^{3-2\epsilon}} \right. \\ &\quad - (2341) - (1432) + (2143) \right) \\ \mathcal{I}_{8(c)}^{(f)} &= \mathrm{i} \frac{\Gamma^{3}(\frac{3}{2}-\epsilon)}{8\pi^{\frac{9}{2}-3\epsilon}} \oint d\tau_{1>2>3>4>5>6} \left(\frac{|\dot{x}_{1}||\dot{x}_{2}|u_{+}(\tau_{1})\gamma_{\mu}u_{-}(\tau_{2})x_{12}^{\mu}}{|x_{12}|^{3-2\epsilon}} \right. \\ &\quad \times \frac{|\dot{x}_{3}||\dot{x}_{4}|u_{+}(\tau_{3})\gamma_{\mu}u_{-}(\tau_{4})x_{34}^{\mu}}{|x_{34}|^{3-2\epsilon}} \frac{|\dot{x}_{5}||\dot{x}_{6}|u_{+}(\tau_{5})\gamma_{\mu}u_{-}(\tau_{6})x_{56}^{\mu}}{|x_{56}|^{3-2\epsilon}} - (234561) \right) \\ \mathcal{J}_{8(c)}^{(f)} &= -\mathrm{i} \frac{\Gamma^{3}(\frac{3}{2}-\epsilon)}{8\pi^{\frac{9}{2}-3\epsilon}} \oint d\tau_{1>2>3>4>5>6} \left(\frac{|\dot{x}_{2}||\dot{x}_{1}|u_{+}(\tau_{2})\gamma_{\mu}u_{-}(\tau_{1})x_{21}^{\mu}}{|x_{21}|^{3-2\epsilon}} \right. \\ &\quad \times \frac{|\dot{x}_{4}||\dot{x}_{3}|u_{+}(\tau_{4})\gamma_{\mu}u_{-}(\tau_{3})x_{43}^{\mu}}{|x_{3}|^{3-2\epsilon}} \frac{|\dot{x}_{6}||\dot{x}_{5}|u_{+}(\tau_{6})\gamma_{\mu}u_{-}(\tau_{5})x_{65}^{\mu}}{|x_{65}|^{3-2\epsilon}} - (234561) \right) \\ \mathcal{K}_{8(c)}^{(f)} &= -\mathrm{i} \frac{\Gamma^{3}(\frac{3}{2}-\epsilon)}{8\pi^{\frac{9}{2}-3\epsilon}} \oint d\tau_{1>2>3>4>5>6} \left(\frac{|\dot{x}_{1}||\dot{x}_{2}|u_{+}(\tau_{1})\gamma_{\mu}u_{-}(\tau_{2})x_{25}^{\mu}}{|x_{12}|^{3-2\epsilon}} \right. \\ &\quad \times \frac{|\dot{x}_{3}||\dot{x}_{6}|u_{+}(\tau_{3})\gamma_{\mu}u_{-}(\tau_{6})x_{36}^{\mu}}{|x_{36}|^{3-2\epsilon}} \frac{|\dot{x}_{5}||\dot{x}_{4}|u_{+}(\tau_{5})\gamma_{\mu}u_{-}(\tau_{4})x_{24}^{\mu}}{|x_{14}|^{3-2\epsilon}} \\ &\quad \times \frac{|\dot{x}_{3}||\dot{x}_{6}|u_{+}(\tau_{5})\gamma_{\mu}u_{-}(\tau_{6})x_{36}^{\mu}}{|x_{36}|^{3-2\epsilon}} \frac{|\dot{x}_{5}||\dot{x}_{5}|u_{+}(\tau_{7})\gamma_{\mu}u_{-}(\tau_{4})x_{24}^{\mu}}{|x_{14}|^{3-2\epsilon}} \\ &\quad \times \frac{|\dot{x}_{5}||\dot{x}_{2}|u_{+}(\tau_{5})\gamma_{\mu}u_{-}(\tau_{6})x_{36}^{\mu}}{|x_{5}|^{3-2\epsilon}} \frac{|\dot{x}_{5}||\dot{x}_{5}|u_{+}(\tau_{5})\gamma_{\mu}u_{-}(\tau_{6})x_{36}^{\mu}}{|x_{5}$$

References

- [1] V. Pestun, Localization of gauge theory on a four-sphere and supersymmetric Wilson loops, Commun. Math. Phys. 313 (2012) 71–129, [0712.2824].
- [2] A. Kapustin, B. Willett and I. Yaakov, Exact Results for Wilson Loops in Superconformal Chern-Simons Theories with Matter, JHEP 1003 (2010) 089, [0909.4559].
- [3] J. M. Maldacena, The Large N limit of superconformal field theories and supergravity, Adv. Theor. Math. Phys. 2 (1998) 231–252, [hep-th/9711200].
- [4] S. Gubser, I. R. Klebanov and A. M. Polyakov, Gauge theory correlators from noncritical string theory, Phys. Lett. B428 (1998) 105–114, [hep-th/9802109].
- [5] E. Witten, Anti-de Sitter space and holography, Adv. Theor. Math. Phys. 2 (1998) 253–291, [hep-th/9802150].

- [6] O. Aharony, O. Bergman, D. L. Jafferis and J. Maldacena, N=6 superconformal Chern-Simons-matter theories, M2-branes and their gravity duals, JHEP 0810 (2008) 091, [0806.1218].
- [7] O. Aharony, O. Bergman and D. L. Jafferis, Fractional M2-branes, JHEP 0811 (2008) 043, [0807.4924].
- [8] D. Gaiotto and X. Yin, Notes on superconformal Chern-Simons-Matter theories, JHEP 0708 (2007) 056, [0704.3740].
- [9] D. Berenstein and D. Trancanelli, *Three-dimensional N=6 SCFT's and their membrane dynamics*, *Phys. Rev.* **D78** (2008) 106009, [0808.2503].
- [10] N. Drukker, J. Plefka and D. Young, Wilson loops in 3-dimensional N=6 supersymmetric Chern-Simons Theory and their string theory duals, JHEP 0811 (2008) 019, [0809.2787].
- [11] B. Chen and J.-B. Wu, Supersymmetric Wilson Loops in N=6 Super Chern-Simons-matter theory, Nucl. Phys. B825 (2010) 38–51, [0809.2863].
- [12] S.-J. Rey, T. Suyama and S. Yamaguchi, Wilson Loops in Superconformal Chern-Simons Theory and Fundamental Strings in Anti-de Sitter Supergravity Dual, JHEP 0903 (2009) 127, [0809.3786].
- [13] N. Drukker and D. Trancanelli, A Supermatrix model for $\mathcal{N}=6$ super Chern-Simons-matter theory, JHEP 1002 (2010) 058, [0912.3006].
- [14] H. Ouyang, J.-B. Wu and J.-j. Zhang, Supersymmetric Wilson loops in $\mathcal{N}=4$ super Chern-Simons-matter theory, JHEP 1511 (2015) 213, [1506.06192].
- [15] M. Cooke, N. Drukker and D. Trancanelli, A profusion of 1/2 BPS Wilson loops in $\mathcal{N}=4$ Chern-Simons-matter theories, JHEP 1510 (2015) 140, [1506.07614].
- [16] H. Ouyang, J.-B. Wu and J.-j. Zhang, Novel BPS Wilson loops in three-dimensional quiver Chern-Simons-matter theories, Phys. Lett. B753 (2016) 215–220, [1510.05475].
- [17] H. Ouyang, J.-B. Wu and J.-j. Zhang, Construction and classification of novel BPS Wilson loops in quiver Chern-Simons-matter theories, Nucl. Phys. B910 (2016) 496–527, [1511.02967].
- [18] A. Mauri, S. Penati and J.-j. Zhang, New BPS Wilson loops in $\mathcal{N}=4$ circular quiver Chern-Simons-matter theories, JHEP 1711 (2017) 174, [1709.03972].
- [19] V. Cardinali, L. Griguolo, G. Martelloni and D. Seminara, New supersymmetric Wilson loops in ABJ(M) theories, Phys. Lett. B718 (2012) 615–619, [1209.4032].
- [20] M. S. Bianchi, L. Griguolo, M. Leoni, S. Penati and D. Seminara, *BPS Wilson loops and Bremsstrahlung function in ABJ(M): a two loop analysis*, *JHEP* **1406** (2014) 123, [1402.4128].
- [21] M. S. Bianchi, L. Griguolo, A. Mauri, S. Penati and D. Seminara, A matrix model for the latitude Wilson loop in ABJM theory, 1802.07742.
- [22] M. Benna, I. Klebanov, T. Klose and M. Smedback, Superconformal Chern-Simons Theories and AdS₄/CFT₃ Correspondence, JHEP **0809** (2008) 072, [0806.1519].
- [23] M. Marino and P. Putrov, Exact Results in ABJM Theory from Topological Strings, JHEP 1006 (2010) 011, [0912.3074].
- [24] N. Drukker, M. Marino and P. Putrov, From weak to strong coupling in ABJM theory, Commun. Math. Phys. 306 (2011) 511–563, [1007.3837].

- [25] M. S. Bianchi, G. Giribet, M. Leoni and S. Penati, 1/2 BPS Wilson loop in N=6 superconformal Chern-Simons theory at two loops, Phys. Rev. D88 (2013) 026009, [1303.6939].
- [26] M. S. Bianchi, G. Giribet, M. Leoni and S. Penati, The 1/2 BPS Wilson loop in ABJ(M) at two loops: The details, JHEP 1310 (2013) 085, [1307.0786].
- [27] L. Griguolo, G. Martelloni, M. Poggi and D. Seminara, Perturbative evaluation of circular 1/2 BPS Wilson loops in N=6 Super Chern-Simons theories, JHEP 1309 (2013) 157, [1307.0787].
- [28] L. Griguolo, M. Leoni, A. Mauri, S. Penati and D. Seminara, Probing Wilson loops in N = 4 Chern-Simons-matter theories at weak coupling, Phys. Lett. B753 (2016) 500-505, [1510.08438].
- [29] M. S. Bianchi, L. Griguolo, M. Leoni, A. Mauri, S. Penati and D. Seminara, Framing and localization in Chern-Simons theories with matter, JHEP 1606 (2016) 133, [1604.00383].
- [30] M. S. Bianchi, A note on multiply wound BPS Wilson loops in ABJM, JHEP 1609 (2016) 047, [1605.01025].
- [31] M. S. Bianchi, L. Griguolo, M. Leoni, A. Mauri, S. Penati and D. Seminara, *The quantum* 1/2 BPS Wilson loop in $\mathcal{N}=4$ Chern-Simons-matter theories, JHEP **1609** (2016) 009, [1606.07058].
- [32] K.-M. Lee and S. Lee, 1/2-BPS Wilson loops and vortices in ABJM model, JHEP 1009 (2010) 004, [1006.5589].
- [33] M. Lietti, A. Mauri, S. Penati and J.-j. Zhang, String theory duals of Wilson loops from Higgsing, JHEP 1708 (2017) 030, [1705.02322].
- [34] D. L. Jafferis, The Exact Superconformal R-Symmetry Extremizes Z, JHEP 1205 (2012) 159, [1012.3210].
- [35] N. Hama, K. Hosomichi and S. Lee, Notes on SUSY Gauge Theories on Three-Sphere, JHEP 1103 (2011) 127, [1012.3512].
- [36] D. L. Jafferis, I. R. Klebanov, S. S. Pufu and B. R. Safdi, Towards the F-Theorem: N=2 Field Theories on the Three-Sphere, JHEP 1106 (2011) 102, [1103.1181].
- [37] J. M. Figueroa-O'Farrill, On the supersymmetries of Anti-de Sitter vacua, Class. Quant. Grav. 16 (1999) 2043–2055, [hep-th/9902066].
- [38] E. Guadagnini, M. Martellini and M. Mintchev, Wilson Lines in Chern-Simons Theory and Link Invariants, Nucl. Phys. B330 (1990) 575-607.
- [39] S.-J. Rey and J.-T. Yee, Macroscopic strings as heavy quarks in large N gauge theory and anti-de Sitter supergravity, Eur. Phys. J. C22 (2001) 379–394, [hep-th/9803001].
- [40] N. Drukker and B. Fiol, All-genus calculation of Wilson loops using D-branes, JHEP 02 (2005) 010, [hep-th/0501109].
- [41] S. Yamaguchi, Wilson loops of anti-symmetric representation and D5-branes, JHEP 0605 (2006) 037, [hep-th/0603208].
- [42] J. Gomis and F. Passerini, Holographic Wilson Loops, JHEP 0608 (2006) 074, [hep-th/0604007].
- [43] J. Gomis and F. Passerini, Wilson Loops as D3-Branes, JHEP 0701 (2007) 097, [hep-th/0612022].

- [44] S. Yamaguchi, Bubbling geometries for half BPS Wilson lines, Int. J. Mod. Phys. A22 (2007) 1353–1374, [hep-th/0601089].
- [45] O. Lunin, On gravitational description of Wilson lines, JHEP **06** (2006) 026, [hep-th/0604133].
- [46] E. D'Hoker, J. Estes and M. Gutperle, Gravity duals of half-BPS Wilson loops, JHEP 06 (2007) 063, [0705.1004].
- [47] Y. Hatsuda, M. Honda, S. Moriyama and K. Okuyama, ABJM Wilson Loops in Arbitrary Representations, JHEP 1310 (2013) 168, [1306.4297].
- [48] H. Ouyang, J.-B. Wu and J.-j. Zhang, BPS Wilson loops in Minkowski spacetime and Euclidean space, Eur. Phys. J. C75 (2015) 606, [1504.06929].
- [49] K. Hosomichi, K.-M. Lee, S. Lee, S. Lee and J. Park, N=5,6 Superconformal Chern-Simons Theories and M2-branes on Orbifolds, JHEP 0809 (2008) 002, [0806.4977].
- [50] D. Gaiotto and E. Witten, Janus Configurations, Chern-Simons Couplings, And The theta-Angle in N=4 Super Yang-Mills Theory, JHEP 1006 (2010) 097, [0804.2907].
- [51] K. Hosomichi, K.-M. Lee, S. Lee, S. Lee and J. Park, N=4 Superconformal Chern-Simons Theories with Hyper and Twisted Hyper Multiplets, JHEP 0807 (2008) 091, [0805.3662].
- [52] Y. Imamura and K. Kimura, On the moduli space of elliptic Maxwell-Chern-Simons theories, Prog. Theor. Phys. 120 (2008) 509-523, [0806.3727].
- [53] S. Terashima and F. Yagi, Orbifolding the Membrane Action, JHEP 0812 (2008) 041, [0807.0368].