Closed-form solutions of the nonlinear Schrödinger equation with arbitrary dispersion and potential

Andrei D. Polyanin^{*}, Nikolay A. Kudryashov^{**}

* Ishlinsky Institute for Problems in Mechanics, Russian Academy of Sciences, pr. Vernadskogo 101, bldg. 1, Moscow, 119526 Russia

 ** Department of Applied Mathematics, National Research Nuclear

University MEPhI, 31 Kashirskoe Shosse, 115409 Moscow, Russia

e-mails: polyanin@ipmnet.ru, nakudryashov@mephi.ru

For the first time, the general nonlinear Schrödinger equation is investigated, in which the chromatic dispersion and potential are specified by two arbitrary functions. The equation in question is a natural generalization of a wide class of related nonlinear partial differential equations that are often used in various areas of theoretical physics, including nonlinear optics, superconductivity and plasma physics. To construct exact solutions, a combination of the method of functional constraints and methods of generalized separation of variables is used. Exact closed-form solutions of the general nonlinear Schrödinger equation, which are expressed in quadratures or elementary functions, are found. One-dimensional non-symmetry reductions are described, which lead the considered nonlinear partial differential equation to a simpler ordinary differential equation or a system of such equations. The exact solutions obtained in this work can be used as test problems intended to assess the accuracy of numerical and approximate analytical methods for integrating nonlinear equations of mathematical physics.

Keywords: nonlinear Schrödinger equations, exact closed-form solutions, solutions in quadratures, solution in elementary functions, method of functional constraints, non-symmety reductions

1 Introduction

1.1 Nonlinear Schrödinger equations

1°. In many areas of theoretical physics, one encounters the classical nonlinear Schrödinger equation with cubic nonlinearity (see, for example, [1–5]):

$$iu_t + u_{xx} + k|u|^2 u = 0, (1)$$

where u = u(x, t) is the desired complex-valued function of real variables, the square of the module of which determines the intensity of light, t is the time, x is the spatial variable, k is a parameter, $i^2 = -1$.

Eq. (1) is used for mathematical modeling of wave propagation in almost all areas of physics where nonlinear wave processes are considered. The theoretical and experimental justification for the use of the nonlinear Schrödinger equation in non-linear optics is given in [6-8]. When describing the propagation of pulses in an optical fiber, the expression with the second derivative is responsible for the dispersion of the pulse, the quadratic function $g(|u|) = k|u|^2$, which sets the potential, is called the Kerr nonlinearity. It characterizes the interaction of a light pulse with a fiber material and specifies the law of refraction of light in a nonlinear medium. The uniqueness of Eq. (1) is explained not only by the fact that this partial differential equation (PDE) is the basic equation for describing the processes of information transfer in an optical medium, but also by the fact that it belongs to the class of integrable partial differential equations [5]. Eq. (1) has an infinite number of conservation laws, Bäcklund transformations, and passes the Painlevé test for partial differential equations [4, 5, 9-11]. The Cauchy problem for Eq. (1) with an initial condition of the general form is solved by the inverse scattering transform [4,5]. The exact solutions of the nonlinear Schrödinger equation (1) are given in |12-14|.

 2° . Related nonlinear partial differential equations of the form

$$iu_t + u_{xx} + g(|u|)u = 0, (2)$$

and other nonlinear Schrödinger-type equations that occur in the scientific literature can be found, for example, in [12–36] (see also Section 2 of this article).

In nonlinear optics, the potential g(|u|) in Eq. (2) determines the law of interaction of a light pulse with a fiber material. Exact solutions of Eq. (2) in the case of power dependence $g(|u|) = k|u|^n$ have been considered, for example, in [12–14]. In plasma theory and laser physics, the PDE (2) with $g(|u|) = k(1 - e^{a|u|})$ is encountered (see, for example, [37]). Exact closed-form solutions of the nonlinear Schrödinger equation (2) for arbitrary function g(|u|) are given in [12, 14].

In this article, we will consider a nonlinear PDE that is much more complex than equation (2). In this equation, instead of the linear dispersion term given by the second derivative u_{xx} , we will include a nonlinear dispersion of the form $[f(u)u_x]_x$, where f(u) is an arbitrary function. Thus, two arbitrary functions f(u)and g(u) will enter into the nonlinear PDE under consideration at once.

1.2 Exact solutions of nonlinear partial differential equations (terminology)

In this article, exact solutions of PDEs are understood as the following solutions [12, 38]:

(a) Solutions which are expressed via the elementary functions.

(b) Solutions which are expressed in quadratures, i.e. through elementary functions, functions included in the equation (this is necessary if the equation contains arbitrary or special functions), and indefinite integrals.

(c) Solutions that are expressed through solutions of ordinary differential equations (ODEs) or systems of such equations.

Various combinations of solutions described in Items (a)–(c) are also allowed. In cases (a) and (b), the exact solution can be presented in explicit, implicit, or parametric form.

Note that exact solutions play an important role as mathematical standards which are often used as test problems to check the adequacy and assess the accuracy of numerical and approximate analytical methods for integrating nonlinear PDEs. The most preferable for these purposes are simple solutions from Items (a) and (b). Several such exact solutions are described later in this article.

Remark 1. Of great interest are also solutions of nonlinear PDEs, which are

expressed through solutions of linear PDEs (examples of such nonlinear PDEs can be found, for example, in [12, 39, 40]), as well as conditionally integrable nonlinear PDEs [40] (which can also be called partially linearizable PDEs).

Exact closed-form solutions of nonlinear PDEs are most often constructed using the classical method of symmetry reductions (see, for example, [39,41–45]), the direct method of symmetry reductions (see, for example, [12, 38, 46–49]), the nonclassical methods of symmetry reductions (see, for example, [49–54]), methods of generalized separation of variables (see, for example, [12, 38, 45, 55–58]), methods of functional separation of variables (see, for example, [12, 38, 52, 59–62]), the method of differential constraints (see, for example, [12, 38, 48, 63–66]), the inverse scattering method (see, for example, [5, 67–70]), the Painlevé analysis of nonlinear PDEs (see, for example, [71–76]), and some other exact analytical methods (see, for example, [12, 77–79]).

It is important to note that for complex nonlinear PDEs depending on one or more arbitrary functions (and it is precisely such nonlinear PDEs that are considered in this article), the vast majority of existing analytical methods for constructing exact solutions are either not applicable at all or are weakly effective. Statistical processing of reference data [12,14] showed that at present the majority of exact solutions of such equations were obtained by methods of generalized and functional separation of variables. Significantly fewer exact solutions of such equations were obtained by nonclassical methods of symmetry reductions and the method of differential constraints, which are much more difficult to use in practice. In general, very few nonlinear PDEs depending on arbitrary functions that admit exact closed-form solutions are known today.

2 General nonlinear Schrödinger equation. Transformations to a system of real PDEs

2.1 The nonlinear Schrödinger equation with arbitrary dispersion and potential

 $1^\circ.$ In this paper we consider the nonlinear Schrödinger equation of the general form

$$iu_t + [f(|u|)u]_{xx} + g(|u|)u = 0, (3)$$

where u = u(x, t) is the complex-value function of real variables, f(z) and g(z) are arbitrary real functions (f is a twice continuously differentiable function, and g is a continuous function), $i^2 = -1$. Eq. (3) with nonlinear dispersion, which is given by the function f(z), is an essential generalization of Eq. (2) with f(z) = const.

The nonlinear Schrödinger equation (3) with power functions

$$f(z) = az^k, \quad g(z) = bz^m + cz^n \tag{4}$$

has been considered in [24], where its solutions of the type of stationary solitons of the form $u = r(x)e^{i\lambda t}$ were described. Some exact solutions of Eq. (3) in the case of arbitrary functions f(|u|) and $g(|u|) \equiv 0$ are presented in [12].

2°. Note that if u(x,t) is a solution of equation (3), than the functions

$$\bar{u} = \pm e^{iC_1} u(\pm x + C_2, t + C_3), \tag{5}$$

where C_1 , C_2 , C_3 are arbitrary real constants, are also solutions of this equation (the signs in Eq. (5) can be chosen independently of each other). From this property follows, that Eq. (3) admits traveling wave solutions of the form u = U(z), $z = x - \lambda t$, where λ is an arbitrary constant (more complex solutions including traveling wave solutions are given in Section 4).

2.2 Transformation of the general nonlinear Schrödinger equation to a system of real PDEs

Let us represent the desired function in exponential form

$$u = r e^{i\varphi}, \quad r = |u|, \tag{6}$$

where $r = r(x, t) \ge 0$ and $\varphi = \varphi(x, t)$ are real functions.

Differentiating (6), we find the derivatives:

$$u_{t} = (r_{t} + ir\varphi_{t})e^{i\varphi},$$

$$[f(|u|)u]_{x} = (h_{x} + ih\varphi_{x})e^{i\varphi}, \quad h = rf(r),$$

$$[f(|u|)u]_{xx} = [h_{xx} - h\varphi_{x}^{2} + i(2h_{x}\varphi_{x} + h\varphi_{xx})]e^{i\varphi}, \quad h = rf(r).$$
(7)

Now we substitute (7) into (3), and then divide all terms by $e^{i\varphi}$. Further equating the real and imaginary parts of the obtained relation to zero, we arrive at the following system of two real partial differential equations:

$$-r\varphi_t + h_{xx} - h\varphi_x^2 + rg(r) = 0,$$

$$r_t + 2h_x\varphi_x + h\varphi_{xx} = 0, \quad h = rf(r).$$
(8)

We use system of equations (8) and expressions (6) to construct exact solutions of the nonlinear Schrödinger equation (3).

Remark 2. Note that the second equation of system (8) can be written in the form of conservation law

$$H_t + (h^2 \varphi_x)_x = 0 \tag{9}$$

where the notation is used

$$H = \int h \, dr.$$

From equation (9), under some standard restrictions, the law of conservation of density follows, $\int_{-\infty}^{\infty} H \, dx = \text{const.}$ In addition, the conservation law (9) can be used to evaluate the accuracy of the results of applying numerical methods of integrating the system (8).

2.3 Method of constructing exact solutions based on functional constraints

The construction of exact closed-form solutions of Eq. (3) is complicated by the fact that it contains two arbitrary functions f(z) and g(z). For such equations and other nonlinear PDEs of the general form that contain arbitrary functions, it is impossible to directly use the standard methods of generalized separation of variables described in books [12, 38, 55]. To find exact solutions of the general nonlinear Schrödinger equation (3), we impose one of the three additional relations (functional constraints) on the argument of arbitrary functions:

$$|u| = \text{const},\tag{10}$$

$$|u| = p(x),\tag{11}$$

$$|u| = q(t), \tag{12}$$

where p(x) and q(t) are some function of one argument. When any of the relations (10)-(12) is satisfied, equation (3) is "linearized". This important fact allows us to further use the standard approach for separating variables applied for linear PDEs or methods of generalized separation of variables used for nonlinear PDEs [12,38,55]. A similar technique, based on the use of additional relations of the type (10)-(12) and called the method of functional constraints, has made it possible to find many exact solutions of nonlinear PDEs with delay [79–82].

After the transition from the complex equation (3) to the system of real PDEs (8), when constructing exact solutions, one should use the relations (10)–(12), setting in them |u| = r (this follows from representation (6)). Thus, the functions p(x) and q(t) in (11) and (12) depend on x and t implicitly and are expressed through the amplitude r.

Remark 3. In Section 4 (see Remark 4) it will be shown that in addition to (10)-(12), a more complex functional constraint based on the transition to a new variable of the traveling wave type can also be used to construct exact solutions.

3 Exact solutions of the general nonlinear Schrödinger equation

In this section exact closed-form solutions of the general nonlinear Schrödinger equation (3) with arbitrary functions f(z) and g(z) are given. To construct these exact solutions we use the functional constraints (10)–(12) and the methods of generalized separation of variables.

3.1 Traveling wave solutions with constant amplitude

First, we use the simplest additional relation (10). Given that |u| = r, we substitute $r = C_1 = \text{const}$ into system (8). From the second equation (8) it follows that $\varphi''_{xx} = 0$. From here, after integration, we have $\varphi(x) = a(t)x + b(t)$, where a(t) and b(t) are arbitrary functions. Substituting this expression into the first equation (8), we find $a(t) = C_2$ and $b(t) = Bt + C_3$. As a result, we arrive at the simple exact solution of the PDE system (8):

$$r = C_1, \quad \varphi = C_2 x + C_3 + Bt, \quad B = g(C_1) - C_2^2 f(C_1),$$
 (13)

where C_1 , C_2 , and C_3 are arbitrary real constants ($C_1 > 0$). Substituting (13) into (6), we have the traveling wave solution of the nonlinear PDE (3):

$$u = C_1 e^{i(C_2 x + C_3 + Bt)}, \quad B = g(C_1) - C_2^2 f(C_1).$$
(14)

This is a solution that is periodic in time and space with constant amplitude C_1 .

In the case of power functions (4) solution (14) takes the form

$$u = C_1 e^{i(C_2 x + C_3 + Bt)}, \quad B = bC_1^m + cC_1^n - aC_1^k C_2^2$$

3.2 Time-periodic solutions with amplitude depending on the spatial variable x

To construct another exact solution, we use the additional relation (11). Given that |u| = r, we substitute r = r(x) into system (8). Integrating the second equation of the resulting system twice, we have

$$\varphi = a(t) \int h^{-2} dx + b(t), \qquad (15)$$

where a = a(t) and b = b(t) are arbitrary functions. By eliminating φ from the first equation of system (8) using (15), we arrive at an equation that, after dividing all terms by r, takes the form

$$-a'_t \int h^{-2} dx - b'_t - a^2 r^{-1} h^{-3} + r^{-1} [h''_{xx} + rg(r)] = 0, \quad h = rf(r).$$
(16)

To solve the functional differential equation (16), which includes the desired functions depending on different variables, we use the methods of generalized

separation of variables [12,38]. Standard analysis shows that two different situations are possible, described below.

1°. The general case when two functions f and g are arbitrary. Setting in (16)

$$a(t) = C_2, \quad b(t) = C_1 t + C_3,$$
(17)

where C_1 , C_2 , and C_3 are arbitrary constants, we obtain a second-order nonlinear ODE of the autonomous form

$$h''_{xx} - C_2^2 h^{-3} - C_1 r + rg(r) = 0, \quad h = rf(r).$$
(18)

Let us first consider the degenerate case when $h''_{xx} \equiv 0$, i.e.

$$h(x) = ax + b, (19)$$

where a and b are arbitrary constants. We substitute the function (19) into (18) and take into account the relation h = rf(r). As a result, we obtain a potential, which in this case is expressed through the function f(r) as

$$g(r) = C_1 + C_2^2 r^{-4} f^{-3}(r).$$
(20)

It follows that of the nonlinear Schrödinger equation

$$iu_t + [f(|u|)u]_{xx} + [C_1 + C_2^2|u|^{-4}f^{-3}(|u|)]u = 0,$$
(21)

where f(r) is an arbitrary function, has an exact solution of the form (6), in which the amplitude r = r(x) is given by the implicit relation

$$ax + b = rf(r), (22)$$

and the phase is found using formula (25) and can be written in the form

$$\varphi = C_1 t + C_2 \int h^{-2} dx + C_3 = C_1 t - \frac{C_2}{a(ax+b)} + C_3.$$
(23)

In the general case of two arbitrary functions f(r) and g(r), the general solution of the second-order nonlinear ODE (18) can be expressed in quadratures in implicit form (detailed explanations are given in Appendix):

$$\int \left[-C_2^2 h^{-2} + 2 \int [C_1 r - rg(r)] \, dh + C_4 \right]^{-1/2} dh = C_5 \pm x,$$

$$h = rf(r), \quad dh = [f(r) + rf'_r(r)] \, dr,$$
(24)

where C_4 and C_5 are arbitrary constants.

In this case, the phase φ is expressed through the amplitude r by the formulas

$$\varphi = C_1 t + C_2 \int h^{-2} dx + C_3 = C_1 t + C_2 \int [rf(r)]^{-2} dx + C_3, \qquad (25)$$

which are obtained by substituting the expressions (17) into (15).

Thus, we have shown that the PDE system (8) admits a time-periodic solution, which is expressed in quadratures using the relations (24) and (25).

Setting $C_2 = 0$ in (24) and (25), we obtain solutions describing stationary solitons, whose phase φ does not depend on the spatial coordinate x.

Example 1. Let us consider the nonlinear Schrödinger equation (3) with a constant potential,

$$g(r) = C_1 = \text{const.} \tag{26}$$

Substituting (26) into solution (24), after integration we have

$$\sqrt{C_4 h^2 - C_2^2} = C_4 (C_5 \pm x). \tag{27}$$

By squaring both parts (26) and taking into account the formula h = rf(r), we obtain the relation

$$C_4 r^2 f^2(r) = C_2^2 + C_4^2 (C_5 \pm x)^2, \qquad (28)$$

which implicitly determines the dependence of the amplitude r on x. Using formulas (25) and (27), we find the phase

$$\varphi = C_1 t + \arctan\left[\frac{C_4(x \pm C_5)}{C_2}\right] + C_3.$$
(29)

Note that in equation (3) (with (26)) and solution (28) the function f = f(r) can be specified arbitrarily.

The general solution of equation (18), represented in quadratures in the implicit form (24), is difficult to analyze due to the presence of two arbitrary functions in it f(r) and g(r). Therefore, we will describe here a simpler and more convenient for analysis inverse (not direct) approach, based on introducing another arbitrary auxiliary function instead of the function g(r) and directly specification of exact solutions in implicit form. Namely, we will consider the function f(r) to

be arbitrary, and the solution r = r(x) of equation (18), taking into account the relation h = rf, will be specified using an arbitrary function h = h(x) in implicit form

$$rf(r) = h(x). \tag{30}$$

The function g = g(r) in this approach is found from equation (18), that leads to the formula

$$g = C_1 + r^{-1} [C_2^2 h^{-3}(x) - h''_{xx}(x)].$$
(31)

The potential function g = g(r) for a given specific function h(x) is determined by eliminating x from relations (30)–(31).

Let us demonstrate with several specific examples how the described approach works in practice. To do this, we will take simple elementary functions h = h(x) and will find the potential functions g = g(r) generated by them. In these cases, the functions g(r) will be expressed through f(r).

Example 2. In relations (30)–(31) we substitute the function

$$h(x) = a(x+b)^k,$$
 (32)

where a, b, and k are arbitrary constants. As a result, we have

$$rf(r) = a(x+b)^{k},$$

$$g = C_{1} + r^{-1}[a^{-3}C_{2}^{2}(x+b)^{-3k} - ak(k-1)(x+b)^{k-2}].$$
(33)

By excluding x from these relations, we obtain the dependence of the potential on r:

$$g(r) = C_1 + C_2^2 r^{-4} f^{-3}(r) - a^{2/k} k(k-1) r^{-2/k} f^{(k-2)/k}(r).$$
(34)

In particular, by setting $C_2 = 0$ and k = -1 in (34), we arrive at the nonlinear Schrödinger equation

$$iu_t + [f(|u|)u]_{xx} + [C_1 - 2a^{-2}r^2f^3(|u|)]u = 0.$$
(35)

This equation, containing an arbitrary function f(|u|), admits an exact solution of the form

$$u = r(x)e^{i(C_1t + C_3)}, (36)$$

in which the amplitude r = r(x) is given by the implicit relation

$$rf(r) = a(x+b)^{-1}.$$

Example 3. In relations (30)–(31) we substitute $h = a \cos(kx+b)$ and $C_2 = 0$. Taking into account that $h''_{xx} = -k^2h$ and h = rf(r), we find the potential $g = C_1 + k^2 f(r)$. Thus, we arrive at the nonlinear Schrödinger equation

$$iu_t + [f(|u|)u]_{xx} + [C_1 + k^2 f(|u|)]u = 0,$$
(37)

which has an exact solution of the form (36), where the amplitude r = r(x) is given by the implicit expression

$$a\cos(kx+b) = rf(r).$$

Example 4. Setting $h = ae^{kx} + be^{-kx}$ and $C_2 = 0$ in (30)–(31), we obtain the Schrödinger equation

$$iu_t + [f(|u|)u]_{xx} + [C_1 - k^2 f(|u|)]u = 0,$$
(38)

which has an exact solution of the form (36), where the amplitude r = r(x) is given implicitly by the relation

$$ae^{kx} + be^{-kx} = rf(r).$$

2°. The case when two functions f and g are given by one arbitrary function. Equation (16) includes two time-dependent functions a = a(t) and b = b(t). Using the splitting principle [12, 38], we impose the following differential constraints on these functions:

$$a'_t = -A_1 a^2 + B_1, \quad b'_t = -A_2 a^2 + B_2,$$
(39)

where A_1 , B_1 , A_2 , and B_2 are arbitrary constants. As a result, we have

$$a^{2}\left(A_{1}\int h^{-2}dx + A_{2} - r^{-1}h^{-3}\right) - B_{1}\int h^{-2}dx - B_{2} + r^{-1}[h_{xx}'' + rg(r)] = 0.$$

By equating the functional factor at a^2 to zero, we obtain two integro-differential equations

$$A_1 \int h^{-2} dx + A_2 - r^{-1} h^{-3} = 0, \quad h = rf(r), \tag{40}$$

$$-B_1 \int h^{-2} dx - B_2 + r^{-1} [h_{xx}'' + rg(r)] = 0.$$
(41)

Depending on the value of the free coefficient A_1 , two different situations may arise. Let us consider them in order.

2.1. For $A_1 \neq 0$, eliminating the integral term from (40)–(41), we arrive at the ODE:

$$h_{xx}'' - \frac{B_1}{A_1}h^{-3} + \left(B_1\frac{A_2}{A_1} - B_2\right)r + rg(r) = 0, \quad h = rf(r), \tag{42}$$

which, up to renaming of constants, coincides with equation (18). Instead of equation (41), we can consider equation (42). Note that equation (40) reduces to an ODE by differentiating with respect to x.

The system of two equations (40)–(41) includes three functions r = r(x), f = f(r), and g = g(r). If we set one of these functions arbitrarily, then the other two can be found from equations (40)–(41) or equations (40) and (42).

The solution of system (40)–(41) can be represented in parametric form by arbitrarily setting the function h = h(x) (which is the product of the functions r and f). Indeed, substituting this function into equations (40) and (42), we obtain

$$r = h^{-3}(x) \left[A_1 \int \frac{dx}{h^2(x)} + A_2 \right]^{-1},$$

$$f = h^4(x) \left[A_1 \int \frac{dx}{h^2(x)} + A_2 \right],$$

$$g = B_2 - B_1 \frac{A_2}{A_1} + \left[\frac{B_1}{A_1} - h^3(x) h_{xx}''(x) \right] \left[A_1 \int \frac{dx}{h^2(x)} + A_2 \right],$$
(43)

where h = h(x) is an arbitrary function, and x plays the role of a parameter here.

For $A_1 \neq 0$, the functions a = a(t) and b = b(t), included in the formula for phase (15), are determined by integrating the first-order ODEs (39) and are expressed through elementary functions:

$$a(t) = \begin{cases} \frac{B_1}{\sqrt{A_1 B_1}} \tanh\left(\sqrt{A_1 B_1} t + C_1\right) & \text{if } A_1 B_1 > 0, \\ \frac{B_1}{\sqrt{|A_1 B_1|}} \operatorname{tg}\left(\sqrt{|A_1 B_1|} t + C_1\right) & \text{if } A_1 B_1 < 0, \\ (A_1 t + C_1)^{-1} & \text{if } B_1 = 0; \end{cases}$$

$$b(t) = \begin{cases} \left(B_2 - A_2 \frac{B_1}{A_1}\right) t + \frac{A_2 B_1}{A_1 \sqrt{A_1 B_1}} \tanh\left(\sqrt{A_1 B_1} t + C_1\right) + C_2 & \text{if } A_1 B_1 > 0, \\ \left(B_2 - A_2 \frac{B_1}{A_1}\right) t + \frac{A_2 B_1}{A_1 \sqrt{|A_1 B_1|}} \operatorname{tg}\left(\sqrt{|A_1 B_1|} t + C_1\right) + C_2 & \text{if } A_1 B_1 < 0, \\ \left(\frac{A_2}{A_1 (A_1 t + C_1)} + B_2 t + C_2 & \text{if } B_1 = 0, \\ (45) \end{cases}$$

where C_1 and C_2 are arbitrary constants.

Example 5. Substitute

$$A_2 = 0, \quad h = k(x + C_3)^{-1/2},$$

into (43). After elementary calculations we get

$$r = \frac{2}{A_1 k} (x + C_3)^{-1/2}, \quad f = \frac{1}{2} A_1 k^2 = \text{const},$$
$$g = B_2 + \frac{1}{2} B_1 k^{-2} (x + C_3)^2 - \frac{3}{8} A_1 k^2 (x + C_3)^{-2}.$$

Eliminating x from here, and for simplicity setting $k = \sqrt{2/A_1}$, we arrive at Eq. (2) with

$$g(r) = B_2 + \frac{B_1}{A_1}r^{-4} - \frac{3}{16}A_1^2r^4, \quad r = |u|.$$
(46)

Equation (2) with potential (46) has the exact solution

$$u = re^{i\varphi}, \quad r = \sqrt{\frac{2}{A_1(x+C_3)}}, \quad \varphi = \frac{1}{4}A_1a(t)(x+C_3)^2 + b(t),$$

where the functions a = a(t) and b(t) are determined by formulas (44) and (45).

2.2. When $A_1 = 0$, it follows from equation (40) that $f(r) = A_2^{-1/3} r^{-4/3}$. Equation (41) can be reduced by differentiation to a third-order nonlinear ODE, which cannot be integrated in closed form. Therefore, to construct some exact solutions of the nonlinear Schrödinger equation (3) we use here the inverse approach (without integrating ODEs) described earlier. For this we first set an arbitrary function h = h(x). Substituting this function into equations (40)–(41) with $A_1 = 0$, we obtain

$$r = A_2^{-1} h^{-3}(x),$$

$$f = A_2 h^4(x),$$

$$g = B_1 \int h^{-2}(x) \, dx + B_2 - A_2 h^3(x) h''_{xx}(x).$$
(47)

These formulas define the solution of system (40)–(41) with $A_1 = 0$, which is given in parametric form using one arbitrary function h = h(x). The dependence of the potential g = g(r) is obtained by eliminating x from the first and third relations (47).

For $A_1 = 0$, the functions a = a(t) and b = b(t), included in the formula for phase (15), are determined by integrating the first-order ODEs (39) and are expressed through elementary functions:

$$a(t) = B_1 t + C_1, \quad b(t) = -\frac{A_2}{3B_1}(B_1 t + C_1)^3 + B_2 t + C_2,$$

where C_1 and C_2 are arbitrary constants.

3.3 Generalized separable solutions with amplitude depending on the time t

To construct an exact solution, we now use the additional relation (12). Given that |u| = r, we substitute r = r(t) into the PDE system (8). By integrating the second equation of the resulting system twice, we arrive at a solution with generalized separation of variables of the following form:

$$r = r(t), \quad \varphi = a(t)x^2 + b(t)x + c(t),$$
(48)

where the function a = a(t) can be expressed in terms of r and f using the formula $a = -r'_t/(2rf)$. Substituting (48) into the first equation (8), we arrive at a quadratic equation with respect to x, the coefficients of which depend on time. As a result, the first equation of the system is reduced to a quadratic equation with

respect to x, the coefficients of which depend on time. By equating the functional coefficients of this quadratic equation to zero and adding the second equation of the system, which in this case depends only on t, we obtain the following system consisting of four ODEs:

$$a'_{t} = -4a^{2}f(r),$$

$$b'_{t} = -4abf(r),$$

$$c'_{t} = -b^{2}f(r) + g(r),$$

$$r'_{t} = -2arf(r).$$

(49)

Here the three first equations were divided by r.

From the first, second and fourth equations of system (49) we can obtain two integrals

$$a = C_1 r^2, \quad b = C_2 r^2,$$
 (50)

where C_1 and C_2 are arbitrary constants. Substitute the first expression (50) into the first equation (49). Integrating the resulting ODE, we find the dependence r = r(t) in implicit form

$$\int \frac{dr}{r^3 f(r)} = -2C_1 t - C_3, \tag{51}$$

where C_3 is an arbitrary constant. Integrating the third equation of system (49), we obtain

$$c = -C_2^2 \int r^4 f(r) \, dt + \int g(r) \, dt + C_4.$$
(52)

For simplicity, assuming in (50) and (52) $C_2 = 0$ and substituting the obtained expressions into (48), we find the functions that determine solution (6): (6):

$$r = r(t), \quad \varphi = C_1 r^2 (x+A)^2 + \int g(r) \, dt + C_4,$$
 (53)

where the function r = r(t) is given implicitly by expression (51). Note that an arbitrary constant A has been added to the second formula, since the system (8) is invariant with respect to an arbitrary constant shift along the spatial variable.

Example 6. For equation (3) with the power dispersion $f(r) = ar^k$ in solution (53) we must set

$$r(t) = \left[a(k+2)(2C_1t + C_3)\right]^{-\frac{1}{k+2}}$$

This formula is derived from (51).

4 Solutions that are nonlinear superpositions of traveling waves

The system of PDEs (8) admits an exact solution of the form

$$r = r(z), \quad \varphi = C_1 t + C_2 x + \theta(z), \quad z = x - \lambda t, \tag{54}$$

where C_1 , C_2 , and λ are arbitrary constants. The special case $C_1 = C_2 = 0$ in (54) corresponds to a traveling wave solution.

Substituting (54) into (8), we obtain the nonlinear system of two ODEs:

$$-r(C_1 - \lambda \theta'_z) + h''_{zz} - h(C_2 + \theta'_z)^2 + rg(r) = 0, -\lambda r'_z + 2h'_z(C_2 + \theta'_z) + h\theta''_{zz} = 0, \quad h = rf(r).$$
(55)

Note that the substitution $\xi = \theta'_z$ allows us to reduce the order of the ODE system (55) by one.

It is easy to verify that the second equation (55) admits the first integral

$$-\lambda \int h(r) \, dr + C_2 h^2 + h^2 \theta'_z = C_3, \quad h = r f(r), \tag{56}$$

where C_3 is an arbitrary constant. Eliminating further the derivative θ'_z from the first equation (55) using (56), we arrive at a second-order nonlinear autonomous ODE (that does not explicitly depend on z) of the form

$$h_{zz}^{\prime\prime} = \Phi(h) + \Psi(r), \qquad (57)$$

where the following notations are used:

$$\Phi(h) = C_3^2 h^{-3}, \quad h = rf(r),$$

$$\Psi(r) = (C_1 + \lambda C_2)r - C_3\lambda rh^{-2} - \lambda^2 rh^{-2}H + 2C_3\lambda h^{-3}H + \lambda^2 h^{-3}H^2 - rg(r), \quad H = \int h \, dr.$$
(58)

The appendix shows that the general solution of equation (57)-(58) can be expressed in quadratures and presented in an implicit form.

By resolving relation (56) with respect to θ'_z , and then integrating, we can find the function $\theta = \theta(z)$,

$$\theta = C_3 \int h^{-2} dz + \lambda \int h^{-2} H \, dz - C_2 z + C_4,$$

where C_4 is an arbitrary constant.

Remark 4. For the physical interpretation of solution (54), it is convenient to represent the phase φ in an equivalent form in two ways:

$$\varphi = C_2(x - \lambda_2 t) + \theta(x - \lambda_1 t), \qquad \lambda_1 = \lambda, \quad \lambda_2 = -C_1/C_2;$$
$$\varphi = (C_2 + C_1 \lambda^{-1})x + \theta_1(x - \lambda t), \qquad \theta_1(z) = \theta(z) - C_1 \lambda^{-1} z.$$

In the first case solution (54) can be interpreted as a nonlinear superposition of two traveling waves with velocities λ_1 and λ_2 , and in the second case as a nonlinear superposition of a stationary wave and a traveling wave with velocity λ .

Remark 5. Solution (54) can be obtained, for example, using the following reasoning. First of all one can note, that one of the two arbitrary constants C_1 or C_2 in (54) can be taken equal to zero without loss of generality (this fact corresponds to the redefinition of the function θ). Next, we pass in equation (3) or system (8) from the variables x, t to new independent variables z, t, where $z = x - \lambda t$. Then, by analogy with (11), we introduce the additional relation |u| = p(z)). After this, the procedure of separation of variables is used.

We will also describe another simpler way to obtain solution (54). First, in Eq. (3) we make the substitution $u = e^{i(C_1t+C_2x)}w$, where C_1 and C_2 are arbitrary real constants. Then a traveling wave solution of the transformed equation is sought, i.e. it is assumed that w = W(z), where $z = x - \lambda t$.

5 Brief conclusion

For the first time, the general nonlinear Schrödinger equation has been investigated, the dispersion and potential of which are specified by two arbitrary functions. New several closed-form solutions of this PDE have been found, which are expressed in quadratures or elementary functions. One-dimensional reductions have been described, which leads the considered nonlinear PDE to simpler ODEs. The obtained solutions can be used as test problems for numerical methods of integrating nonlinear PDEs of mathematical physics. It is important to note that these exact solutions are valid for two arbitrary functions f(z) and g(z) included in the general nonlinear Schrödinger equation (3), so they can be used for a wide variety of problems, specifying a specific form of these functions.

Data availability

No data was used for the research described in the article.

Declaration of competing interest

The authors declare that there is no conflict of interest.

Acknowledgements

The study was supported by the Ministry of Education and Science of the Russian Federation within the framework of the state assignments No. 124012500440-9 and No. FSWU-2023-0031.

Appendix. Solutions of special second-order ODEs

Let us consider a class of second-order autonomous ODEs of the special form

$$h''_{xx} = \Phi(h) + \Psi(r), \qquad h = \Theta(r), \tag{59}$$

where $\Phi(h)$, $\Psi(r)$, and $\Theta(r)$ are given functions. ODE (18) is a particular case of equation (59) with

$$\Phi(h) = C_2^2 h^{-3}, \quad \Psi(r) = C_1 r - rg(r), \quad \Theta(r) = rf(r), \tag{60}$$

For $\Psi(r) \equiv 0$, the general solution of the ODE (59) can be represented in implicit form [83]:

$$\int \left[2 \int \Phi(h) \, dh + A_1 \right]^{-1/2} dh = A_2 \pm x, \quad h = \Theta(r), \tag{61}$$

where A_1 and A_2 are arbitrary constants. In the general case, in the second term of the right-hand side of equation (59) we pass from the variable r to the variable h, setting $\Psi_1(h) = \Psi(r)$, where $h = \Theta(r)$. We have

$$h''_{xx} = \Phi(h) + \Psi_1(h).$$
(62)

The general solution of this equation is determined by the formula (61), in which $\Phi(h)$ must be replaced by $\Phi(h) + \Psi_1(h)$, or

$$\int \left[2 \int \Phi(h) \, dh + 2 \int \Psi_1(h) \, dh + A_1 \right]^{-1/2} dh = A_2 \pm x, \quad h = \Theta(r), \quad (63)$$

We do not change the first inner integral in (63), and in the second inner integral we return from the variable h to the variable r. Taking into account the relations $\Psi_1(h) = \Psi(r)$ and $dh = \Theta'_r(r) dr$, we finally obtain

$$\int \left[2 \int \Phi(h) \, dh + 2 \int \Psi(r) \Theta'_r(r) \, dr + A_1 \right]^{-1/2} dh = A_2 \pm x, \quad h = \Theta(r).$$
(64)

Substituting expressions (60) into (64), we find the general solution of Eq. (18):

$$\int \left[-C_2^2 h^{-2} + 2 \int [C_1 r - rg(r)] dh + A_1 \right]^{-1/2} dh = A_2 \pm x,$$

$$h = rf(r), \quad dh = [f(r) + rf'_r(r)] dr.$$

References

- 1. Agrawal G.P. Nonlinear Fiber Optics, 4th ed. New York: Academic Press, 2007.
- Kivshar Yu.S., Agrawal G.P. Optical Solitons: From Fibers to Photonic Crystals. San Diego: Academic Press, 2003.
- Kodama Y., Hasegawa A. Nonlinear pulse propagation in a monomode dielectric guide. IEEE J. Quantum Electronics 1987; 23(5): 510–524.
- Drazin P.G., Johnson R.S. Solitons: An Introduction. Cambridge: Cambridge Univ. Press, 1989.
- 5. Ablowitz M.J., Clarkson P.A. Solitons, Nonlinear Evolution Equations and Inverse Scattering. Cambridge: Cambridge Univ. Press, 1991.
- Akhmanov S.A., Sukhorukov A.P., Khokhlov R.V. Self-focusing and diffraction of light in a nonliner medium, Sov. Phys. Uspekhi 1968; 10(5): 609–636.

- Hasegawa A., Tappert F. Transmission of stationary nonlinear optical pulses in dispersive dielectric fibers. I. Anomalous dispersion. Appl. Phys. Lett. 1973; 23(3): 142–144.
- Hasegawa A., Tappert F. Transmission of stationary nonlinear optical pulses in dispersive dielectric fibers. II. Normal dispersion. Appl. Phys. Lett. 1973; 23(4): 171–172.
- Weiss J., Tabor M., Carnevale G. The Painlevé property for partial differential equations. J. Math. Phys. 24 (3) (1982) 522–526.
- Kudryashov N.A. Painlevé analysis of the resonant third-order nonlinear Schrödinger equation. Appl. Math. Lett. 2924; 158: 109232.
- Kudryashov N.A. Painlevé analysis of the Sasa–Satsuma equation. Phys. Lett. A 2024; 56: 129900.
- Polyanin A.D., Zaitsev V.F. Handbook of Nonlinear Partial Differential Equations, 2nd ed. Boca Raton: CRC Press, 2012.
- Al Khawaja U., Al Sakkaf L. Handbook of Exact Solutions to the Nonlinear Schrödinger Equations. Bristol: Inst. Phys. Publ., 2019.
- Polyanin A.D. Handbook of Exact Solutions to Mathematical Equations. Boca Raton: CRC Press-Chapman & Hall, 2025.
- 15. Ekici M. Kinky breathers, W-shaped and multi-peak soliton interactions for Kudryashov's quintuple power-law coupled with dual form of non-local refractive index structure. Chaos, Solitons & Fractals 2022; 159: 112172.
- Ekici M., Sonmezoglu A., Biswas A. Stationary optical solitons with Kudryashov's laws of refractive index. Chaos, Solitons & Fractals 2021; 151: 111226.
- Kumar V., Biswas A., Ekici M., Moraru L., Alzahrani A.K., Belic M.R., Timeb⁵ dependent coupled complex short pulse equation: Invariant analysis and complexitons. Chaos, Solitons & Fractals 2021; 150: 111151.

- 18. Sucu N., Ekici M., Biswas A. Stationary optical solitons with nonlinear chromatic dispersion and generalized temporal evolution by extended trial function approach. Chaos, Solitons & Fractals 2021; 147: 110971.
- Zhang J.-H., Qin H.-Q., Si Z.-Z., Jia Y.-H., Kudryashov N.A., Wang Y.-Y., Dai C.-Q. Pure-quartic soliton attracted state and multi-soliton molecules in mode-locked fiber lasers. Chaos, Solitons & Fractals 2024; 187: 115380.
- 20. Jiang J.-H., Si Z.-Z., Kudryashov N.A., Dai C.-Q., Liu W. Prediction of symmetric and asymmetric solitons and model parameters for nonlinear SchrΓ¶dinger equations with competing nonlinearities. Chaos, Solitons & Fractals 2024; 186: 115284.
- Zayed E.M.E., Shohib R.M.A., Alngar M.E.M., Biswas A., Moraru L., Khan S., Yildirim Y., Alshehri H.M., Belic M.R. Dispersive optical solitons with Schrödinger–Hirota model having multiplicative white noise via Ito Calculus. Phys. Lett. A: Gen., Atomic & Solid State Phys. 2022; 445: 128268.
- 22. Kudryashov N.A. Solitons of the complex modified Kortewegb⁵t⁴de Vries hierarchy. Chaos, Solitons & Fractals 2024; 184: 115010.
- Kudryashov N.A., Kutukov A.A., Biswas A., Zhou Q., Yildirim Y., Alshomrani A.S. Optical solitons for the concatenation model: Power-law nonlinearity. Chaos, Solitons & Fractals 2023; 177: 114212.
- Kudryashov N.A. Stationary solitons of the generalized nonlinear Schrödinger equation with nonlinear dispersion and arbitrary refractive index. Appl. Math. Lett. 2022; 128: 107888.
- 25. Malomed B.A. Soliton models: Traditional and novel, one- and multidimensional. Low Temp. Phys. 2022; 48(11): 856–895.
- 26. Zhou Q., Xu M., Sun Y., Zhong Y., Mirzazadeh M. Generation and transformation of dark solitons, anti-dark solitons and dark double-hump solitons. Nonlinear Dynamics 2022; 110(2): 1747–1752.

- 27. Wang M., Zhang J., Li L. The decay mode solutions of the cylindrical/spherical nonlinear Schrödinger equation. Appl. Math. Lett. 2023; 145: 108744.
- 28. Wang H., Li X., Zhou Q., Liu W. Dynamics and spectral analysis of optical rogue waves for a coupled nonlinear Schrödinger equation applicable to pulse propagation in isotropic media. Chaos, Solitons & Fractals 2023; 166: 112924.
- Li P., Dong L., Mihalache D., Malomed B.A. Symmetry-breaking bifurcations of pure-quartic solitons in dual-core couplers. Optics Lett. 2024; 49(20): 5858– 5861.
- 30. Zhong M., Chen Y., Yan Z., Malomed B.A. Suppression of soliton collapses, modulational instability and rogue-wave excitation in two-LΓ©vy-index fractional Kerr media. Proc. Royal Soc. A: Math., Phys. & Eng. Sci. 2024; 480: 2282.
- Kudryashov N.A., Biswas A., Borodina A.G., Yildirim Y., Alshehri H.M. Painlevé analysis and optical solitons for a concatenated model. Optik 2023; 272: 170255.
- 32. Arnous A.H., Biswas A., Yakup Y., Luminita M., Catalina I., Lucian G.P., Asim A. Optical solitons and complexitons for the concatenation model in birefringent fibers. Ukrainian J. Physical Optics 2023; 24(4): 04060–04086.
- 33. Arnous A.H., Biswas A., Kara A.H., Yildirim Y., Moraru L., Iticescu C., Moldovanu S., Alghamdi A.A. Optical solitons and conservation laws for the concatenation model with spatio-temporal dispersion (internet traffic regulation). J. Eur. Optical Soc. 2023; 19: 2.
- 34. Ekici M. Stationary optical solitons with complex Ginzburg–Landau equation having nonlinear chromatic dispersion and Kudryashov's refractive index structures. Phys. Lett. A: Gen. Atomic & Solid State Phys. 2022; 440: 128146.
- 35. Zayed E.M.E., El-Shater M., Arnous A.H., Yildirim Y., Hussein L., Jawad A.J.M., Veni S.S., Biswas A. Quiescent optical solitons with KudryashovB[™]s generalized quintuple-power law and nonlocal nonlinearity having nonlinear

chromatic dispersion with generalized temporal evolution by enhanced direct algebraic method and sub-ODE approach. Eur. Phys. J. Plus 2024; 139(10): 885.

- 36. Han T., Li Z., Li C., Zhao L. Bifurcations, stationary optical solitons and exact solutions for complex Ginzburg–Landau equation with nonlinear chromatic dispersion in non-Kerr law media. J. Optics 2023; 52(2): 831–844.
- 37. Bullough R.K. Solitons. Phys. Bull. 1978; 29(2): 78–82.
- Polyanin A.D., Zhurov A.I. Separation of Variables and Exact Solutions to Nonlinear PDEs. Boca Raton–London: CRC Press, 2022.
- Ibragimov N.H. (ed.). CRC Handbook of Lie Group Analysis of Differential Equations, Vol. 1. Symmetries, Exact Solutions and Conservation Laws. Boca Raton: CRC Press, 1994.
- 40. Broadbridge P., Bradshaw-Hajek B.H., Hutchinson A.J. Conditionally integrable PDEs, non-classical symmetries and applications. Proc. R. Soc. A 2023; 479: 20230209.
- Ovsiannikov L.V. Group Analysis of Differential Equations. Boston: Academic Press, 1982.
- 42. Bluman G.W., Kumei S. Symmetries and Differential Equations. New York: Springer, 1989.
- Olver P.J. Application of Lie Groups to Differential Equations, 2nd ed. New York: Springer, 2000.
- 44. Vaneeva O.O., Popovych R.O., Sophocleous C. Extended group analysis of variable coefficient reaction-diffusion equations with exponential nonlinearities.
 J. Math. Anal. Appl. 2012; 396: 225–242.
- 45. Polyanin A.D., Aksenov A.V. Unsteady magnetohydrodynamics PDE of Mongeb^T Ampére type: Symmetries, closed-form solutions, and reductions. Mathematics 2024; 12(13); 2127.

- 46. Clarkson P.A., Kruskal M.D. New similarity reductions of the Boussinesq equation. J. Math. Phys. 1989; 30: 2201–2213.
- 47. Nucci M.C., Clarkson P.A. The nonclassical method is more general than the direct method for symmetry reductions. An example of the Fitzhugh–Nagumo equation. Phys. Lett. A 1992; 164; 49–56.
- Olver P.J. Direct reduction and differential constraints. Proc. R. Soc. Lond. Ser. A 1994; 444: 509–523.
- 49. Clarkson P.A., Ludlow D.K., Priestley T.J. The classical, direct and nonclassical methods for symmetry reductions of nonlinear partial differential equations. Meth. Appl. Anal. 1997; 4: 173–195.
- Bluman G.W., Cole J.D. The general similarity solution of the heat equation.
 J. Math. Mech. 1969; 18: 1025–1042.
- Arrigo D., Broadbridge P., Hill J.M. Nonclassical symmetry solutions and the methods of Bluman–Cole and Clarkson–Kruskal. J. Math. Phys. 1993; 34: 4692–4703.
- 52. Pucci E., Saccomandi G. Evolution equations, invariant surface conditions and functional separation of variables. Phys. D 2000; 139: 28–47.
- 53. Bradshaw-Hajek D.H. Nonclassical symmetry solutions for non-autonomous reaction-diffusion equations. Symmetry 2019: 11: 208.
- 54. Cherniha R.; Davydovych V. Nonlinear Reaction-Diffusion Systems: Conditional Symmetry, Exact Solutions and Their Applications in Biology. Cham: Springer, 2017.
- 55. Galaktionov V.A., Svirshchevskii S.R. Exact Solutions and Invariant Subspaces of Nonlinear Partial Differential Equations in Mechanics and Physics. Boca Raton: Chapman & Hall/CRC Press, 2007.

- 56. Galaktionov V.A. Invariant subspaces and new explicit solutions to evolution equations with quadratic nonlinearities. Proc. Royal Soc. Edinburgh Sect. A 1995; 125: 225–246.
- 57. Polyanin A.D., Zhurov A. I. Functional and generalized separable solutions to unsteady Navier–Stokes equations. Int. J. Non-Linear Mech. 2016; 79: 88–98.
- 58. Kosov A.A., Semenov E.I. Exact solutions of the generalized Richards equation with power-law nonlinearities. Diff. Equ. 2020; 56: 1119–1129.
- Miller W., Rubel L.A. Functional separation of variables for Laplace equations in two dimensions. J. Phys. A 1993; 26: 1901–1913.
- Doyle P.W., Vassiliou P.J. Separation of variables for the 1-dimensional nonlinear diffusion equation. Int. J. Non-Linear Mech. 1998; 33: 315–326.
- 61. Estevez P.G., Qu C., Zhang S. Separation of variables of a generalized porous medium equation with nonlinear source. J. Math. Anal. Appl. 2002; 275: 44–59.
- 62. Polyanin A.D. Functional separation of variables in nonlinear PDEs: General approach, new solutions of diffusion-type equations. Mathematics 2020; 8(1): 90.
- Sidorov A.F., Shapeev V.P., Yanenko N.N. Method of Differential Constraints and its Applications in Gas Dynamics. Novosibirsk: Nauka, 1984.
- Kaptsov O.V., Verevkin L.V. Differential constraints and exact solutions of nonlinear diffusion equations. J. Phys. A Math. Gen. 2003; 36; 1401–1414.
- Meleshko S.V. Methods for Constructing Exact Solutions of Partial Differential Equations. New York: Springer, 2005.
- Kruglikov B. Symmetry approaches for reductions of PDEs, differential constraints and Lagrange–Charpit method. Acta Appl. Math. 2008; 101: 145– 161.

- Ablowitz M.J., Kaup D.J., Newell A.C., Segur H. The inverse scattering transform–Fourier analysis for nonlinear problems. Stud. Appl. Math. 1974; 53: 249–257.
- Calogero F., Degasperis A. Spectral Transform and Solitons: Tolls to Solve and Investigate Nonlinear Evolution Equations. Amsterdam: North-Holland Publ., 1982.
- Kuo S., Nonlinear Waves and Inverse Scattering Transform. World Scientific Publ., 2023.
- 70. Vu P.L. Inverse Scattering Problems and Their Application to Nonlinear Integrable Equations, 2nd ed. Boca Raton: Chapman & Hall/CRC Press, 2023.
- Kudryashov N.A. Methods of Nonlinear Mathematical Physics. Izd. Dom Intellekt: Dolgoprudnyi, 2010.
- 72. Conte R., Musette M. The Painlevé Handbook, 2nd ed.; Cham: Springer, 2020.
- Jimbo J., Kruskal M.D., Miwa T. Painlevé test for the self-dual Yang–Mills equation. Phys. Lett. A 1982; 92(2): 59–60.
- Weiss J., Tabor M., Carnevale G. The Painlevé property for partial differential equations. J. Math. Phys., 1983; 24: 522–526.
- Weiss J. The Painlevé property for partial differential equations. II: BΓ¤cklund transformation, Lax pairs, and the Schwarzian derivative. J. Math. Phys. 1983; 24: 1405–1413.
- 76. Conte R. Invariant Painlevé analysis of partial differential equations. Phys. Lett. A 1989; 140(7–8); 383–390.
- 77. Kudryashov N.A. Simplest equation method to look for exact solutions of nonlinear differential equations. Chaos, Solitons & Fractals 2005; 24: 1217– 1231.
- Aksenov A.V., Polyanin A.D. Methods for constructing complex solutions of nonlinear PDEs using simpler solutions. Mathematics 2021; 9: 345.

- 79. Polyanin A.D., Sorokin V.G., Zhurov A.I. Delay Ordinary and Partial Differential Equations. Boca Raton–London: CRC Press, 2024.
- 80. Polyanin A.D., Zhurov A.I. Functional constraints method for constructing exact solutions to delay reaction-diffusion equations and more complex nonlinear equations. Commun. Nonlinear Sci. Numer. Simul. 2014; 19(3): 417– 430.
- Polyanin A.D., Zhurov A.I. The functional constraints method: Application to non-linear delay reaction-diffusion equations with varying transfer coefficients. Int. J. Non-Linear Mech. 2014; 67: 267–277.
- Polyanin A.D., Sorokin V.G. Reductions and exact solutions of Lotka–Volterra and more complex reaction-diffusion systems with delays. Appl. Math. Lett. 2022; 125: 107731.
- Polyanin A.D., Zaitsev V.F. Handbook of Ordinary Differential Equations: Exact Solutions, Methods, and Problems. Boca Raton–London: CRC Press, 2018.