

Generalizing the Bierbrauer–Friedman bound for orthogonal arrays*

Denis S. Krotov[‡] Ferruh Özbudak[§] Vladimir N. Potapov[¶]

Abstract

We characterize mixed-level orthogonal arrays in terms of algebraic designs in a special multigraph. We prove a mixed-level analog of the Bierbrauer–Friedman (BF) bound for pure-level orthogonal arrays. For the case when the numbers of levels are powers of the same prime number, we characterize, in terms of multispreads, additive mixed-level orthogonal arrays attaining the BF bound. For pure-level orthogonal arrays, we consider versions of the BF bound obtained by replacing the Hamming graph by its polynomial generalization and show that in some cases this gives a new bound.

Keywords: orthogonal array, algebraic t -design, mixed orthogonal array, completely-regular code, equitable partition, intriguing set, Hamming graph, Bierbrauer–Friedman bound, additive codes.

1 Introduction

The Bierbrauer–Friedman bound for pure-level orthogonal arrays with parameters $OA(N, n, q, t)$ says that

$$N \geq q^n \left(1 - \left(1 - \frac{1}{q} \right) \frac{n}{t+1} \right), \quad (1)$$

see [8] for the case $q = 2$ and [2] for general q . It is easy to see that the bound is nonnegative if and only if $t + 1 > \frac{q-1}{q}n$; so, it is effective for high values of t (in contrast, Rao’s bound [14] is effective for relatively small t). The bound is tight, and there are orthogonal arrays constructed as linear [10, Section 4.3] or additive [1, Section 4.2] codes that attain it. Binary ($q = 2$) orthogonal

*The work of D.K. was supported by the Natural Science Foundation of Hebei Province (project No. A2023205045) and within the framework of the state contract of the Sobolev Institute of Mathematics (FWNF2022-0017).

[‡]School of Mathematical Sciences, Hebei Key Laboratory of Computational Mathematics and Applications, Hebei Normal University, Shijiazhuang 050024, P. R. China

[‡]Sobolev Institute of Mathematics, Novosibirsk 630090, Russia

[§]Faculty of Engineering and Natural Science, Sabancı University, Istanbul, Turkey

[¶]Faculty of Engineering and Natural Science, Sabancı University, Istanbul, Turkey

arrays attaining bound (1) whose size is not a power of 2 can be constructed as completely regular codes (see Definition 2 below) by the Fon-Der-Flaass construction [7]; the first example is $\text{OA}(1536, 13, 2, 7)$. Ternary ($q = 3$) arrays with the similar property were recently discovered, again in terms of CR codes, in [9]; the first example is $\text{OA}(5 \cdot 3^8, 11, 3, 8)$. For $q \geq 4$, the problem of existence of orthogonal arrays attaining bound (1) whose size is not a power of q or of its prime divisor remains open. Similar questions can be considered for mixed-level orthogonal arrays.

In this correspondence, we prove (Section 3) that (1) holds for mixed-level orthogonal arrays $\text{OA}(N, q_1 \cdot q_2 \cdot \dots \cdot q_n, t)$ if we replace q^n by the product of all q_i s and $\frac{1}{q}$ by the average value of $\frac{1}{q_i}$. The arrays (we treat an array as a multiset of rows of length n) attaining this bound are necessarily simple sets (without repeated elements), independent sets (without pairs of elements at Hamming distance 1), and intriguing sets (completely regular codes with covering radius 1, see the definition below). The new bound is tighter than the previously known generalization [5]

$$N \geq q_m^n \left(1 - \frac{(q-1)n}{q_M(t+1) - (q_M - q)n} \right) \quad (2)$$

if $q_M(t+1) - (q_M - q)n > 0$,

where q_m , q , and q_M are respectively the minimum, the average, and the maximum value of q_i , $i = 1, \dots, n$. For example, for $\text{OA}(N, 2^1 4^4, 3)$ the new bound is tight: $N \geq 64 = 2^9 \cdot \left(1 - \left(1 - \frac{3}{10} \right) \frac{5}{3+1} \right)$ (see the construction in Example 1), while (2) gives $N \geq 2^5 \cdot \left(1 - \frac{(3.6-1) \cdot 5}{4 \cdot (3+1) - (4-3.6) \cdot 5} \right) = \frac{16}{7}$.

Further (Section 4), in the case when all q_i are powers of the same prime p , we prove that additive (linear over $\text{GF}(p)$) mixed-level orthogonal arrays attaining the Bierbrauer–Friedman bound are equivalent to special partitions of a vector space into subspaces, called multispreads [12].

Finally (Section 5), we discuss variations of bound (1) that can give positive values in the cases when the original bound (1) is negative. As an example, for $\text{OA}(N, n, 2, \frac{n}{2} - 1)$, n even, we obtain the bound

$$N > 0.409 \cdot n^{-1} \cdot 2^n.$$

2 Definitions and notations

By a graph, we will mean a multigraph, with multiple edges and loops allowed. A graph without loops and edge multiplicities more than 1 is called simple. For a graph G and a positive integer scalar s , sG denotes the graph on the same vertex set with all edge multiplicities multiplied by s . For two graphs $G' = (V', E')$ and $G'' = (V'', E'')$, $G' \square G''$ denotes their Cartesian product, the graph with the vertex set $V' \times V''$ and the edge (multi)set $\{ \{(v_1, v), (v_2, v)\} : \{v_1, v_2\} \in E', v \in V'' \} \cup \{ \{(v, v_1), (v, v_2)\} : \{v_1, v_2\} \in E'', v \in V' \}$.

For a positive integer s , the set $\{0, \dots, s-1\}$ will be denoted by $[s]$.

Definition 1 (adjacency matrix of a graph, eigenfunctions and eigenspaces). For a graph $G = (V, E)$, the *adjacency matrix* A is the symmetric nonnegative integer $|V| \times |V|$ matrix whose rows and columns are indexed by V and the (x, y) th element $A_{x,y}$, $x, y \in V$, equals the multiplicity of $\{x, y\}$ in the edge multiset E . The eigenvectors of A , treated as functions from V to \mathbb{C} , are called *eigenfunctions* of G , or θ -*eigenfunctions*, where θ is the corresponding eigenvalue. An *eigenspace* of G is the subspace of the vector space \mathbb{C}^V consisting of the constantly zero function and all θ -eigenfunctions for some eigenvalue θ . Since the adjacency matrix of a graph is symmetric, the eigenspaces are pairwise orthogonal.

Definition 2 (completely regular code with covering radius 1, $\{b; c\}$ -CR code). A set C of vertices of a regular graph $G = (V, E)$ is called a *completely regular code* with covering radius 1 and intersection array $\{b; c\}$, where $b, c > 0$, or simply a $\{b; c\}$ -*CR code*, if for every vertex in V (in $V \setminus C$) the number of edges that connect it with $V \setminus C$ (respectively, with C), equals b (respectively, c).

Definition 3 ($H(q_1 \cdot q_2 \cdot \dots \cdot q_n)$, Hamming graph). For integers $q_1, q_2, \dots, q_n \geq 2$, the graph $H(q_1 \cdot q_2 \cdot \dots \cdot q_n)$ is the graph on the set of n -tuples from $V = [q_1] \times \dots \times [q_n]$ with two n -tuples forming an edge of multiplicity μ if and only if they differ in only the i th position for some $i \in \{1, \dots, n\}$ and $\mu = Q/q_i$, where $Q = \text{lcm}(q_1, \dots, q_n)$ (least common multiple). If $q_1 = q_2 = \dots = q_n = q$, then $H(q_1 \cdot q_2 \cdot \dots \cdot q_n)$ is a simple graph known as a *Hamming graph* and denoted $H(n, q)$.

Definition 4 (algebraic t -design). For a regular graph $G = (V, E)$ with eigenvalues $\theta_0 > \theta_1 > \dots > \theta_d$ and the corresponding eigenspaces S_0, S_1, \dots, S_d , a multiset C of its vertices is called an *algebraic t -design* (with respect to the natural descending ordering $\theta_0, \theta_1, \dots, \theta_d$ of the graph eigenvalues) if in the decomposition

$$f_C = \varphi_0 + \varphi_1 + \dots + \varphi_d, \quad \varphi_i \in S_i,$$

of the multiplicity function f_C (in the case of a simple set, the characteristic function, indicator) of C we have $\varphi_1 = \dots = \varphi_t \equiv 0$.

Definition 5 (orthogonal arrays, $\text{OA}(N, q_1 \cdot q_2 \cdot \dots \cdot q_n, t)$). A nonempty multiset C of n -tuples from $V = [q_1] \times \dots \times [q_n]$ is called an *orthogonal array* of strength t , $\text{OA}(|C|, q_1 \cdot q_2 \cdot \dots \cdot q_n, t)$, if for any distinct i_1, \dots, i_t from $\{1, \dots, n\}$ and any $a_1 \in q_{i_1}, \dots, a_t \in q_{i_t}$, the number of $(x_1, \dots, x_n) \in C$ such that $x_{i_j} = a_j$, $j = 1, \dots, t$, equals $\frac{|C|}{q_{i_1} q_{i_2} \dots q_{i_t}}$ (i.e., independent on the choice of a_j , $j = 1, \dots, t$). If $q_1 = q_2 = \dots = q_n$, then such orthogonal arrays are called *pure-level* (in some literature, *symmetric*) and also denoted $\text{OA}(N, n, q_1, t)$; otherwise, they are called *mixed-level*, or just *mixed* (in some literature, *asymmetric*). For brevity, in the notation $q_1 \cdot q_2 \cdot \dots \cdot q_n$, equal values of q_i can be grouped using degrees, e.g., $\text{OA}(N, 2 \cdot 2 \cdot 5, t)$ is the same as $\text{OA}(N, 2^2 5^1, t)$, but not the same as $\text{OA}(N, 4 \cdot 5, t) = \text{OA}(N, 4^1 5^1, t)$.

3 Orthogonal arrays and algebraic designs

We first describe the eigenspaces of the graph $H(q_1 \cdot q_2 \cdot \dots \cdot q_n)$.

Lemma 6. *The following functions form an orthogonal basis from eigenfunctions of $H(q_1 \cdot q_2 \cdot \dots \cdot q_n)$:*

$$\chi_{(b_1, b_2, \dots, b_n)}(x_1, x_2, \dots, x_n) = \xi_1^{b_1 x_1} \xi_2^{b_2 x_2} \dots \xi_n^{b_n x_n}, \quad b_i \in [q_i], \quad (3)$$

where ξ_i is the degree- q_i primitive root of 1. Moreover, $\chi_{(b_1, b_2, \dots, b_n)}$ is an eigenfunction corresponding to the eigenvalue $\theta_w = k - wQ$, where

$$k = \sum_{i=1}^n \frac{Q}{q_i} (q_i - 1)$$

is the degree of $H(q_1 \cdot q_2 \cdot \dots \cdot q_n)$, $Q = \text{lcm}(q_1, \dots, q_n)$, and w is the number of nonzeros among b_1, b_2, \dots, b_n .

Proof. The complete graph K_{q_i} has eigenfunctions $\xi_{(b_i)}(x_i) = \xi_i^{b_i x_i}$ corresponding to the eigenvalue $q_i - 1$ if $b_i = 0$ and -1 otherwise. After multiplying (the multiplicity of all edges of) K_{q_i} by $\frac{Q}{q_i}$, we get the same eigenfunctions with eigenvalues $\frac{Q}{q_i}(q_i - 1)$ and $-\frac{Q}{q_i}(q_i - 1)$. We note that the difference Q between these two eigenvalues does not depend on i .

The rest is straightforward from the following easy property of the Cartesian product of graphs, see e.g. [3, Section 1.4.6]: if $\chi'(x')$, $x' \in V'$, and $\chi''(x'')$, $x'' \in V''$, are θ' - and θ'' -eigenfunctions of graphs $\Gamma' = (V', E')$ and $\Gamma'' = (V'', E'')$, respectively, then $\chi(x', x'') = \chi'(x')\chi''(x'')$ is a $(\theta' + \theta'')$ -eigenfunction of $\Gamma' \square \Gamma''$. \square

The following fact generalizes Delsarte's characterization [4, Theorem 4.4] of pure-level orthogonal arrays as algebraic designs.

Theorem 7. *A multiset C of words from $V = [q_1] \times \dots \times [q_n]$ is an orthogonal array $OA(|C|, q_1 \cdot q_2 \cdot \dots \cdot q_n, t)$ if and only if C is an algebraic t -design in $H(q_1 \cdot q_2 \cdot \dots \cdot q_n)$.*

Proof. If. Assume C is an algebraic t -design and f_C is its multiplicity function. We need to show that the sum of f_C over $V_{i_1, \dots, i_t}^{a_1, \dots, a_t}$ does not depend on the choice of a_1, \dots, a_t , where $1 \leq i_1 < \dots < i_t \leq n$, $a_j \in [q_{i_j}]$, and $V_{i_1, \dots, i_t}^{a_1, \dots, a_t}$ denotes the set of all n -tuples (x_1, \dots, x_n) from V such that $x_{i_1} = a_1, \dots, x_{i_t} = a_t$.

We have

$$f_C = \varphi_0 + \varphi_{t+1} + \varphi_{t+2} + \dots + \varphi_n,$$

where φ_i is the zero constant or a θ_i -eigenfunction of $H(q_1 \cdot q_2 \cdot \dots \cdot q_n)$.

For $i > t$, each such eigenfunction is a linear combination of basis eigenfunctions $\chi_{(b_1, \dots, b_n)}$ from (3), where the number of nonzero elements among b_1, \dots, b_n is i . We claim that

(*) for more than t nonzeros in (b_1, \dots, b_n) , the sum of $\chi_{(b_1, \dots, b_n)}$ over $V_{i_1, \dots, i_t}^{a_1, \dots, a_t}$ equals 0. Indeed, denoting by l_1, \dots, l_s the indices from $\{1, \dots, n\} \setminus \{i_1, \dots, i_t\}$, $s = n - t$, we have

$$\begin{aligned} & \sum_{(x_1, \dots, x_n) \in V_{i_1, \dots, i_t}^{a_1, \dots, a_t}} \chi_{(b_1, \dots, b_n)}(x_1, \dots, x_n) \\ &= \xi_{i_1}^{b_{i_1} a_{i_1}} \cdot \dots \cdot \xi_{i_t}^{b_{i_t} a_{i_t}} \cdot \sum_{x_{l_1}=0}^{q_{l_1}} \xi_{l_1}^{b_{l_1} x_{l_1}} \cdot \dots \cdot \sum_{x_{l_s}=0}^{q_{l_s}} \xi_{l_s}^{b_{l_s} x_{l_s}}. \end{aligned} \quad (4)$$

Since the number of nonzeros is larger than t , for at least one j from $\{l_1, \dots, l_s\}$ we have $b_j \neq 0$. It follows that at least one sum in (4) equals 0, which proves (*).

From (*) we conclude that the sum of f_C over $V_{i_1, \dots, i_t}^{a_1, \dots, a_t}$ equals the sum of ϕ_0 over $V_{i_1, \dots, i_t}^{a_1, \dots, a_t}$. Since ϕ_0 is a constant, the sum does not depend on the choice of a_1, \dots, a_t .

Only if. We need to show that f_C is orthogonal to the θ_i -eigenspace for every i in $\{1, \dots, t\}$, i.e., to all $\chi_{(b_1, \dots, b_n)}$ with more than 0 and at most t nonzeros in (b_1, \dots, b_n) . Let all i such that $b_i \neq 0$ lie in $\{i_1, \dots, i_t\}$, where $1 \leq i_1 < \dots < i_t \leq n$. For the standard scalar product $(f_C, \chi_{(b_1, \dots, b_n)})$ of f_C and $\chi_{(b_1, \dots, b_n)}$, we have

$$\begin{aligned} (f_C, \chi_{(b_1, \dots, b_n)}) &= \sum_{(x_1, \dots, x_n) \in V} \chi_{(b_1, \dots, b_n)}(x_1, \dots, x_n) \cdot f_C(x_1, \dots, x_n) \\ &\stackrel{(i)}{=} \sum_{(x_1, \dots, x_n) \in V} \xi_{i_1}^{b_{i_1} x_{i_1}} \cdot \dots \cdot \xi_{i_t}^{b_{i_t} x_{i_t}} \cdot f_C(x_1, \dots, x_n) \\ &= \sum_{a_1=0}^{q_{i_1}-1} \dots \sum_{a_t=0}^{q_{i_t}-1} \sum_{(x_1, \dots, x_n) \in V_{i_1, \dots, i_t}^{a_1, \dots, a_t}} \xi_{i_1}^{b_{i_1} x_{i_1}} \cdot \dots \cdot \xi_{i_t}^{b_{i_t} x_{i_t}} \cdot f_C(x_1, \dots, x_n) \\ &= \sum_{a_1=0}^{q_{i_1}-1} \xi_{i_1}^{b_{i_1} a_1} \cdot \dots \cdot \sum_{a_t=0}^{q_{i_t}-1} \xi_{i_t}^{b_{i_t} a_t} \cdot \sum_{(x_1, \dots, x_n) \in V_{i_1, \dots, i_t}^{a_1, \dots, a_t}} f_C(x_1, \dots, x_n) \\ &\stackrel{(ii)}{=} \sum_{a_1=0}^{q_{i_1}-1} \xi_{i_1}^{b_{i_1} a_1} \cdot \dots \cdot \sum_{a_t=0}^{q_{i_t}-1} \xi_{i_t}^{b_{i_t} a_t} \cdot |C|/q_{i_1} \dots q_{i_t} \stackrel{(iii)}{=} 0, \end{aligned}$$

where in equality (i) we use that $\xi_j^{b_j x_j} = 1$ if $b_j = 0$, in (ii) we use the definition of an orthogonal array, and (iii) holds because for some j in $\{i_1, \dots, i_t\}$ we have $b_j \neq 0$ and hence $\sum_{a=0}^{q_j} \xi_j^{b_j a} = 0$. We have shown that f_C is orthogonal to $\chi_{(b_0, \dots, b_n)}$ for any (b_0, \dots, b_n) with more than 0 and less than $t + 1$ nonzeros. Hence, C is an algebraic t -design. \square

Lemma 8 ([13, Sect. 4.3.1]). *The cardinality of an algebraic t -design C in a k -regular graph $G = (V, E)$ with eigenvalues $k = \theta_0 > \theta_1 > \dots > \theta_d$ satisfies the inequality*

$$\frac{|C|}{|V|} \geq \frac{-\theta_{t+1}}{k - \theta_{t+1}}. \quad (5)$$

Moreover, a multiset C of vertices of G is an algebraic t -design meeting (5) with equality if and only if C is a simple set (without multiplicities more than 1) and a $\{k; -\theta_{t+1}\}$ -CR code.

The following bound was proved in [8] for $q_1 = \dots = q_s = 2$ and in [2] for $q_1 = \dots = q_s = q$ for any q . The theorem generalizes the mentioned results to the case of mixed-level orthogonal arrays.

Theorem 9. *For an orthogonal array C with parameters $OA(N, q_1 \dots q_n, t)$, we have*

$$N \geq q_1 q_2 \dots q_n \left(1 - \left(1 - \frac{1}{q}\right) \frac{n}{t+1}\right), \quad (6)$$

where q is the harmonic mean of all q_i , i.e., $\frac{1}{q} = \frac{1}{n} \sum_{i=1}^n \frac{1}{q_i}$.

Moreover, a multiset of vertices of the graph $H = H(q_1 \dots q_n)$ is an $OA(|C|, q_1 \dots q_n, t)$ meeting (6) with equality if and only if C is a simple set (without multiplicities more than 1) and a $\{k; -\theta_{t+1}\}$ -CR code, where $k = Qn(1 - \frac{1}{q})$ (the degree of H) and $\theta_{t+1} = k - (t+1)Q$, $Q = \text{lcm}(q_1, \dots, q_n)$.

Proof. Taking into account Theorem 7 and Lemma 8, it remains to check that (5) and (6) are the same for $G = H$. Indeed, the degree of H is

$$k = \frac{Q}{q_1}(q_1 - 1) + \dots + \frac{Q}{q_n}(q_n - 1) = Qn - Q\left(\frac{1}{q_1} + \dots + \frac{1}{q_n}\right) = Qn\left(1 - \frac{1}{q}\right)$$

and $\theta_i = k - iQ$. So, we find

$$\frac{-\theta_{t+1}}{k - \theta_{t+1}} = \frac{(t+1)Q - k}{(t+1)Q} = 1 - \left(1 - \frac{1}{q}\right) \frac{n}{t+1}.$$

□

Remark 1. As noted in [5], the definition of an orthogonal array implies that N must be a multiple of

$$S = \text{LCM}\{q_{i_1} q_{i_2} \dots q_{i_t} : 0 < i_1 < \dots < i_t\}.$$

In particular, this gives the bound $N \geq S$ (by similarity of arguments, it can be considered as an analog of the Singleton bound for error-correcting codes), but also means that any bound of form $N \geq B(q_1 \dots q_n, t)$ can be rounded to

$$N \geq \lceil B(q_1 \dots q_n, t)/S \rceil \cdot S. \quad (7)$$

However, in some cases bound (6) is already divisible by S , and the problem of existence orthogonal arrays attaining it arises. In the next section, we show how to construct such arrays in the case when all q_i are powers of the same prime number.

4 Additive mixed orthogonal arrays attaining the Bierbrauer–Friedman bound

In this section, all q_i are powers of the same prime p , and $[p^i]$ is associated with the vector space \mathbb{F}_p^i (to be explicit, an integer a in $[p^i]$ can be associated with its p -based notation, treated as a vector from \mathbb{F}_p^i). We should warn the reader about the following difference in notation: now q_i denotes p^i , not the number of levels (alphabet size) in the i th position. This is because in this section, it is convenient to write parameters in the form $\text{OA}(N, q_1^{n_1} q_2^{n_2} \dots q_s^{n_s}, t)$ (all q_i are different, but some n_i , $i < s$, can be zero), while in Section 3 the preferred form was $\text{OA}(N, q_1 \cdot q_2 \cdot \dots \cdot q_n, t)$, where q_i are not necessarily distinct.

Definition 10. A set of $(n_1 + \dots + n_s)$ -tuples from $V = (\mathbb{F}_p^1)^{n_1} \times (\mathbb{F}_p^2)^{n_2} \times \dots \times (\mathbb{F}_p^s)^{n_s}$ is called *additive* (an additive code, or an additive orthogonal array if we consider it as an orthogonal array) if it is closed with respect to the coordinatewise addition, i.e., forms a subspace of the $(n_1 + 2n_2 + \dots + sn_s)$ -dimensional vectorspace V over \mathbb{F}_p .

Any k -dimensional vector subspace C of an n -dimensional vector space S_n can be represented as the null-space (kernel) of a homomorphism from S to an $(n - k)$ -dimensional vector space S_{n-k} over the same field. If the bases in S_n and S_{n-k} , are fixed, such a homomorphism is represented by an $(n - k) \times n$ matrix, called a *check matrix* of C . In our case the dimension of the space is $(n_1 + 2n_2 + \dots + sn_s)$, we have a natural basis, and the coordinates of a vector, as well as the columns of a check matrix, are naturally divided into $n_1 + n_2 + \dots + n_s$ groups, *blocks*, the first n_1 blocks of size 1, the next n_2 blocks of size 2, and so on. For a check matrix H , by $H_{i,j}$ we denote the space spanned by the columns from the j th block of size i .

Definition 11 (multifold partition of a space). A multiset D of subspaces of a vector space S is called a μ -fold *partition* of S if every nonzero vector of S belongs to exactly μ subspaces from D , respecting the multiplicities.

Theorem 12. Let H be an $m \times (n_1 + 2n_2 + \dots + sn_s)$ rank- m matrix over \mathbb{F}_p , and let C be the null-space of H . Denote

$$k = \sum_{i=1}^s n_i(p^s - p^{s-i}) = p^s \sum_{i=1}^s n_i(1 - p^{-i}), \quad \mu = \frac{k}{p^m - 1}, \quad q_i = p^i. \quad (8)$$

The following assertions are equivalent:

- (i) C is an $\text{OA}(|C|, q_1^{n_1} q_2^{n_2} \dots q_s^{n_s}, t)$ attaining bound (6);
- (i') C is an $\text{OA}(|C|, q_1^{n_1} q_2^{n_2} \dots q_s^{n_s}, t)$ with $t = \frac{\mu+k}{p^s} - 1$;
- (ii) C is a $\{k, \mu\}$ -CR code in $H(q_1^{n_1} q_2^{n_2} \dots q_s^{n_s})$;
- (iii) the multiset $M = \{p^{s-i} \times H_{i,j}\}_{i=1}^s \}_{j=1}^{n_i}$ of subspaces of \mathbb{F}_p^m is a μ -fold partition ($p^{s-i} \times H_{i,j}$ denotes that $H_{i,j}$ is added p^{s-i} times in the multiset).

Proof. Since C is a null-space of H , we have $|C| = p^{n_1+2n_2+\dots+sn_s}/p^m = q_1^{n_1} \cdot q_2^{n_2} \cdot \dots \cdot q_s^{n_s}/p^m$.

(i) \iff (i') Taking into account the expression for $|C|$ above, (6) has the form

$$p^{-m} \geq \left(1 - \left(1 - \frac{1}{q}\right) \frac{n}{t+1}\right). \quad (9)$$

(recall that q is the harmonic mean of $\underbrace{q_1, \dots, q_1}_{n_1 \text{ times}}, \underbrace{q_2, \dots, q_2}_{n_2 \text{ times}}, \dots, \underbrace{q_s, \dots, q_s}_{n_s \text{ times}}$). From

the equality in (9), we find

$$t+1 = \frac{n(1 - \frac{1}{q})}{1 - p^{-m}} = \frac{p^{-s}k}{1 - p^{-m}} = p^{-s}k \left(1 + \frac{1}{p^m - 1}\right) = \frac{k + \mu}{p^s}.$$

Inversely, substituting $t = \frac{k+\mu}{p^s} - 1$ turns (9) to equality.

(i) \iff (ii) By Theorem 9, C attains (6) if and only if it is a $\{k; -\theta_{t+1}\}$ -CR code, where k is the degree of $H(q_1^{n_1} q_2^{n_2} \dots q_s^{n_s})$ and $\theta_{t+1} = k - p^s(t+1)$ (the $(t+1)$ th largest eigenvalue of $H(q_1^{n_1} q_2^{n_2} \dots q_s^{n_s})$, counting from 0).

(ii) \iff (iii) Assume that (iii) holds and we have to show (ii). We first check that

(*) $H_{i,j}$ has dimension i for each i, j .

Indeed, if it is so, then $H_{i,j}$ has $(p^i - 1)$ nonzero points, $p^{s-i} \times H_{i,j}$ has $(p^s - p^{s-i})$ nonzero points, and M has $\sum_{i=1}^s n_i(p^s - p^{s-i})$, i.e., k , nonzero points. This is exactly the number of points we need to cover all $p^m - 1$ nonzero points of \mathbb{F}_p^m with multiplicity $\mu = \frac{k}{p^m - 1}$.

But if one $H_{i,j}$ has dimension smaller than i , then the number of points in M is not enough to make a μ -fold partition of \mathbb{F}_p^m . So, (*) holds.

Next, (*) means that C is independent. Indeed, if there are two adjacent vectors x and y in C , then their difference $x - y$ has nonzero values only in coordinates from one block, say (i, j) th. In this case, these values are coefficients of a nontrivial linear dependency between the corresponding columns, and hence $H_{i,j}$ has dimension smaller than i , contradicting (*).

The independence of C implies that every its element is connected to elements not in C by k edges (k is the degree of the graph), and it remains to confirm the second parameter μ of the $\{k; \mu\}$ -CR code.

For a vector v not in C denote by s its *syndrome* $H \cdot v^T$ (since $v \notin C$, the syndrome s is nonzero). A vector $v - e$ is adjacent to v and belongs to C if and only if $H \cdot e^T = s$ and e has zeros in all positions out of one block, say (i, j) th. The last is equivalent to $s \in H_{i,j}$. Since the subspace $H_{i,j}$ is counted with multiplicity p^{s-i} in the multiset M , and the multiplicity of the edge $\{v, v - e\}$ is also p^{s-i} , the number of edges between v and C is equal to the multiplicity of s in M , i.e., μ .

By reversing the arguments, we ensure that (ii) implies (iii). \square

A collection of subspaces of \mathbb{F}_p^m of dimension at most s satisfying the condition of Theorem 12 is called a $(\lambda, \mu)_p^{s,m}$ -multispread [12], where $\lambda = n_1(p^{s-1} - 1) + n_2(p^{s-2} - 1) + \dots + n_{s-1}(p^1 - 1)$ (the multiplicity of the zero vector in M). Theorem 12 means that such multispreads with n_i subspaces of dimension i , $i = 1, \dots, s$, corresponds to orthogonal arrays in $(\mathbb{F}_p^1)^{n_1} \times \dots \times (\mathbb{F}_p^s)^{n_s}$ attaining the generalized Bierbrauer–Friedman bound (6) in Theorem 9. Note that n_1, \dots, n_s are not uniquely determined from the parameters p, s, m, λ, μ of a $(\lambda, \mu)_p^{s,m}$ -multispread, and even without fixing n_1, \dots, n_s the problem of existence of $(\lambda, \mu)_p^{s,m}$ -multispread is open in general (in [12], it is completely solved for $s = 2$, any p , and for $p^s \in \{2^3, 2^4, 3^3\}$). The pure-level subcase of Theorem 12 was proved in [1, Theorem 4.8]; in that case, $\lambda = 0$, and multispreads are μ -fold spreads, whose parameters are characterized, see [11, p.83], [6, Corollary 8].

Example 1. Consider the $3 \times (1 \cdot 1 + 4 \cdot 2)$ matrix

$$H = \begin{pmatrix} 1 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 0 & 1 & 0 & 1 & 1 \\ 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 1 \end{pmatrix}$$

over \mathbb{F}_2 . Its null-space is an additive OA(64, $2^1 4^4$, 3). Indeed, $64 = 2^9(1 - (1 - \frac{3}{10})\frac{5}{3+1})$; the degree of the graph $H(2^1 4^4) = 2K_2 \times K_4 \times K_4 \times K_4 \times K_4$ is $1+4 \cdot 3 = 14$; the eigenvalue $\theta_{t+1} = \theta_4$ equals $14 - 4 \cdot 4 = -2$; the subspace $H_{1,1} = \langle (1, 1, 1) \rangle$, corresponding to the first column, is taken with multiplicity 2 and covers the vector $(1, 1, 1)$ exactly $2 = -\theta_4$ times; each of the other nonzero vectors from \mathbb{F}_2^2 is covered exactly $2 = -\theta_4$ times by the subspaces $\langle (1, 0, 0), (0, 1, 0), \langle (1, 0, 0), (0, 0, 1) \rangle, \langle (0, 1, 0), (0, 0, 1) \rangle, \langle (1, 1, 0), (0, 1, 1) \rangle$, formed by the last four blocks of columns.

5 Polynomial generalization of the Bierbrauer–Friedman bound

For a connected graph G , $d(G)$ denotes its diameter, and $G^{(i)}$, denotes a simple graph where two vertices are adjacent if and only if the distance between them in G is i .

Definition 13 (distance-regular graph). A simple connected graph is called a *distance-regular graph* if for $i = 1, \dots, d(G)$ the product $A_1 A_i$ is a linear combination of A_{i-1} , A_i , and A_{i+1} .

Corollary 14. For every distance-regular graph G , there are degree- i polynomials $K^{(i)}$, $i = 0, \dots, d(G)$, such that $G^{(i)} = K^{(i)}(G)$.

Let G be a distance-regular graph. Consider a nonnegative integer linear combination of the polynomials $K^{(i)}$, i.e., $P = \sum_i a_i K^{(i)}$, $a_i \geq 0$. Then $P(A)$ is the adjacency matrix of the multigraph $P(G)$ with the same set V of vertices. If vertices u and v are at distance i from each other in G , then the edge $\{u, v\}$ has multiplicity a_i in $P(G)$.

Proposition 15. (a) *The set of eigenfunctions of $P(G)$ includes the set of eigenfunctions of G . If P is strictly monotonic function, then these sets coincide.*

(b) *If φ is an eigenfunction of G with eigenvalue θ , then φ is also an eigenfunction of $P(G)$ with eigenvalue $P(\theta)$.*

(c) *If P strictly increases, then sets of algebraic t -designs of G and $P(G)$ coincide.*

The next lemma is a variant of Lemma 8.

Lemma 16. *Let $G = (V, E)$ be a distance-regular graph with eigenvalues $k = \theta_0, \theta_1, \dots, \theta_{d(G)}$, in the decreasing order, let the polynomial P be a linear combination*

$$P(x) = \alpha_1 K^{(1)}(x) + \dots + \alpha_{d(G)} K^{(d(G))}(x)$$

of the polynomials $K^{(1)}$ from Corollary 14, where $\alpha_i \geq 0$, $i = 1, \dots, d(G)$, and let

$$\mu = \max \{P(\theta_{t+1}), \dots, P(\theta_{d(G)})\}.$$

If $\mu < 0$, then the cardinality of an algebraic t -design C satisfies the inequality

$$\frac{|C|}{|V|} \geq \frac{-\mu}{P(k) - \mu}. \quad (10)$$

Proof. If all α_i are integer, then $P(x)$ is a graph, for which Lemma 8 gives (10).

If all α_i are rational, we divide P by their greatest common divisor, which makes all α_i integer and does not affect the value in (10).

If some α_i are irrational, we can approximate them by rational values with any accuracy and get (10) as the limit of bounds corresponding to polynomials P with rational coefficients. \square

Our aim is to find a polynomial P such that $\theta_{t+1} \geq 0$ but $\mu < 0$. In this case, the lower bound from Lemma 8 is non-positive and hence trivial, while the lower bound from Lemma 16 is positive and can be valuable. Actually, to obtain the best lower bound in (10), we need to solve the following linear programming problem in the variables $\alpha_1, \dots, \alpha_{d(G)}, \mu$ (the last one is an additional variable, which plays the role of $\max\{\theta_j\}_{j \geq t+1}$):

$$\left\{ \begin{array}{l} \sum_{i=1}^{d(G)} \alpha_i K^{(i)}(\theta_j) \leq \mu, \quad j = t+1, \dots, d(G), \\ \alpha_i \geq 0, \quad i = 1, \dots, d(G), \\ \sum_{i=1}^{d(G)} \alpha_i K^{(i)}(\theta_0) = \text{const} \quad (\text{the degree of } P(G)), \\ \mu \rightarrow \min. \end{array} \right.$$

By the proof of Lemma 16, we do not need to care about the integrity of α_i . By Lemmas 8, 16, and Proposition 15(c) we can conclude the following.

Proposition 17. *Let a collection $a_1, \dots, a_{\text{d}(G)}$ be the solution of the linear program and let P be the corresponding polynomial. Suppose that $P(\theta_{t+1}) \neq P(\theta_j)$ for $j \neq t+1$. If some t -design C attain the bound (10), then C is a $\{b; c\}$ -CR code in G , for some b and c .*

For the Hamming graph $G = H(n, q)$, the degree- w polynomial $K^{(w)}$ is

$$K^{(w)}(\cdot) = P_w(P_1^{-1}(\cdot)),$$

$$\text{where } P_w(x) = P_w(x; n, q) = \sum_{j=0}^w (-1)^j (q-1)^{w-j} \binom{x}{j} \binom{n-x}{w-j}$$

is the *Krawtchouk polynomial*; $P_1(x) = (q-1)n - qx$, $P_1^{-1}(y) = \frac{(q-1)n - y}{q}$.

Example 2. Consider $G = H(2m, 2)$, $t = m - 1$. Let us derive a bound on the size of $\text{AO}(N, 2m, 2, m-1)$ by combining only three polynomials $K^{(i)}$, $i = 1, 2, 3$. We have $G = H(2m, 2)$, $\theta_m = 0$,

$$K^{(1)}(x) = x, \quad K^{(2)}(x) = \frac{1}{2}(x^2 - 2m), \quad K^{(3)}(x) = \frac{1}{6}(x^3 - (6m-2)x).$$

We are looking for a polynomial $P = K^{(3)} + \beta K^{(2)} + \alpha K^{(1)}$ where $\alpha \geq 0$, $\beta \geq 0$, and $P'(x) \geq 0$ for all $x \in [-2m, 0]$. Put $\alpha = \frac{\beta^2}{2} + m - \frac{1}{3}$. Then $6P(x) = (x + \beta)^3 - \beta^3 - 6\beta m$. It is easy to see that P increases monotonically. Substituting $\beta = \frac{m}{\sqrt{3}}$, we obtain $6P(x) = x^3 + \sqrt{3}mx^2 + m^2x - 2\sqrt{3}m^2$ and, by (10),

$$\frac{|C|}{|V|} \geq \frac{-6P(0)}{6P(2m) - 6P(0)} = \frac{2\sqrt{3}m^2}{10m^3 + 4\sqrt{3}m^3} = \frac{1}{m(5 \cdot 3^{-\frac{1}{2}} + 2)} > \frac{0.2046}{m}.$$

6 Conclusion

We have considered two possibilities to generalize the Bierbrauer–Friedman bound for orthogonal arrays. One of them concerns mixed-level orthogonal arrays, and we have also show how to construct arrays attaining the generalized bound. However, the constructed arrays have one restriction because of their additive structure over a finite field; namely, the size of such an array is always a prime power. As was mentioned in the introduction, the only known arrays attaining the Bierbrauer–Friedman bound with the size not being a prime power are pure-level binary or ternary. With generalizing the bound the problem of the existence of such arrays also expands to mixed-level case. In particular: do there exist mixed-level $\text{OA}(N, 2^{n_1} 3^{n_2}, t)$ attaining bound (6)? Orthogonal arrays attaining the considered bound (in both pure- and mixed-level cases) are of special interest because of their relation with such regular structures as CR-1 codes (equivalently, intriguing sets, equitable 2-partitions, perfect 2-colorings).

References

- [1] E. A. Beshpalov, D. S. Krotov, A. A. Matushev, A. A. Taranenko, and K. V. Vorob'ev. Perfect 2-colorings of Hamming graphs. *J. Comb. Des.*, 29(6):367–396, June 2021. <https://doi.org/10.1002/jcd.21771>.
- [2] J. Bierbrauer. Bounds on orthogonal arrays and resilient functions. *J. Comb. Des.*, 3(3):179–183, 1995. <https://doi.org/10.1002/jcd.3180030304>.
- [3] A. E. Brouwer and W. H. Haemers. *Spectra of Graphs*. Springer-Verlag, New York, 2012. <https://doi.org/10.1007/978-1-4614-1939-6>.
- [4] P. Delsarte. *An Algebraic Approach to Association Schemes of Coding Theory*, volume 10 of *Philips Res. Rep., Supplement*. N.V. Philips' Gloeilampenfabrieken, Eindhoven, 1973.
- [5] W. S. Diestelkamp. Parameter inequalities for orthogonal arrays with mixed levels. *Des. Codes Cryptography*, 33(3):187–197, 2004. <https://doi.org/10.1023/B:DESI.0000036245.92856.f0>.
- [6] S. El-Zanati, G. Seelinger, P. Sissokho, L. Spence, and C. Vanden Eynden. On λ -fold partitions of finite vector spaces and duality. *Discrete Math.*, 311(4):307–318, 2011. <https://doi.org/10.1016/j.disc.2010.10.026>.
- [7] D. G. Fon-Der-Flaass. Perfect 2-colorings of a hypercube. *Sib. Math. J.*, 48(4):740–745, 2007. <https://doi.org/10.1007/s11202-007-0075-4> translated from *Sib. Mat. Zh.* 48(4) (2007), 923–930.
- [8] J. Friedman. On the bit extraction problem. In *Foundations of Computer Science, IEEE Annual Symposium on*, pages 314–319, Los Alamitos, CA, USA, 1992. IEEE Computer Society. <https://doi.org/10.1109/SFCS.1992.267760>.
- [9] S. Goryainov and D. Krotov. Completely regular codes in graphs covered by a Hamming graph. arXiv eprint 2411.09698, DOI10.48550/arXiv.2411.09698, 2024.
- [10] A. S. Hedayat, N. J. A. Sloane, and J. Stufken. *Orthogonal Arrays. Theory and Applications*. Springer Series in Statistics. Springer, New York, NY, 1999. <https://doi.org/10.1007/978-1-4612-1478-6>.
- [11] J. W. P. Hirschfeld. *Projective Geometries Over Finite Fields*. Oxford University Press, New York, 1979.
- [12] D. S. Krotov and I. Yu. Mogilnykh. Multispreads. E-print 2312.07883, arXiv.org, 2023. Available at <http://arxiv.org/abs/2312.07883>.
- [13] D. S. Krotov and V. N. Potapov. Completely regular codes and equitable partitions. In M. Shi and P. Solé, editors, *Completely Regular Codes in Distance Regular Graphs*, chapter 1. CRC Press, 2024. To appear.

- [14] C. R. Rao. Factorial experiments derivable from combinatorial arrangements of arrays. *J. R. Stat. Soc., Suppl.*, 9(1):128–139, 1947. <https://doi.org/10.2307/2983576>.