

# RX J1420.4+5334 – another tidal disruption event? \*

J. Greiner<sup>1</sup>, R. Schwarz<sup>1</sup>, S. Zharikov<sup>2,3</sup>, M. Orio<sup>4,5</sup>

<sup>1</sup> Astrophysical Institute Potsdam, An der Sternwarte 16, 14482 Potsdam, Germany

<sup>2</sup> Special Astrophysical Observatory, 357147 Nizhnij Arkhyz, Russia

<sup>3</sup> OAN, Instituto de Astronomía, UNAM, 22860 Ensenada, México

<sup>4</sup> Osservatorio Astronomico di Torino, Strada Osservatorio 20, I-10125 Pino Torinese (TO), Italy

<sup>5</sup> Department of Physics, University of Wisconsin, 1150 Univ. Av., Madison, WI 53706, USA

Received 18 August 2000; accepted 25 September 2000

**Abstract.** We have discovered a transient X-ray source, RX J1420.4+5334, which displays a ROSAT flux variation of  $\gtrsim 150$  between the ROSAT All-Sky-Survey in 1990 and a preceding pointed ROSAT observations in July 1990. Optical observations suggest a non-active galaxy as the only visible counterpart. We therefore tentatively identify RX J1420.4+5334 as a tidal disruption event in a non-active galaxy.

**Key words:** X-rays: galaxies – Accretion – Galaxies: nuclei – Galaxies: individual: RX J1420.4+5334

## 1. Introduction

Tidal disruption of stars in the gravitational potential of massive black holes in the centers of galaxies has been recognized long ago as a way to prove the existence of massive black holes in otherwise non-active galaxies. During the tidal disruption about 50% of the mass of the star are stripped off and get unbound from the star (mostly independent of impact parameter, masses and velocities; Ayal et al. 2000). This material can eventually be accreted by the black hole, leading to a flare of optical to X-ray emission lasting a few months to years. While originally proposed to explain the activity in AGN (Hills 1975), a first calculation of the frequency, temporal behaviour and spectral signatures of tidal disruption events (Frank & Rees 1976, Young et al. 1977, Kato & Hoshi 1978, Lidskii & Ozernoi 1979, Gurzadyan & Ozernoi 1980, Gurzadyan & Ozernoi 1981) revealed both, the potential of such events to learn more about the inner regions of galaxies as well as the problems in observing these rare events (Rees 1988).

Send offprint requests to: J. Greiner, jgreiner@aip.de


\* Partly based on observations collected at the German-Spanish Astronomical Centre, Calar Alto, operated by the MPI für Astronomie, Heidelberg, jointly with the Spanish National Commission for Astronomy, and the WIYN telescope, operated by the University of Wisconsin, Indiana University, Yale University, and the National Optical Astronomy Observatories.

Over the recent few years, tidal disruption has attracted renewed attention, both theoretically and observationally. Loeb & Ulmer (1997) have studied the case that a surrounding gaseous envelope could reprocess most of the radiation from the accretion process into a thermal spectrum of  $10^4$  K, thus being easily detectable in optical surveys rather than in X-ray surveys. Magorrian & Tremaine (1999) have recalculated the stellar disruption rates in detailed dynamical models of real galaxies, taking into account the refilling of the loss cone of stars on disruptable orbits by two-body relaxation and tidal forces in non-spherical galaxies. Ayal et al. (2000) have performed simulations of the long-term evolution of tidally disrupted stars and find that only about 10% of the stellar mass are actually accreted by the central black hole.

On the observational side, the discovery of time-variable UV or X-ray emission in distant, otherwise non-active galaxies may provide the first indirect evidence of tidal disruption events. Besides the early report on a UV flare in NGC 4552 (Renzini et al. 1995), a few very good candidates have been detected with ROSAT during the recent years, such as the  $\sim 2 \times 10^{43}$  erg/s outburst in NGC 5905 (Bade et al. 1996, Komossa & Bade 1999), the  $> 9 \times 10^{43}$  erg/s flare in RXJ 1242.6–1119 (Komossa & Greiner 1999), the  $\sim 10^{44}$  erg/s flare in RXJ 1624.9+7554 (Grupe et al. 1999), or the  $\sim 10^{44}$  erg/s flare in RXJ 1331.9–3243 (Reiprich & Greiner 2000). Interestingly, in all these ROSAT discoveries the X-ray spectrum of the flares is extremely soft, of the order of 50–100 eV effective temperature. The time scales of these events, though not in all cases well constrained, are of the order of months to years. All these properties, together with the fact that none of the corresponding galaxies shows any optical sign of activity have led to the tentative identification as being associated with disruption events.

Here we report on the discovery of another transient X-ray source and the optical and X-ray follow-up observations.

## 2. Observational results




**Fig. 1.** Long-term X-ray light curve of RX J1420.4+5334 with the only secure detection during the ROSAT All-Sky-Survey. Because of the very soft spectrum we have used a factor 7.8 for the HRI to PSPC count rate conversion (Greiner et al. 1997). Upper limits are shown with arrows, and the 1995 HRI detection of three photons is marked with an open triangle. The inset shows the light curve during the three days of the All-Sky-Survey. The large, vertical dashes at the bottom denote the times of optical observations.

### 2.1. ROSAT All-Sky-Survey

The search for supersoft X-ray sources in the ROSAT PSPC All-Sky-Survey (Greiner 1996) revealed a few sources for which neither confirmation through a ROSAT follow-up observation nor a reasonable optical counterpart could be found. One of these sources is RX J1420.4+5334  $\equiv$  1RXS J142024.4+533403  $\equiv$  RBS 1376 (Schwope et al. 2000). It was detected during the PSPC All-Sky-Survey between December 5–8, 1990 at a count rate of 0.28 cts/s, resulting in the collection of a total of 255 counts. There is no significant short-term variability within the 3 day scanning coverage. The source position has been determined using only photons above 0.25 keV (to avoid position deterioration by ghost images) to be: RA (2000.0) =  $14^{\text{h}}20^{\text{m}}24^{\text{s}}.2$ , Decl. (2000.0) =  $+53^{\circ}34'11''$  with an error of  $\pm 20''$ .

The spectrum is extremely soft as indicated by the hardness ratio:  $HR1 = -0.92 \pm 0.03$  which is defined as  $(H-S)/(H+S)$ , with H (S) being the counts above (below) 0.4 keV over the full PSPC range of 0.1–2.4 keV. Source photons were extracted with a radius of  $4'$ . The background was chosen at the same ecliptic longitude at  $\approx 1^{\circ}$  distance, corresponding to background photons collected typically 15 sec before or after the time of the source photons. Standard corrections were applied using the dedicated EXSAS software package (Zimmermann et al. 1994). The spectrum fitted with a blackbody model is shown in Fig. 2, and Tab. 2 summarizes the fit parameters, also including fits with a power law and an accretion disk model.



**Fig. 2.** Fit of a black body model to the X-ray spectrum of RX J1420.4+5334 as measured during the ROSAT All-Sky-Survey. The lower left panel shows the deviation between model and data in units of  $\chi^2$  per spectral bin.


### 2.2. ROSAT pointed observations

Follow-up pointed ROSAT observations were performed on July 10/11, 1995 and again on June 3, 1997. In both cases the HRI was used. In the 1995 observation we detect three X-ray photons at the position of RX J1420.4+5334, corresponding to a likelihood of 6.2 ( $2.5 \sigma$ ). Because of this marginal significance, we do not claim a detection, but rather use these detected photons as an upper limit (Fig. 1) on the count rate:  $5.5 \times 10^{-4}$  HRI cts/s. The mean position of these photons, RA (2000.0) =  $14^{\text{h}}20^{\text{m}}23^{\text{s}}.6$ , Decl. (2000.0) =  $+53^{\circ}34'16''$  with an error of  $\pm 10''$  is shown in Fig. 3 for comparison purposes. The 1997 observation, with less than half of the 1995 exposure, results in an upper limit of  $1.2 \times 10^{-3}$  HRI cts/s.

RX J1420.4+5334 is serendipitously in the field of view of a ROSAT PSPC pointed observation performed during the calibration phase, though at a very large off-axis angle of  $49'$ . No source was detected (using standard EXSAS commands), giving a  $3\sigma$  upper limit of  $1.8 \times 10^{-3}$  PSPC cts/s (Fig. 1).

### 2.3. Optical observations

Investigation of the X-ray source position on the POSS revealed only one bright optical object (labelled “A” in Fig. 3), at RA (2000.0) =  $14^{\text{h}}20^{\text{m}}24^{\text{s}}.4$ , Decl. (2000) =  $53^{\circ}34'12''$ . We obtained a low-resolution spectrum of this R  $\approx 18.8$  mag source using the 3.5m telescope at Calar Alto on June 13, 1993. We used the Cassegrain spectrograph with a  $1024 \times 640$  RCA chip (pixel size  $15 \mu\text{m}$ ). A grating with  $240 \text{ \AA}/\text{mm}$  was used, allowing the range 4000–8500  $\text{\AA}$  to be observed with a FWHM resolution of 10  $\text{\AA}$ . The spectrum was bias and flat-field corrected and calibrated in wavelength using standard MIDAS reduction packages. BD  $+25^{\circ}4655$  was used for flux calibration.



**Fig. 3.** Finding chart of RX J1420.4+5334 based on the sum of four B band images (total exposure of 2000 sec) taken on May 26, 1996 at the 3.5m WIYN telescope. The large circle marks the  $3\sigma$  X-ray error box of the ROSAT All-Sky-Survey detection ( $20''$  radius), while the smaller corresponds to the position of the photons detected in the HRI ( $10''$  radius). The inset shows a zoomed portion of the central part.

The optical spectrum is characterized by strong absorption lines of NaI  $5175\text{\AA}$ , MgI  $5890\text{\AA}$  and  $H\alpha$  (though it is near the atmospheric A band), typical of an elliptical or early spiral type. Using the MgI, NaI and  $H\alpha$  line we derive a redshift of  $z = 0.147 \pm 0.001$ . The Balmer absorption lines and the strong drop of the flux beyond the CaII H/K break argue against a classification as BL Lac object. No AGN-like forbidden emission lines, like  $[\text{OIII}]\lambda 5007$ , are detected.

In order to search for optical variability, we embarked on a more extensive photometry programme. Observations were done at three locations, the Sonneberg Observatory (Germany), the Special Astrophysical Observatory (Russia) and at the WIYN telescope at Kitt Peak (USA). Tab. 1 summarizes the logistic details. Throughout these observations we did not found any significant optical variability.

The imaging revealed a second, much fainter ( $R \approx 20.6$  mag) source within the X-ray error circle (labelled “B” in Fig. 3). During some of the WIYN observations, at very good seeing conditions of  $0''.9$ , this source is read-

**Table 1.** Log of Observations

Telescope <sup>(1)</sup>	Date	Filter/Wavelength	D <sup>(2)</sup> (hrs)	T <sub>Int</sub> (sec)
<b>X-ray</b>				
R 150046P	1990 Jul 19–22	0.1–2.4 keV	77.7	11 475
R survey	1990 Dec 5–8	0.1–2.4 keV	77.8	910
R 201999P	1995 Jul 10/11	0.1–2.4 keV	30.7	5560
R 300577H	1997 Jun 3	0.1–2.4 keV	0.6	2150
<b>Optical</b>				
CA 3.5m	1993 Jun 13	4000–8500		3600
SO 0.6 m	1994 Mar 30	white	2.5	30
SO 0.6 m	1994 Apr 14	white	2.9	60
SO 0.6 m	1994 Apr 19	white	3.2	30
SO 0.6 m	1994 May 27	white	3.0	60
WIYN 3.5m	1996 May 25	R/B	1.0	270
WIYN 3.5m	1996 May 26	B	0.3	500
WIYN 3.5m	1996 May 27	B	0.1	500
SAO 1.0m	1996 Jul 14	V	1.5	300

<sup>(1)</sup> The abbreviations mean: R = ROSAT, SO = Sonneberg Observatory (Germany), CA = Calar Alto (Spain), SAO = Special Astrophysical Observatory (Russia).


<sup>(2)</sup> Duration of the coverage.

ily seen to be extended (see inset of Fig. 3). Moreover, the intensity distribution over the galaxy disk is very flat. Since active galaxies typically have a very peaked intensity profile (dominated by a bright central source), this flat distribution argues for a non-active galaxy. The shape and radial profile suggest an elliptical or early spiral subtype. Size and luminosity arguments then suggest that its distance is about a factor 3–5 larger than that of source “A”. A lower limit of  $z \sim 0.07$  can be placed by assuming  $M_V \sim -16$  mag, corresponding to the faint end of the galaxy luminosity distribution.

Certainly, a spectrum of this galaxy would be very helpful to verify these suggestions.

### 3. Discussion

We have presented evidence for a transient X-ray source with an amplitude of a factor of  $>150$  within 6 months. The only optical counterparts down to 23rd magnitude are two galaxies, the brighter of which is at a redshift of  $z=0.147$ , and the fainter one probably a factor 3–5 more distant. If the X-ray flare occurred in the brighter galaxy, the unabsorbed, bolometric X-ray luminosity is  $2.5 \times 10^{44}$  erg/s (using the disk blackbody model with only galactic foreground absorption and a normalization which fits the  $z=0.147$  distance; see last line in Tab. 2). It should be mentioned that this is a lower limit since most probably we did not catch the peak of the flare, and the X-ray emission may have suffered additional intrinsic absorption. If the flare occurred in the fainter galaxy (B) then the luminosity would probably be even larger, but certainly not lower than  $6 \times 10^{43}$  erg/s (if at  $z \sim 0.07$ ).



**Fig. 4.** Optical spectrum of RX J1420.4+5334, source “A”. Prominent absorption lines are marked.

**Table 2.** Spectral fit results of RX J1420.4+5334

Model <sup>(1)</sup>	Fit parameters <sup>(2)</sup>		$\chi^2_{red}$	Flux <sup>(3)</sup> (0.1-2.4 keV)	Flux <sup>(3)</sup> (bol.)
bbdy, free fit	$kT = 28 \pm 10$	$N_H = 2.0 \pm 0.1$	0.96	$7.9 \times 10^{-12}$	$1.6 \times 10^{-11}$
bbdy, $N_H = N_H^{gal}$	$kT = 38 \pm 10$		0.90	$8.8 \times 10^{-13}$	$4.1 \times 10^{-12}$
powl, free fit	$\alpha = -7.5 \pm 1.5$	$N_H = 2.8 \pm 1.0$	0.96	$2.4 \times 10^{-12}$	–
powl, $N_H = N_H^{gal}$	$\alpha = -4.8 \pm 0.5$		1.07	$8.9 \times 10^{-13}$	–
dibb, free fit	$\dot{M} = 0.01 \pm 0.005$	$M = 10^6$ (fix) $N_H = 6.6 \pm 1.5$	0.94	$5.0 \times 10^{-10}$	$1.8 \times 10^{-6}$
dibb, $N_H = N_H^{gal}$	$\dot{M} = 2.6 \pm 0.1$	$M = 10^6$ (fix)	0.90	$9.2 \times 10^{-13}$	$6.2 \times 10^{-12}$
dibb, $N_H = N_H^{gal}$	$\dot{M} = 1.8 \pm 0.1$	$M = 7 \times 10^5$ Norm <sup>(4)</sup> = $2.9 \times 10^{-10}$ (fix)	0.90	$9.2 \times 10^{-13}$	$6.4 \times 10^{-12}$

<sup>(1)</sup> The galactic column density is  $N_H^{gal} = 1.2 \times 10^{20} \text{ cm}^{-2}$  (Dickey & Lockman 1990). The model abbreviations have the following meaning: bbdy = black body, powl = power law, dibb = disk black body model.

<sup>(2)</sup> Units are eV for temperatures,  $M_\odot$  for masses,  $10^{20} \text{ cm}^{-2}$  for  $N_H$ .

<sup>(3)</sup> Observed fluxes in units of  $\text{erg cm}^{-2} \text{ s}^{-1}$  (absorption-corrected).

<sup>(4)</sup> Normalization is chosen such that distance is equal to  $D_L = c/H * z = 588 \text{ Mpc}$  (and inclination assumed to be face-on).

Though it is impossible to claim with certainty that this evidence proves a tidal disruption origin, we will in the following, by the principle of exclusion, discuss possible alternatives (for a more extensive discussion of alternatives see Komossa & Bade 1999). (1) An active galaxy as the most common origin of large X-ray variability is very unlikely given our optical observations (see above section 2.3). (2) X-ray afterglows of gamma-ray bursts (GRB) may appear as new, bright X-ray sources on the sky. However, all of the observed afterglows have faded rapidly (with time<sup>-0.8...-2.1</sup>; see Greiner 2000). The ROSAT all-sky survey data suggests a rather flat light curve over about 3 days, inconsistent with a GRB afterglow. (3) This flat, 3-day long light curve also excludes neutron star merger events which are predicted to reach peak luminosities of  $10^{44} \text{ erg/s}$ , but should fade by a factor of  $\sim 3$  within  $\sim 2$  days (Li & Paczynski 1998).

Thus, a tidal disruption scenario seems a logical consequence. Moreover, it is even plausible: Spectral fitting of a simple disk blackbody model (Shakura & Sunyaev 1973) gives a consistent picture concerning X-ray temperature, mass of the central object ( $\sim 10^6 M_\odot$ ), luminosity and distance (see last row of Tab. 2).

*Acknowledgements.* We are grateful to E. Pavlenko (Crimean Observatory) and W. Wenzel (Sonneberg) for initial optical observations in 1994, as well as to N. Borisov for help in observations conducted at SAO. We appreciate enlightening discussions with M. Salvato (AIP) on the radial profiles of galaxies. JG was partly supported by the German Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie (BMBF/DLR) under contract 50 QQ 96 02 3. The ROSAT project was supported by BMBF/DLR and the Max-Planck-Society.

## References

- Ayal S., Livio M., Piran T., 2000, ApJ (in press; astro-ph/0002499)
- Bade N., Komossa S., Dahlem M., 1996, A&A 309, L35
- Dickey J.M., Lockman F.J., 1990, ARAA 28, 215
- Frank J., Rees M.J., 1976, MNRAS 176, 633
- Greiner J., 1996, in *Supersoft X-ray Sources*, ed. J. Greiner, Lect. Notes in Phys. 472, Springer, p. 285
- Greiner J., Bickert K.F., Luthardt R., Viotti R., Altamore A., González-Riestra R., Stencel R.E., 1997, A&A 322, 576
- Greiner J., 2000, <http://www.aip.de/People/JGreiner/grb.html>
- Grupe D., Thomas H.-C., Leighly K.M., 1999, A&A 350, L31
- Gurzadyan V.G., Ozernoi L.M., 1980, A&A 86, 315
- Gurzadyan V.G., Ozernoi L.M., 1981, A&A 95, 39
- Kato M., Hoshi R., 1978, Prog. Theor. Phys. 60, 1692
- Komossa S., Bade N., 1999, A&A 343, 775
- Komossa S., Greiner J., 1999, A&A 349, L45
- Li L.-X., Paczynski B., 1998, ApJ 507, L59
- Lidskii V.V., Ozernoi L.M., 1979, Pis'ma Astron. Zh. 5, 28 (Soviet Astron. Lett. 5, 16)
- Magorrian J., Tremaine S., 1999, MNRAS 309, 447
- Rees M., 1988, Nature 333, 523
- Reiprich T.H., Greiner J., 2000, in *Black Holes in Binaries and Galactic Nuclei*, eds. L. Kaper, E.P.J. van den Heuvel, and P.A. Woudt, Springer (in press)
- Renzini A., Greggio L., Di Serego-Alighieri S. et al., 1995, Nat. 378, 39
- Schwope A.D., Hasinger G., Lehmann I., et al. 2000, AN 321, 1
- Shakura N.I., Sunyaev R.A., 1973, A&A 24, 337
- Young P.J., Shields G.A., Wheeler J.C., 1977, ApJ 212, 367
- Zimmermann H.U., Becker W., Belloni T., Döbereiner S., Izzo C., Kahabka P., Schwentker O., 1994, MPE report 257