

Pressure dependence and non-universal effects of microscopic couplings on the spin-Peierls transition in CuGeO_3

R. Raupach, A. Klümper, F. Schönfeld

Institut für Theoretische Physik, Universität zu Köln, Zùlpicher Str. 77, 50937 Köln, Germany

The theory by Cross and Fisher (CF) is by now commonly accepted for the description of the spin-Peierls transition within an adiabatic approach. The dimerization susceptibility as the essential quantity, however, is approximated by means of a continuum description. Several important experimental observations can not be understood within this scope. Using density matrix renormalization group (DMRG) techniques we are able to treat the spin system exactly up to numerical inaccuracies. Thus we find the correct dependence of the equation of state on the spin-spin interaction constant J , still in an adiabatic approach. We focus on the pressure dependence of the critical temperature which is absent in the CF theory as the only energy scale with considerable pressure dependence is J which drops out completely. Comparing the theoretical findings to the experimentally measured pressure dependence of the spin-Peierls temperature we obtain information on the variation of the frustration parameter with pressure. Furthermore, the ratio of the spectral gap and the transition temperature is analyzed.

I. INTRODUCTION

Low dimensional quantum systems are currently of considerable interest mainly due to the fascinating phase transitions driven by strong quantum fluctuations. The continuous interest from the theoretical side is provoked by the discovery of many experimental systems realizing quasi one-dimensional quantum systems. In the field of spin-Peierls systems the discovery of the inorganic compound CuGeO_3 realized a milestone as many measurements have been performed with high accuracy since. Therefore, CuGeO_3 has attracted much attention in experimental as well as in theoretical works. The high temperature behaviour of CuGeO_3 was found to be modelled adequately by one-dimensional frustrated Heisenberg chains [1–4]. In the dimerized phase, many features were shown to be consistent within an adiabatic description of the phonon degrees of freedom. This observation comprises zero temperature [5–8] as well as thermodynamic properties [4].

Even in one space dimension only a few exact results exist particularly concerning thermodynamics. For integrable systems the thermodynamical potentials and asymptotic behaviour of correlation functions are known. A notorious problem is posed by response functions and non-integrable systems in general. With respect to this, the recently developed transfer matrix DMRG (TMRG) [9–11] on the basis of transfer matrices [12] provides a very powerful method to calculate thermodynamic quantities of spin chains without any use of perturbative methods. This has been demonstrated in several applications [4,11,13–17].

In this paper we study the influence of microscopic coupling constants on the spin-Peierls transition temperature. This allows for an understanding of the considerable pressure dependence of the phase diagram

along the following line of reasoning. It is known that external pressure affects the magnetic properties of CuGeO_3 considerably [3,18,19]. Fits for the magnetic susceptibility yield the change of the nearest-neighbour spin interaction J and estimates for the frustration parameter α . Using these data for J as function of pressure we are able to explain the observed increase of the spin-Peierls temperature and estimate the pressure dependence of the next-nearest-neighbour exchange

The outline of the paper is as follows. In section II we present the model and a motivation for our description of the experimentally studied spin-Peierls systems. We study the static dimerization susceptibility in section III. Section IV is devoted to the computation of the critical temperature as function of the spin exchange couplings and external pressure, respectively. We give a comparison of our results with experimental measurements. In section V we investigate the spectral gap and its ratio to the spin-Peierls temperature. The conclusion is given in section VI.

II. MODEL

In the inorganic spin-Peierls compound CuGeO_3 the magnetic interactions are attributed to Heisenberg spin exchange. There is numerous evidence that in addition to the nearest-neighbour interaction (J) a next-nearest-neighbour exchange $J' = \alpha J$ [1–3] with $\alpha = 0.35$ has to be taken into account. Usually the constant α is referred to as frustration parameter.

At the spin-Peierls transition the system undergoes a structural phase transition driven by the quantum spin system coupled to the phonons. The spin-phonon coupling is modelled by spin exchange integrals depending linearly on the local displacements. The adiabatic treat-

ment yields a quantum spin system coupled to just one phonon mode with the commonly used Hamiltonian

$$\hat{H} = \sum_i \left\{ J[(1 + \delta_i)\mathbf{S}_i\mathbf{S}_{i+1} + \alpha\mathbf{S}_i\mathbf{S}_{i+2}] + \frac{K}{2}\delta_i^2 \right\}, \quad (1)$$

where \mathbf{S}_i are spin 1/2 operators, $\delta_i = (-1)^i\delta$ denotes the modulation of the magnetic exchange couplings in the dimerized phase. Here, we restrict ourselves to vanishing external magnetic field where the system shows a phase transition from the uniform (U), i.e. $\delta = 0$ to the dimerized (D) phase ($\delta > 0$).

The elastic energy can be expressed in terms of microscopic constants rendering (1) equivalent to an RPA treatment of the phonon propagator for the full spin-phonon system. Within RPA the condition for the phase transition is identical to that for (1) as formulated in (4) below if K is adjusted to the following value [20]

$$K = \frac{J^2}{2} \left(\sum_i \frac{\lambda_i^2}{M_i\Omega_i^2} \right)^{-1} =: \mathcal{C}J^2. \quad (2)$$

The sum runs over the spin-Peierls active modes. For each mode i , M_i denotes the effective mass of the unit cell, Ω_i the frequency of the spin-Peierls active phonon and λ_i the spin-phonon coupling constant. In particular, K is proportional to J^2 . The constant \mathcal{C} contains only the microscopic parameters of the underlying lattice.

Our numerical investigations do not improve over those of CF [21] with respect to the RPA treatment. However, within the RPA approximation we deal with the complete dynamics of the quantum spin system. Note that in Ref. [21] a continuum description of the spin system was used with a subsequent bosonization treatment which is believed to capture only the long distance asymptotics of the correlation functions.

III. RESPONSE FUNCTIONS

The static dimerization susceptibility of the spin system is defined by

$$A_\alpha(x) = -J^{-1} \lim_{\delta \rightarrow 0} \frac{\partial^2 f_\alpha(x, \delta)}{\partial \delta^2}, \quad (3)$$

where $x = T/J$, and $f_\alpha(x, \delta)$ is the free energy per site for system (1) with fixed dimerization δ , frustration α and K set to zero. The response function A is nothing but the correlation of the nearest neighbour spin exchange $\mathbf{S}_i\mathbf{S}_{i+1}$ (dimer operator) at momentum $q = \pi$ and energy $\omega = 0$. The U/D phase transition takes place for

$$A_\alpha(x_{\text{SP}}) = K/J = \mathcal{C}J. \quad (4)$$

For details the reader is referred to [4].

Let us now review the results obtained by CF. Within the bosonization approach they find $x \cdot A_{\alpha=0}(x) = \chi_0$ with $\chi_0 \approx 0.26$. As a direct consequence by use of the inversion of (4),

$$T_{\text{SP}} = JA_\alpha^{-1}(\mathcal{C}J), \quad (5)$$

this yields $dT_{\text{SP}}/dJ = 0$. Of course, this is also clear from the fact that the energy scale of the spin system completely drops out due to scale invariance.

Fig. 1 shows a comparison of our TMRG results¹ for the function $x \cdot A_\alpha(x)$ for various values of α with the findings of CF and exact data for free fermions. The enormous progress achieved by the numerical analysis is the correct treatment of the spin system on the lattice at practically all length scales. This improves over the continuum limit approach in which the asymptotics of correlation functions is incorrectly extended to short distances. For the unfrustrated Heisenberg model, i.e. $\alpha = 0$, we are able to observe directly the deviations between the continuum limit and a lattice treatment of the spin degrees of freedom.




FIG. 1. Depiction of the function $x \cdot A(x)$ with $x = T/J$ for: free fermions, Heisenberg model in continuum limit (CF), and TMRG results for frustration parameter $\alpha = 0, 0.241, 0.35$. The circles denote the relevant values for CuGeO_3 (see text).

With respect to CuGeO_3 we fix J by the requirement that the experimental magnetic susceptibility equals that of the strictly one-dimensional model at the critical point leading to $J = 130\text{K}$ ($\alpha = 0$), $J = 150\text{K}$ ($\alpha \approx \alpha_c \approx 0.2412$ [1,22,23]), $J = 160\text{K}$ ($\alpha = 0.35$), and $J = 350\text{K}$ (free fermions). The circles in Fig. 1 denote

¹We have used 24 states in the renormalization step and have set the accuracy with respect to Trotter decomposition to $(TM)^{-1} = 0.05$.

the values of $x \cdot A_\alpha(x)$ at the experimentally determined spin-Peierls temperature $T_{\text{SP}}^0 = 14.4\text{K}$ for CuGeO_3 (see e.g. [24–26]). These values imply constants (cf. Eq. 4) $\mathcal{C}_0 \approx 0.019\text{K}^{-1}$ ($\alpha = 0$), $\mathcal{C}_{0.241} \approx 0.040\text{K}^{-1}$ ($\alpha = 0.241$), $\mathcal{C}_{0.35} \approx 0.070\text{K}^{-1}$ ($\alpha = 0.35$), and $\mathcal{C}_{\text{ff}} \approx 0.0027\text{K}^{-1}$ (free fermions).

For the unfrustrated model the value $x_{\text{SP}} \cdot A_{\alpha=0}(x_{\text{SP}})$ almost coincides with the results of CF. However the agreement happens only fortuitously since χ_0 is a zero temperature quantity. For other frustration parameters qualitative and quantitative deviations from the CF-line appear. The divergence of $A(x)$ for free fermions, Heisenberg model with $\alpha < \alpha_c$, $\alpha = \alpha_c$, and $\alpha > \alpha_c$, is $\log(x)$, $1/x \times \log$. corrections [4,27], $1/x$ (see [22] and references therein), and exponential [4], respectively. The quantitative values for $x_{\text{SP}} \cdot A(x_{\text{SP}})$ also differ from each other. We must conclude that it is risky to deduce quantitative results from the bosonization approach as already pointed out by CF. For applications to CuGeO_3 it is furthermore uncertain if the pure Heisenberg chain can even yield the qualitative results correctly. From the analysis of the susceptibility data at higher temperature we are led to favour the frustration parameter $\alpha = 0.35$ [2–4] for which the behaviour of the response function deviates considerably from χ_0/x .

IV. PRESSURE DEPENDENCE OF THE MAGNETIC SYSTEM

From (5) it is straight forward to deduce the dependence of the spin-Peierls temperature on the variation of the spin coupling constants. The relation is valid for every fixed value of α . We first focus on the system with $\alpha_0 = 0.35$ at ambient pressure.

The dependence of J and α on the external hydrostatic pressure have already been obtained from the relation between magnetostriction and the pressure dependence of the magnetic susceptibility χ [3]. From our TMRG data we find $\frac{\partial \chi}{\partial \ln J} \gg \frac{\partial \chi}{\partial \ln \alpha}$ at constant temperature. These two quantities appear in the following expression, $\frac{d\chi}{dp} = \sum_i \frac{\partial \chi}{\partial \ln x_i} \frac{\partial \ln x_i}{\partial p}$, where $x_1 = J$ and $x_2 = \alpha$. Therefore, we conclude the value for the pressure dependence of J to be more reliable than that of α . The authors of Ref. [3] deduced a value of $\frac{d \ln J}{dp} = -7.0(5)\%/GPa$ using the relation between magnetostriction and pressure dependencies of J and α . Using in addition the experimentally determined value for the pressure dependence of the spin-Peierls temperature $\frac{dT_{\text{SP}}}{dp} \approx 4.8\text{K}/GPa$ [18] we obtain the relation $T_{\text{SP}}(J)$ (thick black line in Fig. 2).

Due to the particular geometry involved in the super-exchange mechanism we expect the magnetic exchange energies to respond much more sensitively to the pressure (bond-bending mechanism [28,29]) than the phonon frequencies or the spin-phonon coupling constants. We therefore consider \mathcal{C} as independent of pressure which

applicability for CuGeO_3 will break down at higher pressure. The numerical results with $\mathcal{C} = \mathcal{C}_{0.35}$ are shown in Fig. 2. Obviously, a constant value of $\alpha = 0.35$ (dashed-dotted line) can not explain the observed behaviour, even yielding a change of the critical temperature to opposite direction.




FIG. 2. Variation of the spin-Peierls temperature versus the relative change of the magnetic exchange coupling. The lines show TMRG results for constant α and fixed $\mathcal{C}_{0.35}$. The thick black line displays the experimental behaviour as an implicit function of pressure. The dashed (dashed-dotted) line shows the theoretical results for $dJ'/dp = 0$ ($d\alpha/dp = 0$). The corresponding dependencies of α on the pressure are shown in the inset.

From geometrical reasons we expect J' to be non-decreasing. Assuming $J' = \text{constant}$ to be realized, the increase of T_{SP} is too small by a factor of approximately 4 (as displayed by the thick dashed line in Fig. 2). Consequently, the main effect must be a strong increase of J' . The deduced dependence of α on the hydrostatic pressure fits well with $\frac{d \ln \alpha}{dp} = 24 \pm 2\%/GPa$ (or $\frac{d \ln J'}{dp} = 17 \pm 2\%/GPa$ respectively) up to about 1.4 GPa as displayed in the inset of Fig. 2. The error was determined assuming that the value of $\frac{dT_{\text{SP}}}{dp}$ involves an error of $\pm 0.5\text{K}/GPa$, where the uncertainty in $\frac{d \ln J}{dp}$ only plays a secondary role. In general, already the weak requirement that J is non-increasing with hydrostatic pressure gives a minimum pressure dependence of $\frac{d \ln \alpha}{dp} \geq 20\%/GPa$. The pressure dependence of α has already been investigated in Ref. [3], however, with large error bounds which are respected by our results. The divergence of $\frac{d \ln \alpha}{dp}$ at a finite pressure or in other words, an upper limit of the critical temperature, is a physical prediction of the chosen one-dimensional approach. It is mostly based on a limited spontaneous gap as a function of α . But as already mentioned above, we expect an agreement with CuGeO_3 only in the low pressure region.

A similar analysis can be done on the basis of the unfrustrated Heisenberg model and free fermions. Evaluating the magnetic susceptibility data of Takahashi et al. [18] we derive $\frac{d \ln J}{dp} \approx -5\%/GPa$ ($\alpha = 0$) and $\frac{d \ln J}{dp} \approx -6\%/GPa$ (free fermions). In contrast to the frustrated case, a change of α under pressure for these initially unfrustrated models is not reasonable. The deduced theoretical BCS-ratio is too small by a factor of about 3 for free fermions (or equivalently Hartree-Fock calculations), in the case of the unfrustrated Heisenberg model even by a factor of ≈ 15 (see Tab. I).

We must conclude that neither the unfrustrated Heisenberg model nor Hartree-Fock results are able to describe the physics of $CuGeO_3$ correctly. In contrast to this a Heisenberg model with parameters $\alpha_0 = 0.35$ ans $\frac{d \ln \alpha}{dp} = 24\%/GPa$ turns out to reproduce the experimental findings up to about 1.4 GPa.

V. "BCS-RATIO"

Another interesting quantity is the ratio of the singlet-triplet gap to the spin-Peierls temperature, known as "BCS-ratio". Combining the TMRG data for A with zero temperature DRMG calculations for the singlet-triplet gap as a function of α and the dimerization one gets a relation between Δ_{ST} and T_{SP} . The results are shown in the bottom section of Fig. 3.




FIG. 3. Bottom: Singlet-triplet gap Δ_{ST} as a function of the spin-Peierls temperature. Symbols denote the DRMG results, the solid line shows the exact result for free fermions. The thin lines show the expected behaviour from the scaling of A (see text). Top left: Ratio of the spin gap and spin-Peierls temperature (BCS ratio). Top right: BCS ratio, experiment versus theory.

Using the scaling of A [4], the definition of the critical temperature and the dependence of the ground state

energy on small saturation dimerizations $\delta_0 = \delta(T = 0)$ one finds

$$\delta_0 \simeq T_{SP}^{3/2} \exp(-\frac{\Delta_{ST}^0}{2}/T_{SP}), \quad (6)$$

ignoring logarithmic corrections for $\alpha < \alpha_c$. Here Δ_{ST}^0 denotes the singlet-triplet gap for vanishing dimerization, which is zero for $\alpha \leq \alpha_c$. In the next step we apply $\Delta_{ST} - \Delta_{ST}^0 \simeq \delta_0^{2/3}$ [30,31] to derive an explicit asymptotic relation between the gap and the spin-Peierls temperature,

$$\Delta_{ST} - \Delta_{ST}^0 \simeq T_{SP} \exp(-\frac{\Delta_{ST}^0}{3}/T_{SP}), \quad (7)$$

again neglecting logarithmic corrections for $\alpha < \alpha_c$. The thin lines in Fig. 3 show fits according to Eq. (7) in the range up to $T/J = 0.15$. For $\alpha = 0.5$ we used the known value $\Delta_{ST}^0 = 0.2338J$ [32,33], hence only performed a one-parameter fit in the presented region. For $\alpha = 0.35$ a spontaneous gap of $\Delta_{ST}^0 = 0.035J$ is used, also derived by $T = 0$ DRMG. For $\alpha = 0$ the predicted linear behaviour (cf. (7)) can not be seen due to the logarithmic corrections. A linear extrapolation of our data points shows a positive offset. However, this is consistent with the presence of logarithmic corrections since we expect a zero limit at $T = 0$ with infinite slope. The free fermions show the well known behaviour $\Delta_{ST}/T_{SP} \approx 1.76$. We now derive the BCS-ratio as a function of the spin-Peierls temperature simply by dividing by T_{SP} (upper left of Fig. 3). Firstly we observe that the Heisenberg like models have a distinctly larger BCS-ratio than free fermions.

For comparison to experiment we again fix the constant \mathcal{C} . Furthermore, the pressure dependence of J and the frustration derived in section IV is taken into account here. The results are shown in the top right of Fig. 3. Interestingly, the experimental result [18] compares well with the low temperature asymptotics for free fermions but showing a pronounced increase with pressure which is not reproduced. From the investigations in Ref. [4] we already know for Heisenberg models that the gap and therefore the BCS-ratio is larger than the experimentally observed one. This happens due to the strictly one-dimensional treatment of $CuGeO_3$, i.e. the neglect of the dispersion perpendicular to the chain which will lower the true gap. The unfrustrated chain even yields a qualitatively incorrect tendency of a decreasing BCS-ratio under pressure the appropriately frustrated system is at least able to explain the increase.

VI. CONCLUSION

The TMRG analysis of the spin-Peierls phase transition allows a complete treatment of the quantum dynamics. In contrast to continuum descriptions correlations

are respected at all length scales, which leads to the exact response functions. Within the scope of an adiabatic description, equivalent to RPA as in the CF theory, we are able to study the influence of the spin-spin exchange energy scale J on the critical temperature. We like to emphasize that these results are a non-trivial improvement over the CF theory which shows no dependence on J at all. Moreover, frustrated models can be investigated as well.

Using the pressure dependence of J it is possible to study the dependence of the spin-Peierls temperature on pressure. Neither Hartree-Fock calculations nor the unfrustrated Heisenberg chain yield the strong increase as measured. Once again, our investigations favour a frustration of $\alpha = 0.35$ for CuGeO_3 at ambient pressure. We find a rather strong dependence of the frustration on pressure, $\frac{d \ln \alpha}{dp} = 24 \pm 2\%/GPa$, which agrees with earlier studies [3].

The analysis of the “BCS-ratio” also gives a clear indication that frustration is present in CuGeO_3 , even though some quantitative deviations can only be explained by residual perpendicular couplings. There is evidence for a strong dependence of α on pressure from the “BCS-ratio”.

Acknowledgement

We like to thank U. Löw for carefully reading the manuscript. The work was supported by the Deutsche Forschungsgemeinschaft through SFB 341.

[1] G. Castilla, S. Chakravarty and V. J. Emery, Phys. Rev. Lett. **75**, 1823 (1995)
[2] J. Riera and A. Dobry, Phys. Rev. B **51**, 16098 (1995)
[3] K. Fabricius et al., Phys. Rev. B **57**, 1102 (1998)
[4] A. Klümper, R. Raupach, and F. Schönfeld, Phys. Rev. B **59**, 3612 (1998)
[5] G. Bouzerar, A. P. Kampf and F. Schönfeld, cond-mat/9701176
[6] T. Lorenz *et al.*, Phys. Rev. Lett. **81**, 148 (1998)
[7] G.S. Uhrig, F. Schönfeld, and J. Boucher, Europhys. Lett. **41**, 431 (1998)
[8] G.S. Uhrig, F. Schönfeld, J.-P. Boucher, and M. Horvatic, Phys. Rev. B in press
[9] R.J. Bursill, T. Xiang and G.A. Gehring, J. Phys. Cond. Mat **8**, L583 (1996)
[10] X. Wang and T. Xiang, Phys. Rev. B **56**, 5061 (1997)
[11] N. Shibata, J. Phys. Soc. Jap. **66**, 2221 (1997)
[12] M. Suzuki and M. Inoue, Prog. Theor. Phys. **78**, 787 (1987)
[13] K. Maisinger, U. Schollwöck, Phys. Rev. Lett **81**, 445 (1998)
[14] N. Shibata, B. Ammon, M. Troyer, M. Sigrist, and K. Ueda, Phys. Soc. Jap. **67**, 1086 (1998)

[15] D. Coombes, T. Xiang, and G.A. Gehring, J. Phys. C **10**, L159 (1998)
[16] T. Xiang, Phys. Rev. B **58**, 9142 (1998)
[17] S. Eggert, and S. Rommer, Phys. Rev. Lett. **81**, 1690 (1998)
[18] H. Takahashi, N. Mōri, O. Fujita, J. Akimitsu and T. Matsumoto, Sol. State Commun. B **95**, 817 (1995)
[19] P.H.M. van Loosdrecht *et al.*, Phys. Rev. Lett. **78**, 487 (1997)
[20] R. Werner, C. Gros, and M. Braden, cond-mat/9810038
[21] M.C. Cross and D.S. Fisher, Phys. Rev. B **19**, 402 (1979)
[22] S. Eggert, Phys. Rev. B **54**, R9612 (1996)
[23] K. Okamoto and K. Nomura, Phys. Lett. A **169**, 433 (1993)
[24] K. Hirota *et al.*, Phys. Rev. Lett. **73**, 736 (1994)
[25] T. Lorenz *et al.*, Phys. Rev. B **55**, 5914 (1997)
[26] J.P. Pouget *et al.*, Phys. Rev. Lett. **72**, 4037 (1994)
[27] A. Klümper, to be published elsewhere
[28] W. Geertsma and D. Khomskii, Phys. Rev. B **54**, 3011 (1996)
[29] D. Khomskii, W. Geertsma, and M. Mostovoy, Czech. Journ. Phys. **46**, 3239 (1996)
[30] I. Affleck, D. Gepner, H.J. Schulz, and T. Ziman, J. Phys. A **22**, 511 (1989)
[31] G.S. Uhrig, F. Schönfeld, M. Laukamp, and E. Dagotto, Eur. Phys. J. B **7**, 67 (1999)
[32] W. J. Caspers, K. M. Emmett and W. Magnus, J. Phys. A **17**, 2687 (1984)
[33] G.S. Uhrig and E. Müller-Hartmann, unpublished

	$\frac{d \ln J}{dp} [\%/GPa]$	$\frac{d \ln \alpha}{dp} [\%/GPa]$	$\frac{dT_{SP}}{dp} [K/GPa]$
free fermions	≈ -6	(0)	$\approx \mathbf{1.8}$
$\alpha = 0$	≈ -5	(0)	$\approx \mathbf{0.3}$
$\alpha_0 = 0.35$	-7.0 ± 0.4	$\mathbf{24 \pm 2}$	4.8 ± 0.5

TABLE I. Pressure dependences of the spin system parameters and the spin-Peierls temperature as explained in the text. Numbers in bold face denote results obtained by our DMRG calculations, all other quantities were used as input data. Note that models based on free fermions or unfrustrated Heisenberg chains can not explain the rather large pressure dependence of the critical temperature.