

EQUIVARIANT EMBEDDINGS OF TREES INTO HYPERBOLIC SPACES

MARC BURGER, ALESSANDRA IOZZI, AND NICOLAS MONOD

1. INTRODUCTION

For every cardinal $\alpha \geq 2$ there are three complete constant curvature model manifolds of Hilbert dimension α : the sphere \mathbf{S}^α , the Euclidean space \mathbf{E}^α and the hyperbolic space \mathbf{H}^α . Studying isometric actions on these spaces corresponds in the first case to studying orthogonal representations and in the second case to studying cohomology in degree one with orthogonal representations as coefficients. In this paper we address the third case and, in particular, we study isometric actions of automorphisms groups of trees on \mathbf{H}^α .

The goal of this paper is twofold: first we exhibit, for every tree \mathcal{T} , a one-parameter family of equivariant embeddings (with respect to appropriate representations of $\text{Aut}(\mathcal{T})$) into an infinite dimensional hyperbolic space which are, up to rescaling, asymptotically isometric and have convex cobounded image. Secondly, in the case in which the tree is regular of finite valence at least 3 and $G < \text{Aut}(\mathcal{T})$ is a closed subgroup satisfying appropriate transitivity properties, we show that the representations constructed above give the unique irreducible nonelementary actions of G by isometries on a hyperbolic space of appropriate infinite dimension.

Quadratic forms of finite index, and in particular of index one, can be studied on real vector spaces of arbitrary dimension. A quadratic form of index one leads, *via* its cone of negative vectors, to a geodesic CAT(-1) space which is then complete if and only if the quadratic form satisfies a strong nondegeneracy condition. For any dimension α , there is one such space \mathbf{H}^α with ideal boundary $\partial\mathbf{H}^\alpha$ and bordification $\overline{\mathbf{H}}^\alpha = \mathbf{H}^\alpha \cup \partial\mathbf{H}^\alpha$. Then we have:

THEOREM A. *Let V be the set of vertices of a tree \mathcal{T} with $|V| = \alpha + 1$. Then for every $\lambda > 1$ there is an embedding $\Psi_\lambda : V \rightarrow \mathbf{H}^\alpha$ and a representation $\pi_\lambda : \text{Aut}(\mathcal{T}) \rightarrow \text{Isom}(\mathbf{H}^\alpha)$ such that:*

- (i) *The map Ψ_λ is π_λ -equivariant and extends equivariantly to a boundary map $\partial\Psi_\lambda : \partial\mathcal{T} \rightarrow \partial\mathbf{H}^\alpha$ which is a homeomorphism onto its image.*

N.M. partially supported by FNS grant 8220-067641 and NSF grant DMS 0204601.

- (ii) For any two vertices $x, y \in V$ there is a precise relation between the combinatorial distance $d_{\mathcal{T}}$ and the Riemannian distance $d_{\mathbf{H}^\alpha}$:

$$\lambda^{d_{\mathcal{T}}(x,y)} = \cosh d_{\mathbf{H}^\alpha}(\Psi_\lambda x, \Psi_\lambda y).$$

- (iii) The set $\Psi_\lambda(V)$ has finite codiameter in the convex hull $\mathcal{C} \subseteq \mathbf{H}^\alpha$ of the image of $\partial\Psi_\lambda$.

This result is the outcome of our attempt to understand certain claims of Gromov [G, Section 6.A], to the extent that nontrivial amalgams admit actions with unbounded orbits on infinite dimensional hyperbolic spaces.

Theorem A applies in particular to the automorphism group $\text{Aut}(\mathcal{T}_r)$ of an r -regular tree \mathcal{T}_r . By taking products and by denoting by \mathbf{H}^∞ the real hyperbolic space of countable dimension $\alpha = \aleph_0$, we obtain a family of metrically proper convex cobounded actions of $\text{Aut}(\mathcal{T}_r \times \mathcal{T}_s)$ on $\mathbf{H}^\infty \times \mathbf{H}^\infty$. This leads immediately to the following:

COROLLARY B. *Any cocompact lattice $\Gamma < \text{Aut}(\mathcal{T}_r \times \mathcal{T}_s)$ admits a metrically proper convex cobounded action on the product $\mathbf{H}^\infty \times \mathbf{H}^\infty$ of two hyperbolic spaces of countable dimension.*

Recall that in [BM1] and [BM2] these type of lattices were studied systematically and examples of torsion-free simple groups Γ were obtained. These Γ 's are then fundamental groups of finite aspherical (two-dimensional) complexes; on the other hand, the question to which extent there are compact aspherical manifolds (with or without boundary) with simple fundamental group is open. Here we obtain metrically proper convex cobounded actions of Γ on $\mathbf{H}^\infty \times \mathbf{H}^\infty$; in particular, $\Gamma \backslash (\mathbf{H}^\infty \times \mathbf{H}^\infty)$ retracts to a convex bounded (infinite dimensional) aspherical manifold with boundary. In contrast with the algebraic aspect of this situation, observe that if Λ is a group acting in a metrically proper convex cobounded way on \mathbf{H}^∞ , then Λ is a nonelementary Gromov hyperbolic group, and hence SQ-universal [O]; in particular, it admits many normal subgroups.

Turning to the classification of isometric actions, we recall that an action of a group G on \mathbf{H}^α by isometries is *elementary* if it preserves a point in $\overline{\mathbf{H}^\alpha}$ or a geodesic. Hence the study of elementary actions on \mathbf{H}^α reduces essentially to the study of isometric actions on the Euclidean space $\mathbf{E}^{\alpha-1}$ or on the sphere $\mathbf{S}^{\alpha-1}$. (Conversely, as observed by Gromov [G, 7.A], any isometric action on $\mathbf{E}^{\alpha-1}$ can be extended to an *elementary* action on \mathbf{H}^α by realizing $\mathbf{E}^{\alpha-1}$ as a horosphere based at a fixed point at infinity; similarly, isometric actions on $\mathbf{S}^{\alpha-1}$ extend obviously to actions with a fixed point in \mathbf{H}^α .)

Since any nonelementary action admits a unique minimal G -invariant hyperbolic subspace (Proposition 4.3), we say that a (nonelementary) action $G \rightarrow \text{Isom}(\mathbf{H}^\alpha)$ is *irreducible* if there is no G -invariant hyperbolic subspace other than \mathbf{H}^α .

Let now \mathcal{T}_r be the regular tree of finite valence $r \geq 3$, $G < \text{Aut}(\mathcal{T}_r)$ a closed subgroup and $\pi : G \rightarrow \text{Isom}(\mathbf{H}^\alpha)$ a nonelementary action of G on \mathbf{H}^α . If G acts triply transitively on $\partial\mathcal{T}_r$, we shall see that the image $\pi(g)$ of any hyperbolic automorphism $g \in G$ is a hyperbolic isometry of \mathbf{H}^α of translation length ℓ_π independent of g , provided g is of translation length one in \mathcal{T}_r . With this in mind, we can state the following:

THEOREM C. *Let $G < \text{Aut}(\mathcal{T}_r)$ be a closed subgroup which acts triply transitively on $\partial\mathcal{T}_r$. For every $\ell > 0$ there exists, up to equivalence, a unique irreducible nonelementary continuous homomorphism $\pi : G \rightarrow \text{Isom}(\mathbf{H}^\infty)$ with $\ell_\pi = \ell$.*

It follows that the representation π in Theorem C is exactly the irreducible component of $\pi_\lambda|_G$ for $\lambda = e^{\ell\pi}$ in Theorem A.

The structure and unitary representation theory of closed subgroups of $\text{Aut}(\mathcal{T}_r)$ with some transitivity conditions on their action at infinity is the object of intensive study. We refer to [FN], [A], and the references therein for a more comprehensive picture. A first notable set of examples to which Theorem C applies is given by the topological group $G = \mathbf{PGL}_2(k)$, where k is a non-Archimedean local field; indeed, if q is the cardinality of the residue field of k , then the action of $\mathbf{PGL}_2(k)$ on the associated Bruhat–Tits tree \mathcal{T}_{q+1} identifies it with a closed subgroup of $\text{Aut}(\mathcal{T}_{q+1})$ which acts triply transitively on $\partial\mathcal{T}_{q+1}$.

Another important class of examples of closed subgroups of $\text{Aut}(\mathcal{T}_r)$ are the universal groups introduced in [BM1]. Recall that when $\mathcal{T}_r = (X, Y)$ is a r -regular tree, one can label its edges in such a way that for every vertex the edges issued from it are labelled $\{1, 2, \dots, r\}$. Thus, for any $g \in \text{Aut}(\mathcal{T}_r)$ and vertex $x \in X$, one obtains a permutation $c(g, x) \in S_r$ representing g “locally” at x . To a permutation group $F < S_r$ one can then associate $U(F)$, the closed subgroup of $\text{Aut}(\mathcal{T}_r)$ consisting of all $g \in \text{Aut}(\mathcal{T}_r)$, such that $c(g, x) \in F$ for all $x \in X$. Then $U(F)$ does not depend, up to conjugation, on the labelling of the edges. It acts transitively on X and at every vertex it induces the full permutation group F on the edges issued from x and is, by construction, maximal with respect to this property. The group $U(F)$ satisfies Tits’ independence condition [T] and, in fact, all closed vertex transitive subgroup of $\text{Aut}(\mathcal{T}_r)$ satisfying Tits’ independence condition are of the form $U(F)$.

Many properties of $U(F)$ can be read off the finite permutation group $F < S_r$. For example, for $n = 2$ and 3 , $U(F)$ is n -transitive on $\partial\mathcal{T}_r$ if and only if F is n -transitive. In the case in which F is doubly transitive, the unitary dual of $U(F)$ has been determined by O. Amann [A]. When F is triply transitive, the above Theorem C applies to $U(F)$.

REMARK. We have been informed by A. Valette that the algebraic part of the construction of Theorem A can also be derived elegantly from the “tree cocycles” that he proposes in [V].

The structure of the paper is as follows. In Section 2, modelling on the finite dimensional case, we discuss basic properties of quadratic forms of finite index on a real vector space of arbitrary dimension, we single out the notion of strongly nondegenerate form and show that strongly nondegenerate forms are determined by their signature. In Section 3 we associate a hyperbolic space to every nondegenerate quadratic form of index one; this is a geodesic CAT(-1) space which is complete if and only if the form is strongly nondegenerate. This leads to the existence and the uniqueness of \mathbf{H}^α for every cardinal α . In Section 4 we discuss the existence of irreducible hyperbolic subspaces (Proposition 4.3) and establish the description of elementary actions in terms of orthogonal representations and cocycles in degree one (Proposition 4.4). In Section 5, 6 and 7 we turn more specifically to the study of actions on \mathbf{H}^α of automorphisms groups of trees. In Sections 5 and 6 we study more closely actions on \mathbf{H}^α of certain locally compact groups occurring as stabilizers of ends of trees, *i.e.* topological ascending HNN-extensions. These actions turn out to be elementary and hence substantial use of Section 4 is made. In Section 7 the proof of Theorem C is completed by showing that the irreducible part of the action under consideration is determined by its restriction to any parabolic subgroup. In Section 8 we give the explicit construction in Theorem A. Finally, the Appendix contains some of the explicit matrix representations used throughout the paper.

2. QUADRATIC FORMS OF FINITE INDEX

A *quadratic space* is a pair (\mathcal{H}, Q) consisting of a real vector space \mathcal{H} and a quadratic form $Q : \mathcal{H} \rightarrow \mathbf{R}$. As usual, Q is *positive definite* if $Q(x) > 0$ for all $x \neq 0$ and *negative definite* if $-Q$ is positive definite; $\dim \mathcal{H}$ denotes the cardinal of any \mathbf{R} -basis of \mathcal{H} . Define

$$i_{\pm}(Q) = \sup \left\{ \dim W : W \text{ is a subspace of } \mathcal{H} \text{ and } Q|_W \text{ is pos./neg. definite} \right\}$$

and the *index* of Q as

$$i(Q) = \sup \left\{ \dim W : W \text{ is an isotropic subspace of } \mathcal{H} \right\}.$$

Let $B : \mathcal{H} \times \mathcal{H} \rightarrow \mathbf{R}$ be the bilinear (symmetric) form associated to Q . For any subset $S \subseteq \mathcal{H}$ write

$${}^{\perp}S = \{x \in \mathcal{H} : B(x, s) = 0 \forall s \in S\}.$$

We say that Q is *nondegenerate* if ${}^{\perp}\mathcal{H} = 0$ and that the quadratic space is of *finite index* if $i(Q) \in \mathbf{N}$ (we agree that $0 \in \mathbf{N}$).

Just like in the case of finite dimensional quadratic spaces, we have:

PROPOSITION 2.1. *Let (\mathcal{H}, Q) be a nondegenerate quadratic space of finite index. Then*

$$(i) \quad i(Q) = \min\{i_-(Q), i_+(Q)\}.$$

Assume now $i(Q) = i_-(Q)$.

- (ii) If $W_- \subseteq \mathcal{H}$ is a negative definite subspace with $\dim W_- = i(Q)$, then $W_+ := {}^\perp W_-$ is positive definite and $\mathcal{H} = W_- \oplus W_+$.
- (iii) If $\mathcal{H} = W'_- \oplus W'_+$ is an orthogonal direct sum with W'_\pm pos./neg. definite, then $\dim W'_- = i(Q)$.

We precede the proof of the proposition by a couple of lemmas.

LEMMA 2.2. Let (\mathcal{H}, Q) be a quadratic space and $W \subseteq \mathcal{H}$ a finite dimensional subspace such that $Q|_W$ is nondegenerate. Then $\mathcal{H} = W \oplus {}^\perp W$. If moreover Q is nondegenerate then $Q|_{{}^\perp W}$ is so too.

Proof. The kernel of the linear map $\mathcal{H} \rightarrow W^*$ induced by B is ${}^\perp W$; since W has finite dimension we have

$$(2.1) \quad \dim(\mathcal{H}/{}^\perp W) \leq \dim W^* = \dim W.$$

On the other hand, since $W \cap {}^\perp W = 0$, the canonical projection $W \rightarrow \mathcal{H}/{}^\perp W$ is injective; hence it is an isomorphism by (2.1). \square

LEMMA 2.3. Let (\mathcal{H}, Q) be a quadratic space with $Q(x) \geq 0$ for all $x \in \mathcal{H}$. Then ${}^\perp \mathcal{H} = \{x \in \mathcal{H} : Q(x) = 0\}$.

Proof. If $Q(x) = 0$, then for all $y \in \mathcal{H}$ and all $\lambda \in \mathbf{R}$ we have

$$0 \leq Q(\lambda x + y) = 2\lambda B(x, y) + Q(y)$$

hence $B(x, y) = 0$ for all y . \square

Proof of Proposition 2.1. Let $A_\pm \subseteq \mathcal{H}$ be pos./neg. definite subspaces of finite dimension and set $A = A_- + A_+$. Then $i(Q|_A) \leq i(Q)$ and the theory of finite dimensional quadratic spaces implies

$$\min\{\dim A_-, \dim A_+\} \leq i(Q|_A) \leq i(Q),$$

whence $\min\{i_-(Q), i_+(Q)\} \leq i(Q)$. Assume without loss of generality that $i_-(Q) \leq i_+(Q)$, pick a negative definite subspace W_- of dimension $i_-(Q)$ and let $W_+ := {}^\perp W_-$. Since $Q|_{W_-}$ is nondegenerate, $\mathcal{H} = W_- \oplus W_+$ and $Q|_{W_+}$ is nondegenerate (Lemma 2.2). Since $\dim W_- = i_-(Q)$, we have $Q(x) \geq 0$ for all $x \in W_+$ and hence, by Lemma 2.3, W_+ is positive definite. If now W is an isotropic subspace with $\dim W = i(Q)$, then $W \cap W_+ = 0$ and thus the canonical projection $W \rightarrow \mathcal{H}/W_+ \cong W_-$ is injective; hence $i(Q) \leq i_-(Q)$. This proves (1) and (2). As for (3), if W_- is negative definite with $\dim W_- = i(Q)$, then $W_- \rightarrow \mathcal{H}/W'_+ \cong W'_-$ is injective and hence an isomorphism since $\dim W'_- \leq i_-(Q) = i(Q)$. \square

In view of Proposition 2.1 we call \pm -decomposition of (\mathcal{H}, Q) any orthogonal direct sum decomposition $\mathcal{H} = W_- \oplus W_+$ where W_\pm are pos./neg. definite. We

associate to such a decomposition the scalar product $\langle \cdot, \cdot \rangle_{\pm}$ defined for $x, y \in \mathcal{H}$ by

$$\langle x, y \rangle_{\pm} := B(x_+, y_+) - B(x_-, y_-)$$

where $x = x_- + x_+, y = y_- + y_+$ are the corresponding decompositions.

LEMMA 2.4. *Let (\mathcal{H}, Q) be a nondegenerate quadratic space of finite index and $\mathcal{H} = W_- \oplus W_+ = W'_- \oplus W'_+$ two \pm -decompositions. Then $(\mathcal{H}, \langle \cdot, \cdot \rangle_{\pm})$ is a Hilbert space if and only if $(\mathcal{H}, \langle \cdot, \cdot \rangle'_{\pm})$ is a Hilbert space, in which case the two scalar products are equivalent.*

We need the following:

LEMMA 2.5. *Let \mathcal{H} be a real vector space, $\langle \cdot, \cdot \rangle_1, \langle \cdot, \cdot \rangle_2$ two scalar products and $\mathcal{H}' \subseteq \mathcal{H}$ a subspace such that*

- (1) $\langle \cdot, \cdot \rangle_1, \langle \cdot, \cdot \rangle_2$ coincide on \mathcal{H}' and \mathcal{H}' is complete;
- (2) \mathcal{H}' is of finite codimension.

Then $\langle \cdot, \cdot \rangle_1, \langle \cdot, \cdot \rangle_2$ are equivalent and \mathcal{H} is a Hilbert space.

Proof. Let \mathcal{H}'_1 be the orthogonal of \mathcal{H}' for $\langle \cdot, \cdot \rangle_1$. Since \mathcal{H}' is complete, we have $\mathcal{H} = \mathcal{H}' \oplus \mathcal{H}'_1$. But \mathcal{H}'_1 is complete because it is finite dimensional and hence $(\mathcal{H}, \langle \cdot, \cdot \rangle_1)$ is a Hilbert space. For any $x \in \mathcal{H}$, write $x = x' + x'_1$ according to the above decomposition. Since $\|\cdot\|_1$ and $\|\cdot\|_2$ are equivalent on \mathcal{H}'_1 , there is $c > 0$ with $\|x'_1\|_1^2 \geq c\|x'_1\|_2^2$ for all x . We may chose $c \leq 1$ and now

$$\begin{aligned} \|x\|_1^2 &= \|x'\|_1^2 + \|x'_1\|_1^2 \geq \|x'\|_1^2 + c\|x'_1\|_2^2 \geq \\ &\geq c(\|x'\|_1^2 + \|x'_1\|_2^2) \geq \frac{c}{2}(\|x'\|_1 + \|x'_1\|_2)^2 \geq \frac{c}{2}\|x\|_2^2. \end{aligned}$$

□

Proof of Lemma 2.4. Assume $i(Q) = i_-(Q)$. Since B is continuous with respect to both $\|\cdot\|_{\pm}$ and $\|\cdot\|'_{\pm}$, all subspaces considered are closed for both topologies and so is in particular $W_+ \cap W'_+$. Moreover, the latter is of codimension at most $2i(Q)$, hence we conclude by Lemma 2.5. □

DEFINITION 2.6. A nondegenerate quadratic space of finite index (\mathcal{H}, Q) is *strongly nondegenerate* if for some (hence any) \pm -decomposition $\mathcal{H} = W_- \oplus W_+$ the space $(\mathcal{H}, \langle \cdot, \cdot \rangle_{\pm})$ is a Hilbert space.

We denote by $d(\mathcal{L})$ the cardinal of any Hilbert basis of a Hilbert space \mathcal{L} . Observing that $d(\mathcal{L})$ depends only on the equivalence class of the scalar product, we deduce that the pair $(d(W_+), d(W_-))$ is independent of the choice of a \pm -decomposition $\mathcal{H} = W_- \oplus W_+$. We call $(d(W_+), d(W_-))$ the *signature* of the strongly nondegenerate quadratic space (\mathcal{H}, Q) .

Two quadratic spaces (\mathcal{H}_1, Q_1) and (\mathcal{H}_2, Q_2) are *isomorphic* if there is a vector space isomorphism $T : \mathcal{H}_1 \rightarrow \mathcal{H}_2$ with $Q_1 = Q_2 \circ T$. Observe that if (\mathcal{H}_1, Q_1)

is nondegenerate of finite index, then so is (\mathcal{H}_2, Q_2) . Since the image of a \pm -decomposition for Q_1 is a \pm -decomposition for Q_2 , we see that Q_2 is strongly nondegenerate if Q_1 is so; in that case T is automatically continuous. In particular the orthogonal group $\mathbf{O}(Q)$ of a strongly nondegenerate form Q of finite index consists of bounded linear operators.

PROPOSITION 2.7. *For strongly nondegenerate forms of finite index, the signature is a complete invariant of isomorphisms.*

Proof. Using \pm -decompositions, this follows immediately from the fact that $d(\mathcal{L})$ determines completely the Hilbert spaces \mathcal{L} up to isomorphisms. \square

Let (\mathcal{H}, Q) be a strongly nondegenerate form of finite index, \mathcal{H}^* the topological dual and $A : \mathcal{H} \rightarrow \mathcal{H}^*$ the continuous morphism associated to B . Applying the Riesz representation theorem to the restrictions of B to W_\pm for a \pm -decomposition $\mathcal{H} = W_- \oplus W_+$, we deduce that A is an isomorphism (of topological vector spaces).

PROPOSITION 2.8. *Let (\mathcal{H}, Q) be a strongly nondegenerate form of finite index and $V \subseteq \mathcal{H}$ a closed subspace such that $Q|_V$ is nondegenerate. Then $(V, Q|_V)$ is strongly nondegenerate and $\mathcal{H} = V \oplus {}^\perp V$.*

Proof. Assume $i(Q) = i_-(Q)$ and let $V = U_- \oplus U_+$ be a \pm -decomposition of V with U_\pm pos./neg. definite (which exists by Proposition 2.1). Since V is closed and B continuous, $U_+ = {}^\perp U_- \cap V$ is closed. Let now $\mathcal{H} = W_- \oplus W_+$ be any \pm -decomposition of \mathcal{H} . Then $W_+ \cap U_+$ is closed, of finite codimension in V and B coincides with $\langle \cdot, \cdot \rangle_\pm$ on it. By Lemma 2.5 with $\langle \cdot, \cdot \rangle_1 = \langle \cdot, \cdot \rangle_2$, we deduce that V is a Hilbert space and hence $(V, Q|_V)$ is strongly nondegenerate. To conclude, the nondegeneracy of $(V, Q|_V)$ implies, as observed above, that the morphism $A_V : V \rightarrow V^*$ associated to $B|_V$ is a topological isomorphism. In particular, for every $x \in \mathcal{H}$ there is $x_V = A_V^{-1}B(x, \cdot)|_V \in V^*$ such that $B(x, y) = B(x_V, y)$ for all $y \in V$. Thus $x \in {}^\perp V + x_V$ and the claim follows. \square

3. REAL HYPERBOLIC SPACE

Let (\mathcal{H}, Q) be a nondegenerate quadratic space of index one; we assume more specifically that $i(Q) = i_-(Q) = 1$. Let $C_- := \{x \in \mathcal{H} : Q(x) < 0\}$ be the cone of negative vectors and $\mathbf{H} := \mathbf{R}^* \setminus C_-$ the set of negative lines. One proves as usual the reverse Cauchy–Schwartz inequality

$$B(x, y)^2 \geq Q(x)Q(y), \quad \forall x, y \in C_-$$

with equality if and only if $\mathbf{R}^*x = \mathbf{R}^*y$. This allows to define $\tilde{d} : C_- \times C_- \rightarrow \mathbf{R}_+$ by

$$\cosh^2 \tilde{d}(x, y) = \frac{B(x, y)^2}{Q(x)Q(y)},$$

which descends to a well defined function $d : \mathbf{H} \times \mathbf{H} \rightarrow \mathbf{R}_+$.

REMARK 3.1. A useful geometric fact is that for any finite set S of negative lines, the restriction $Q|_{\mathcal{H}_S}$ of Q to the span $\mathcal{H}_S \subseteq \mathcal{H}$ of S is equivalent to the standard real quadratic form of signature $(\dim \mathcal{H}_S - 1, 1)$ and that, under this isomorphism, the restriction of d to the image \mathbf{H}_S of \mathcal{H}_S in \mathbf{H} corresponds to the standard distance on the finite dimensional real hyperbolic space of dimension $|S| - 1$.

As a consequence of the above remark and [BH, Theorem 10.10], we have the following:

PROPOSITION 3.2. *The function d is a distance function with respect to which \mathbf{H} is a geodesic CAT(-1) space.* \square

Let ℓ_- be a vector of length -1 and let $\mathcal{H} = \mathcal{H}_+ \oplus \mathbf{R}\ell_-$ be the orthogonal decomposition, where $Q|_{\mathcal{H}_+}$ is positive definite (see Proposition 2.1). The exponential map $\exp : \mathcal{H}_+ \rightarrow \mathbf{H}$ is defined as follows. For every $v \in \mathcal{H}_+$ there is a unique $t > 0$ such that $x := v + t\ell_-$ has length -1 ; we define $\exp(v) := [x]$ to be the image of x in \mathbf{H} . A straightforward computation gives

$$\cosh d(\exp(v), \exp(w)) = \left| -B(v, w) + \sqrt{1 + Q(v)}\sqrt{1 + Q(w)} \right|,$$

and, in particular, $\cosh d(\exp(v), [\ell_-]) = \sqrt{1 + Q(v)}$.

PROPOSITION 3.3. *The CAT(-1) space \mathbf{H} is complete if and only if (\mathcal{H}, Q) is strongly nondegenerate.*

Proof. Using the above formulæ, one checks that \exp is for all $R > 0$ a bi-Lipschitz bijection between the ball in $(\mathcal{H}_+, Q|_{\mathcal{H}_+})$ of radius $(\sinh R)^2$ centered at 0 and the ball in (\mathbf{H}, d) of radius R centered at $[\ell_-]$. Thus (\mathbf{H}, d) is complete if and only if \mathcal{H}_+ is complete, which in view of Lemma 2.5 (with $\langle \cdot, \cdot \rangle_1 = \langle \cdot, \cdot \rangle_2$) is equivalent to the quadratic space (\mathcal{H}, Q) being strongly nondegenerate. \square

Observe that any orthogonal transformation $T \in \mathbf{O}(Q)$ preserves C_- and descends to an isometry of \mathbf{H} . The group $\mathbf{O}(Q)$ is a direct product $\mathbf{O}_+(Q) \cdot \{\pm Id\}$, where $\mathbf{O}_+(Q)$ is the subgroup preserving the (two) connected components of C_- . Let $\mathbf{PO}(Q) = \mathbf{O}(Q) / \pm Id$. Then:

PROPOSITION 3.4. *The homomorphism $\mathbf{O}(Q) \rightarrow \text{Isom}(\mathbf{H})$ induces isomorphisms*

$$\mathbf{O}_+(Q) \rightarrow \mathbf{PO}(Q) \rightarrow \text{Isom}(\mathbf{H}).$$

REMARK 3.5. Let G be a topological group and $\pi : G \rightarrow \mathbf{O}(Q)$ a group homomorphism. We call π *continuous* if the action map $G \times \mathcal{H} \rightarrow \mathcal{H}$ is continuous; then, the resulting action $G \times \mathbf{H} \rightarrow \mathbf{H}$ is continuous. Conversely, given a continuous action $G \times \mathbf{H} \rightarrow \mathbf{H}$, one verifies that the resulting homomorphism $G \rightarrow \mathbf{O}_+(Q)$ deduced from Proposition 3.4 is continuous.

LEMMA 3.6. *The $\mathbf{O}(Q)$ -action on \mathbf{H} is transitive.*

Proof of the lemma. Let $L, L' \subseteq \mathcal{H}$ be two negative lines. Then $L \oplus {}^\perp L$ and $L' \oplus {}^\perp L'$ are two \pm -decompositions, and ${}^\perp L, {}^\perp L'$ are isomorphic Hilbert spaces. Hence there is (compare also Proposition 2.7) an isomorphism of (\mathcal{H}, Q) bringing L to L' . \square

Proof of Proposition 3.4. Let $T \in \text{Isom}(\mathbf{H})$. By Lemma 3.6, we may assume that T fixes $[\ell_-]$. Define a map $U : \mathcal{H}_+ \rightarrow \mathcal{H}_+$ by $\exp(U(v)) = T(\exp(v))$. It follows from the above formulæ that U is a bijection, fixes 0 and preserves $B|_{\mathcal{H}_+}$. Hence U is a linear orthogonal transformation of \mathcal{H}_+ . Defining $S := U \oplus \text{Id}$, one verifies that $S \in \mathbf{O}_+(Q)$ corresponds to T via $\mathbf{O}(Q) \rightarrow \text{Isom}(\mathbf{H})$ and the statement follows. \square

One proves similarly:

PROPOSITION 3.7. *Let (\mathcal{H}_i, Q_i) be strongly nondegenerate quadratic spaces of signature $(\alpha_i, 1)$ and let \mathbf{H}_i be the associated hyperbolic spaces (for $i = 1, 2$). The following are equivalent:*

- (\mathcal{H}_1, Q_1) is isomorphic to (\mathcal{H}_2, Q_2) .
- $\alpha_1 = \alpha_2$.
- \mathbf{H}_1 is isometric to \mathbf{H}_2 . \square

Thus we obtain for each cardinal α “the” real hyperbolic space \mathbf{H}^α .

3.1. Bordification. Let again $\mathbf{H} = \mathbf{R}^* \setminus C_-$ be the real hyperbolic space associated to a strongly nondegenerate quadratic space (\mathcal{H}, Q) of signature $(\alpha, 1)$. Let $\partial\mathbf{H}$ be the boundary of the CAT(-1) space (\mathbf{H}, d) defined as usual as classes of asymptotic rays. Set

$$C_0 := \{x \in \mathcal{H} : Q(x) = 0, x \neq 0\}, \quad C_{\leq 0} := \{x \in \mathcal{H} : Q(x) \leq 0, x \neq 0\}.$$

Using that any configuration of finitely many geodesics in \mathbf{H} is contained in a finite dimensional hyperbolic subspace (see Remark 3.1), we obtain a bijection identifying the bordification $\overline{\mathbf{H}} = \mathbf{H} \sqcup \partial\mathbf{H}$ with the set $\mathbf{R}^* \setminus C_{\leq 0}$. We relate this to the description of $\overline{\mathbf{H}}$ in terms of Busemann cocycles: for every $x \in C_{\leq 0}$, define $\tilde{b}_x : C_- \times C_- \rightarrow \mathbf{R}$ by

$$(3.1) \quad \tilde{b}_x(y, z) := \begin{cases} \tilde{d}(x, y) - \tilde{d}(x, z) & \text{if } x \in C_-, \\ \frac{1}{2} \ln \frac{B(x, y)^2 Q(z)}{B(x, z)^2 Q(y)} & \text{if } x \in C_0. \end{cases}$$

Then, for every $x \in C_{\leq 0}$, \tilde{b}_x satisfies the cocycle identity

$$\tilde{b}_x(y_2, y_3) - \tilde{b}_x(y_1, y_3) + \tilde{b}_x(y_1, y_2) = 0.$$

Moreover, \tilde{b}_x gives a well defined function $b : \mathbf{R}^* \setminus C_{\leq 0} \times \mathbf{H} \times \mathbf{H} \rightarrow \mathbf{R}$ which coincides on $\mathbf{H} \times \mathbf{H} \times \mathbf{H}$ with $(x, y, z) \mapsto d(x, y) - d(x, z)$.

For every $x \in \mathbf{R}^* \setminus C_{\leq 0}$ and $z_1, z_2 \in \mathbf{H}$, the cocycle property implies that the continuous functions $y \mapsto b_x(y, z_1)$ and $y \mapsto b_x(y, z_2)$, defined on \mathbf{H} , differ by a constant. Thus we obtain, for every $x \in \mathbf{R}^* \setminus C_{\leq 0}$, a well defined class $B(x) \in C(\mathbf{H})/\mathbf{R}$,

where $C(\mathbf{H})$ is the space of continuous functions on \mathbf{H} . Endowing $C(\mathbf{H})/\mathbf{R}$ with the topology coming from the topology on $C(\mathbf{H})$ of uniform convergence on bounded sets, and denoting by $\overline{B(\mathbf{H})}$ the closure of $B(\mathbf{H})$ in $C(\mathbf{H})/\mathbf{R}$, we have the following:

PROPOSITION 3.8. *The map $B : \mathbf{R}^* \setminus C_{\leq 0} \rightarrow \overline{B(\mathbf{H})}$ is a homeomorphism when $\mathbf{R}^* \setminus C_{\leq 0}$ is endowed with the quotient of the norm topology.*

One can verify that in this topology $\overline{\mathbf{H}}$ is compact if and only if \mathbf{H} is finite dimensional.

By a slight abuse of terminology, we call *horospheres* in \mathcal{H} (respectively, in \mathbf{H}) centered at $x \in C_0$ (respectively, at $\xi \in \partial\mathbf{H}$) the level sets of $\tilde{b}_x(\cdot, z)$ (respectively, $B(\xi)$).

4. NONELEMENTARY AND ELEMENTARY ACTIONS

In this section we study basic properties of group actions on hyperbolic spaces. First we establish that any nonelementary action has a unique minimal invariant hyperbolic subspace, and then we turn to the description of elementary actions in terms of orthogonal representations, characters, and continuous cocycles.

Let X be a metric space. Recall that a *semicontraction* is a map $T : X \rightarrow X$ such that $d(Tx, Ty) \leq d(x, y)$ for all $x, y \in X$. Recall the

PROPOSITION 4.1. *Let X be a complete $CAT(-1)$ space and $T : X \rightarrow X$ a semicontraction. Then one of the following holds:*

- (i) *The set $\{T^n x : n \geq 1\}$ is bounded for some (hence any) $x \in X$ and the set $X^T \subseteq X$ of T -fixed points is not empty.*
- (ii) *The set $\{T^n x : n \geq 1\}$ is unbounded for some (hence any) $x \in X$ and there exists a subsequence $\{n_k\}_{k \geq 1}$ and $\xi \in \partial X$ with $\lim_{k \rightarrow \infty} T^{n_k} x = \xi$ and $T\xi = \xi$. Moreover, $|(\partial X)^T| = 1$ or 2 .*

(A general semicontraction need not extend to infinity; the notation $T\xi = \xi$ means that $T : X \rightarrow X$ extends by continuity to $X \cup \{\xi\}$ endowed with the topology induced by \overline{X} .)

Proof of the proposition. The case (ii) follows from the argument given by A. Karlsson (proof of [K, 5.1]); see [KN, §3] for the additional statement on $|(\partial X)^T|$. If on the other hand the T -orbits are bounded, then it is known that X^T is non-empty; indeed, one verifies that for any $x \in X$ the circumcentre of the set $\{T^k(x) : k \geq n\}$ converges to a T -fixed point as $n \rightarrow \infty$. \square

This result applies in particular to the case where T is an isometry and is the basis for the classification of isometries.

DEFINITION 4.2. An isometry T is called:

- *Elliptic* if $\{T^n x : n \geq 1\}$ is bounded.
- *Parabolic* if $\{T^n x : n \geq 1\}$ is unbounded and $|(\partial X)^T| = 1$.
- *Hyperbolic* if $\{T^n x : n \geq 1\}$ is unbounded and $|(\partial X)^T| = 2$.

If T is hyperbolic, then $(\partial X)^T = \{\xi_-, \xi_+\}$ with $\lim_{n \rightarrow \pm\infty} T^n x = \xi_{\pm}$ for all $x \in X$ (see again [KN, §3]). However, when X is not proper and T is parabolic, the sequence $T^n x$ might not converge in \overline{X} .

As usual a group action $G \times X \rightarrow X$ by isometries is called *elementary* if G preserves a nonempty finite subset of \overline{X} . This is equivalent to saying that either G fixes a point in \overline{X} or it preserves a geodesic.

Let now \mathbf{H} be the hyperbolic space associated to a strongly nondegenerate quadratic space (\mathcal{H}, Q) of signature $(\alpha, 1)$. In the sequel we shall study nonelementary and elementary actions and we shall prove the following

PROPOSITION 4.3. *Let $\pi : G \rightarrow \mathbf{O}(Q)$ be a homomorphism. Then, one of the following holds:*

- (i) G preserves an isotropic line and all horospheres in \mathcal{H} centered at it.
- (ii) G preserves a negative line.
- (iii) There is a unique minimal nondegenerate closed G -invariant subspace $\mathcal{H}_1 \subseteq \mathcal{H}$ of index one. Any nondegenerate closed G -invariant subspace of index one contains \mathcal{H}_1 .

First we prove the proposition in the case where the associated action on \mathbf{H} is nonelementary, which excludes of course (i) and (ii). The remaining will be a consequence of a closer analysis of elementary actions.

Proof in the nonelementary case. We need to show that (iii) holds. Let \mathcal{P} be the set of G -invariant closed positive definite subspaces of \mathcal{H} , ordered by inclusion, let $\mathcal{C} \subseteq \mathcal{P}$ be a maximal chain and $L := \overline{\bigcup \mathcal{C}}$. Then L is closed, G -invariant, and $Q|_L \geq 0$. By Lemma 2.3 applied to L , we have that $\{x \in L : Q(x) = 0\} = L \cap {}^\perp L$; if the latter were not zero, it would be a G -invariant isotropic line, contradicting the assumption that the action is nonelementary. Thus L is a G -invariant closed maximal positive definite subspace of \mathcal{H} . Set $\mathcal{H}_0 := L$, $\mathcal{H}_1 := {}^\perp L$. Then, by Proposition 2.8, $\mathcal{H} = \mathcal{H}_0 \oplus \mathcal{H}_1$ and \mathcal{H}_1 is a G -invariant closed nondegenerate subspace of index one which is minimal with respect to these properties.

Let now $\mathcal{H} = \mathcal{H}'_0 \oplus \mathcal{H}'_1$ be any other orthogonal decomposition into G -invariant closed subspaces where \mathcal{H}'_0 is positive definite and \mathcal{H}'_1 of index one. We need to show that $\mathcal{H}'_1 \supseteq \mathcal{H}_1$. Consider $J := \mathcal{H}_1 \cap \mathcal{H}'_1$. Again, since the G -action is not elementary, J is nondegenerate. There are two cases:

J is indefinite. Since $J \subseteq \mathcal{H}_1$, we have either $J = \mathcal{H}_1$, whence $\mathcal{H}'_1 \supseteq \mathcal{H}_1$ and we are done; or $J = 0$.

J is positive definite. Then $\mathcal{H}_0 \oplus J$ would be a G -invariant closed positive definite subspace and, by maximality, we would have $\mathcal{H}_0 \oplus J \subseteq \mathcal{H}_0$ and hence, once again, $J = 0$.

Thus, we may assume (for a contradiction) $\mathcal{H}_1 \cap \mathcal{H}'_1 = 0$. Let $\mathbf{H}_1, \mathbf{H}'_1 \subseteq \mathbf{H}$ be the corresponding hyperbolic subspaces and consider the orthogonal projections $p : \mathbf{H} \rightarrow \mathbf{H}_1$ and $p' : \mathbf{H} \rightarrow \mathbf{H}'_1$ given by the nearest point retraction. Since $\mathbf{H}_1 \cap \mathbf{H}'_1 = \emptyset$ and \mathbf{H} is CAT(-1), both $p|_{\mathbf{H}'_1}$ and $p'|_{\mathbf{H}_1}$ are contractions. Hence the map $f : \mathbf{H}_1 \rightarrow \mathbf{H}_1$, defined by $f := p|_{\mathbf{H}'_1} \circ p'|_{\mathbf{H}_1}$, is a G -equivariant contraction, that is, $d(f(x), f(y)) < d(x, y)$ for all distinct $x, y \in \mathbf{H}_1$. If for some $x \in \mathbf{H}_1$ the set $\{d(f^n(x), x) : n \geq 1\}$ were bounded, Proposition 4.1(i) would imply the existence of an f -fixed point in \mathbf{H} . Since f is G -equivariant, the set of its fixed points \mathbf{H}_1^f is G -invariant. However, since f is a contraction, \mathbf{H}_1^f consists of one point, which is hence G -fixed, contradicting the assumption that the G -action is nonelementary.

Thus, the f -orbits are unbounded and by Proposition 4.1(ii) there is a subsequence $\{n_k\}$ and $\xi \in \partial\mathbf{H}_1$ with $\lim_{k \rightarrow \infty} f^{n_k}(x) = \xi$. However, for every given $g \in G$ and for $x \in \mathbf{H}_1$

$$d(gf^n(x), f^n(x)) = d(f^n(gx), f^n(x)) < d(gx, x)$$

is bounded independently of n , thus, by passing to subsequences, the sequences $gf^n(x)$ and $f^n(x)$ are at bounded distance and hence define the same point at infinity, namely $g\xi = \xi$. Since this contradicts again the assumption that the action is nonelementary, the proof in this case is complete. \square

4.1. Elementary Actions. As before, let (\mathcal{H}, Q) be a strongly nondegenerate quadratic space of signature $(\alpha, 1)$. We shall study, for a topological group G , the elementary actions on \mathbf{H}^α , and more specifically, the actions fixing a point in $\partial\mathbf{H}^\alpha$. Thus, fix L_+ a isotropic line in \mathcal{H} , let $\mathbf{O}_{L_+}(Q)$ be its stabilizer in $\mathbf{O}(Q)$, and let $\text{Rep}(G, \mathbf{O}_{L_+}(Q))$ be the set of continuous representations (see Remark 3.5). Then G acts on L_+ by multiplication by a continuous character $\chi : G \rightarrow \mathbf{R}^*$. The bilinear form B induces a Hilbert space structure of Hilbert dimension $\alpha - 1$ on ${}^\perp L_+/L_+$ and we may thus fix a real Hilbert space E of Hilbert dimension $\alpha - 1$ and an isomorphism $i : {}^\perp L_+/L_+ \rightarrow E$. Since $\pi : G \rightarrow \mathbf{O}_{L_+}$ is continuous and G preserves L_+ and ${}^\perp L_+$, it induces on ${}^\perp L_+/L_+$ a continuous orthogonal representation which we transport to E via i , obtaining $\varrho : G \rightarrow \mathbf{O}(E)$. The space $\mathcal{H}/{}^\perp L_+$ is one-dimensional; since G preserves B , it acts on that space by multiplication by χ^{-1} .

We define a new G -module structure on \mathcal{H} by means of the continuous representation $\chi \otimes \pi$. Then we have a short exact sequence of G -modules

$$0 \longrightarrow {}^\perp L_+/L_+ \longrightarrow \mathcal{H}/L_+ \longrightarrow \mathcal{H}/{}^\perp L_+ \longrightarrow 0$$

in which the last term is a trivial G -module of dimension one. Thus, applying the corresponding transgression map in degree zero, the image of the trivial module $\mathcal{H}/{}^\perp L_+$ in the continuous cohomology $H_c^1(G, {}^\perp L_+/L_+)$ yields *via* i a subspace

$$\mathbf{R} \cdot \eta \subseteq H_c^1(G, \chi \otimes \varrho).$$

Thus, with i fixed, we associated to every continuous homomorphism $\pi : G \rightarrow \mathbf{O}_{L_+}(Q)$ the following data:

- a continuous homomorphism $\chi_\pi \in \text{Hom}_c(G, \mathbf{R}^*)$,
- a continuous orthogonal representation $\varrho_\pi \in \text{Hom}_c(G, \mathbf{O}(E))$, and
- a continuous class $\eta_\pi \in H_c^1(G, \chi \otimes \varrho)$, well defined up to scalar multiplication.

Denoting by $\mathcal{Z}(G, E)$ the set of all triples (χ, ϱ, η) with $\chi \in \text{Hom}_c(G, \mathbf{R}^*)$, $\varrho \in \text{Hom}_c(G, \mathbf{O}(E))$ and $\eta \in H_c^1(G, \chi \otimes \varrho)$, we have that $\mathbf{O}(E) \times \mathbf{R}^*$ acts on $\mathcal{Z}(G, E)$ by $(T, \lambda)(\chi, \varrho, \eta) = (\chi, T\varrho T^{-1}, \lambda T\eta)$.

PROPOSITION 4.4. (i) *The map $\text{Rep}(G, \mathbf{O}_{L_+}(Q)) \rightarrow \mathcal{Z}(G, E)$, $\pi \mapsto (\chi_\pi, \varrho_\pi, \eta_\pi)$ induces a bijection*

$$\text{Rep}(G, \mathbf{O}_{L_+}(Q))/\mathbf{O}_{L_+}(Q) \xrightarrow{\cong} [\mathbf{O}(E) \times \mathbf{R}^*] \setminus \mathcal{Z}(G, E).$$

(ii) *The representation π leaves all horospheres centered at L_+ invariant if and only if $|\chi_\pi| = 1$.*

Proof. Given $\chi : G \rightarrow \mathbf{R}^*$, $\varrho : G \rightarrow \mathbf{O}(E)$ and $\eta \in H_c^1(G, \chi \otimes \varrho)$, we indicate how to reconstruct $\pi \in \text{Rep}(G, \mathbf{O}_{L_+}(Q))$. Fix an isotropic line $L_- \neq L_+$ and let $f : G \rightarrow E$ be a continuous cocycle representing η . Set $F = {}^\perp(L_+ \oplus L_-)$, and denote by j the isomorphism of Hilbert spaces obtained by composing $F \rightarrow {}^\perp L_+ \rightarrow E$. Fix $\ell_\pm \in L_\pm$ with $B(\ell_-, \ell_+) = 1$. Using the notation of Appendix A, we define

$$\pi(g) = \begin{pmatrix} \pi(g)_1 & \pi(g)_2^+ \\ 0 & \pi(g)_3^- & \pi(g)_4 \end{pmatrix}$$

by

$$\pi(g)_1 = \begin{pmatrix} \chi(g) & a(g) \\ 0 & \chi(g)^{-1} \end{pmatrix}$$

where

$$(4.1) \quad \begin{aligned} a(g) &= -\frac{1}{2}\chi(g)\|f(g)\|_E^2 \\ \pi(g)_2^+(v) &= -\langle \varrho(g)j(v), f(g) \rangle_E, \quad \forall v \in F \\ \pi(g)_3^- &= \chi(g)^{-1}j^{-1}(f(g)) \\ \pi(g)_4 &= j^{-1}\varrho(g)j. \end{aligned}$$

The rest of the proposition is now a verification left to the reader and uses the fact that $|\chi_\pi|$ is the exponential of the Busemann character associated to the fixed point L_+ . \square

We turn now to representations $\pi : G \rightarrow \mathbf{O}_{L_+}(Q)$ for which $|\chi|$ is not identically 1 (write $\chi = \chi_\pi$). We fix once and for all χ and $a \in G$ such that $|\chi(a)| \neq 1$.

DEFINITION 4.5. We shall say that a continuous cocycle $f : G \rightarrow E$ for $\chi \otimes \varrho$ is *standard* if $f(a) = 0$.

LEMMA 4.6. Let $\varrho : G \rightarrow \mathbf{O}(E)$ be a continuous orthogonal representation and $\chi : G \rightarrow \mathbf{R}^\times$ a continuous homomorphism with $|\chi(a)| \neq 1$.

- (i) Every class in $H_c^1(G, \chi \otimes \varrho)$ admits a unique standard representative.
- (ii) If $M < G$ is a compact subgroup normalised by a , then any standard cocycle vanishes on M .

Proof. (i) Set $\tau = \chi \otimes \varrho$. Recall that $|\chi(a)| \neq 1$ implies that $1 - \tau(a)$ is invertible and hence $H^1(\langle a \rangle, \tau)$ vanishes. Therefore, for any cocycle $f' : G \rightarrow E$ there exists $v \in E$ such that $f'(a^n) = \tau(a^n)v - v$. Now $f(g) := f'(g) + v - \tau(g)v$ defines a standard cocycle. If f_1 and f_2 are any two cohomologous cocycles, there exists $v \in E$ such that $f_1(g) = f_2(g) + \tau(g)v - v$. If in addition f_1 and f_2 are standard, then $\tau(a)v = v$, which implies, since $|\chi(a)| \neq 1$, that $v = 0$ and hence $f_1 = f_2$.

(ii) Let f be a standard cocycle. Since M is compact, $C := \sup_{k \in M} \|f(g)\|$ is finite. We have for all $k \in M$

$$f(k) = \tau(k)f(a) + f(k) = f(ka) = f(aa^{-1}ka) = \tau(a)f(a^{-1}ka) + f(a) = \tau(a)f(a^{-1}ka),$$

which implies that $C = |\chi(a)|C$ and hence $C = 0$. \square

Let $\eta \in H_c^1(G, \chi \otimes \varrho)$ and let $\pi : G \rightarrow \mathbf{O}_{L_+}(Q)$ be the homomorphism associated to (χ, ϱ, η) by the above construction. Then $\pi(a)$ is hyperbolic with fixed points L_- and L_+ . If $f : G \rightarrow E$ is the standard cocycle representing the class η , define

$$I_\eta := \overline{\langle f(g) : g \in G \rangle},$$

which is a closed G -invariant subspace of E . With these definitions, we have the following:

PROPOSITION 4.7. *There is a G -invariant orthogonal decomposition*

$$\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_0,$$

where

$$\mathcal{H}_1 = (L_+ \oplus L_-) \oplus j^{-1}(I_\eta), \quad \mathcal{H}_0 = {}^\perp \mathcal{H}_1.$$

Moreover, the subspace \mathcal{H}_1 is nondegenerate of index one, \mathcal{H}_0 is positive definite and if $\mathcal{H}'_1 \subseteq \mathcal{H}$ is any closed nondegenerate G -invariant subspace of index one, then $\mathcal{H}'_1 \supset \mathcal{H}_1$.

Proof. We verify the last assertion: let $\mathbf{H}'_1 \subseteq \mathbf{H}_1$ be the hyperbolic subspace associated to \mathcal{H}'_1 . Without loss of generality, $|\chi(a)| > 1$. Then $\lim_{n \rightarrow \pm\infty} \pi(a^n)x = L_\pm$ for all $x \in \mathbf{H}'_1$, hence $L_\pm \in \partial\mathbf{H}'_1$. If $L := L_+ \oplus L_-$ and $\text{Pr}_L : \mathcal{H} \rightarrow L$ is the orthogonal projection, this implies that $\pi(g)\ell_- - \text{Pr}_L(\pi(g)\ell_-)$ is in \mathcal{H}'_1 for all $g \in G$; therefore, using the formulæ in (4.1) and Appendix A, $\mathcal{H}'_1 \supseteq j^{-1}(I_\eta)$. \square

End of proof of Proposition 4.3. We know already that the alternative (iii) of the proposition holds in the nonelementary case. If the G -action is elementary, then there are the following possibilities:

G fixes a point in \mathbf{H} . This corresponds to (ii).

G leaves a geodesic line in \mathbf{H} invariant. Denote this line by $\mathbf{H}_1 \subseteq \mathbf{H}$. This means that there is a G -invariant two-dimensional subspace $\mathcal{H}_1 \subseteq \mathcal{H}$ of index one. In fact, $\mathcal{H}_1 = L_+ \oplus L_-$, where $\partial\mathbf{H}_1 = \{L_-, L_+\}$. We may assume that G has no fixed point in \mathbf{H}_1 , so there is $g \in G$ acting hyperbolically: say, $\lim_{n \rightarrow \pm\infty} \pi(g)^n x = L_\pm$. But then, if \mathcal{H}'_1 is any G -invariant closed nondegenerate subspace of index one and $\mathbf{H}'_1 \subseteq \mathbf{H}$ the associated hyperbolic subspace, we get $\partial\mathbf{H}'_1 \ni L_\pm$, hence $\mathcal{H}'_1 \supseteq \mathcal{H}_1$.

G fixes an isotropic line L_+ . Let then $\chi : G \rightarrow \mathbf{R}^*$ be the associated character. Either $|\chi| = 1$ and we are in alternative (i), or $|\chi| \neq 1$ and we can apply Proposition 4.7, so that alternative (iii) holds once again. \square

5. ACTIONS OF CERTAIN HNN-EXTENSIONS

Let $P = \langle a \rangle \rtimes N$ be a locally compact group, semidirect product of an infinite cyclic subgroup $\langle a \rangle$ with generator a and a closed normal subgroup $N = \bigcup_{n \in \mathbf{Z}} K_n$ which is the increasing union of compact open subgroups $K_n < K_{n+1}$ such that $aK_n a^{-1} = K_{n+1}$ for all $n \in \mathbf{Z}$. In other words, P is the topological ascending HNN-extension $P = K_0 *_a$.

In this section we study more in detail elementary actions of P on a hyperbolic space \mathbf{H} , using the results in Section 4, in particular Proposition 4.4. We begin with the following general fact:

LEMMA 5.1. *Let X be a complete CAT(-1) space and $P \times X \rightarrow X$ a continuous action by isometries. Then there is a P -fixed point in \overline{X} and, in fact, one of the following holds:*

- (i) a is elliptic and $X^P \neq \emptyset$.
- (ii) a is parabolic and $|(\partial X)^P| = 1$.
- (iii) a is hyperbolic and the attracting fixed point of a is P -fixed.

Proof. For (i), we have for all $x \in X$

$$\sup_{g \in N} d(gx, x) = \sup_{k \in K_0, n \geq 0} d(ka^{-n}x, a^{-n}x).$$

Since by assumption a has bounded orbits and K_0 is compact, the latter quantity is bounded and hence N has bounded orbits; the setwise decomposition $P = \langle a \rangle \cdot N$ implies that P itself has bounded orbits and hence admits a fixed point in X .

In cases (ii) and (iii), the set $\{a^n x : n \geq 0\}$ is unbounded for all $x \in X$. Pick $\xi \in \partial X$ and a subsequence $\{n_k\}$ of \mathbf{N} with $\lim_{k \rightarrow \infty} a^{n_k} x = \xi$ for all $x \in X$ (see Proposition 4.1). It is enough to show that ξ is P -fixed. Setting $F_j := \overline{X}^{K_j}$, we have

$$\begin{aligned} F_j \cap X &\neq \emptyset & \forall j \in \mathbf{Z} \\ F_j &\supseteq F_\ell & \forall j \leq \ell \\ a^n F_j &= F_{j+n} & \forall j, n \in \mathbf{Z}. \end{aligned}$$

Picking $x \in F_j$, we have $a^{n_k} x \in F_{j+n_k} \subseteq F_j$ for all $k \geq 0$ and hence $\xi = \lim_{k \rightarrow \infty} a^{n_k} x \in F_j$ since F_j is closed in \overline{X} . Thus $\xi \in \bigcap_{j \in \mathbf{Z}} F_j = \overline{X}^N$. It follows that ξ is indeed P -fixed. \square

We now turn back to the particular case where $X = \mathbf{H}$ is the hyperbolic space attached to a strongly nondegenerate quadratic space (\mathcal{H}, Q) of signature $(\alpha, 1)$. We will focus on the study of continuous elementary representations $\pi : P \rightarrow \mathbf{O}_{L_+}(Q)$ for which $\pi(a)$ is hyperbolic with attracting fixed point L_+ ; that is, $|\chi(a)| > 1$.

PROPOSITION 5.2. *Let $\chi : P \rightarrow \mathbf{R}^*$ be a continuous homomorphism with $|\chi(a)| > 1$, let $\varrho : P \rightarrow \mathbf{O}(E)$ be a continuous orthogonal representation and $\tau := \chi \otimes \varrho$.*

- (i) *The orthogonal complement $E^{K_0} \ominus E^{K_0}$ of E^{K_0} in E^{K_0} is isomorphic to $\mathbf{H}_c^1(P, \tau)$, with isomorphism given by $v \mapsto f_{\sigma(v)}$, where for $v \in E^{K_0} \ominus E^{K_0}$,*

$$\sigma(v) := \sum_{n \leq -1} \tau(a)^{n+1} v$$

and $f_{\sigma(v)}$ is the standard cocycle uniquely determined by

$$f_{\sigma(v)}(k) = \tau(k)\sigma(v) - \sigma(v). \quad (\forall k \in K_0)$$

- (ii) *For $\eta \in \mathbf{H}_c^1(P, \tau)$, let $f_{\sigma(v)}$ be the standard cocycle representing η (with $v \in E^{K_0} \ominus E^{K_0}$). Then the subspace $I_\eta = \overline{\langle f_{\sigma(v)}(p) : p \in P \rangle}$ coincides with the closed cyclic subspace generated by v and, in fact,*

$$I_\eta = \overline{\langle f_{\sigma(v)}(n) : n \in N \setminus \bigcap_{j \in \mathbf{Z}} K_j \rangle}.$$

Proof. (i) The proof consists of two steps.

Claim 1: There is an isomorphism of topological vector spaces

$$\begin{aligned} \{v \in E : (Id - \tau(a))v \in E^{K_0}\} / E^{K_0} &\longrightarrow \mathbf{H}_c^1(P, \tau) \\ v &\longmapsto [f_v] \end{aligned}$$

where f_v is the standard cocycle determined uniquely by

$$f_v(k) := \tau(k)v - v \quad \forall k \in K_0.$$

Though this follows immediately from the Mayer-Vietoris sequence associated to topological HNN-extensions in continuous cohomology, we give an explicit proof:

Let f be a standard cocycle. Then repeated applications of the cocycle identity imply that for all $n \in \mathbf{Z}$ and $g \in P$ one has

$$(5.1) \quad f(a^n g a^{-n}) = \tau(a) f(g),$$

which, applied to $n \geq 0$ and $g \in K_0$ shows that f is determined by its restriction to K_0 . Since K_0 is compact, there is $v \in E$, uniquely determined modulo E^{K_0} , such that, for all $k \in K_0$,

$$(5.2) \quad f(k) = \tau(k)v - v.$$

Substituting (5.2) into (5.1) with $n = -1$ and $g \in K_0$, we get

$$(5.3) \quad (Id - \tau(a))v \in E^{K_0}.$$

Conversely, any cocycle f on K_0 defined by (5.2) with v satisfying (5.3) extends uniquely to a standard cocycle on P , which hence completes the proof of Claim 1.

Claim 2: The map

$$\begin{aligned} \sigma : E^{K_{-1}} \oplus E^{K_0} &\longrightarrow \{v \in E : (Id - \tau(a))v \in E^{K_0}\} / E^{K_0} \\ v &\longmapsto \sum_{n \leq -1} \tau(a)^{n+1} v \quad \text{mod } E^{K_0} \end{aligned}$$

is an isomorphism of topological vector spaces.

To start the proof of the Claim, observe that, since $|\chi(a)| > 1$, the operator $S := Id - \tau(a) = \tau(a)(\tau(a)^{-1} - Id)$ has a (bounded) inverse given by $S^{-1} = -\sum_{n=0}^{\infty} \tau(a)^{-(n+1)}$. Since $\tau(a)^\ell E^{K_0} = E^{K_\ell}$, this implies that

$$(5.4) \quad J := \overline{\bigcup_{\ell \leq -1} E^{K_\ell}} \supseteq \{v \in E : Sv \in E^{K_0}\}.$$

Defining $E_j := E^{K_j} \oplus E^{K_{j+1}}$, we have an orthogonal decomposition

$$(5.5) \quad J = E^{K_0} \oplus \bigoplus_{j \leq -1}^{\wedge} E_j.$$

Let $v = v_0 + \sum_{j \leq -1} v_j$ with $Sv \in E^{K_0}$; since we need to determine $v \text{ mod } E^{K_0}$, we may assume $v_0 = 0$. Then

$$Sv = -\tau(a)v_{-1} - \sum_{j=1}^{\infty} (\tau(a)v_{-(j+1)} - v_{-j})$$

where $\tau(a)v_{-1} \in E^{K_0}$ and $\tau(a)v_{-(j+1)} - v_{-j} \in E_{-j}$. In view of (5.4) and (5.5), saying that $Sv \in E^{K_0}$ is equivalent to saying that

$$\tau(a)v_{-(j+1)} = v_{-j}, \quad j \geq 1$$

which implies $v = \sum_{n \leq -1} \tau(a)^{n+1} v_{-1}$ and hence yields Claim 2.

(ii) It is clear that I_η is contained in $\overline{\langle \tau(p)v : p \in P \rangle}$. Conversely, since for $k \in K_0$ we have $f_{\sigma(v)}(k) = \tau(k)\sigma(v) - \sigma(v)$, it follows that

$$- \int_{K_0} f_{\sigma(v)}(k) dk = \sigma(v) - \text{Pr}_{E^{K_0}}(\sigma(v)) = \sigma(v)$$

and

$$- \int_{K_{-1}} \tau(u) \int_{K_0} f_{\sigma(v)}(k) dk du = v.$$

Thus $v \in I_\eta$, which is a closed invariant subspace, and hence contains the closure of its orbit $\overline{\langle \tau(p)v : p \in P \rangle}$. The additional formula for I_η follows since $f_{\sigma(v)}$ vanishes on $M = \bigcap_{j \in \mathbf{Z}} K_j$ by Lemma 4.6(ii). \square

6. REPRESENTATIONS OF CERTAIN PARABOLIC SUBGROUPS

6.1. Gelfand Pairs. We start this section recalling some definitions and facts about Gelfand pairs which will be essential in the sequel. We refer to [Si, Sec. 24], for example, for a complete discussion and proofs. Whilst this theory is generally presented for unitary representations, it carries over without changes to orthogonal representations; this will be our viewpoint here.

Let G be a locally compact group, $K < G$ a compact subgroup and let $C_c(G)^{\natural K}$ be the convolution algebra of bi- K -invariant functions on G with compact support. Then (G, K) is a Gelfand pair if $C_c(G)^{\natural K}$ is commutative. It is easy to see that the condition $x^{-1} \in KxK$ for all $x \in G$ is sufficient for (G, K) to be a Gelfand pair.

If (G, K) is a Gelfand pair, a continuous bi- K -invariant function $\varphi \in C(G)^{\natural K}$ is a spherical function if

- (1) $\varphi(e) = 1$, and
- (2) for all $f \in C_c(G)^{\natural K}$ there exists a constant c_f such that $\varphi * f = c_f \varphi$.

An irreducible orthogonal representation of G is K -spherical if there exists a nonzero K -invariant vector. If (G, K) is a Gelfand pair and (π, \mathcal{H}) is any irreducible orthogonal representation of G , then $\dim \mathcal{H}^K \leq 1$, and hence (π, \mathcal{H}) is K -spherical if and only if $\dim \mathcal{H}^K = 1$. Moreover, (equivalence classes of) K -spherical representations of a Gelfand pair (G, K) are in bijective correspondence with positive definite spherical functions, with the correspondence given by $\varphi(g) = \langle \pi(g)v, v \rangle$, where $v \in \mathcal{H}$ is a K -invariant vector of norm one and $\langle \cdot, \cdot \rangle$ is the inner product in \mathcal{H} .

6.2. In the remainder of this section, we shall consider a closed subgroup $P < \text{Aut}(\mathcal{T})$ of the automorphism group of a locally finite tree \mathcal{T} satisfying the following conditions:

- (1) P fixes a point $\xi \in \partial \mathcal{T}$;

(2) P acts doubly transitively on $\partial\mathcal{T} \setminus \{\xi\}$.

We shall assume throughout that the vertices of \mathcal{T} have valence at least three; observe that under these hypotheses it follows that P acts transitively on the vertices of \mathcal{T} , which is therefore a regular tree (and ξ is uniquely determined). Fix a geodesic line $c : \mathbf{Z} \rightarrow \mathcal{T}$ with $c(+\infty) = \xi$; by (2) there is a hyperbolic element $a \in P$ with axis c , translation length one and attracting fixed point ξ . We denote by K_j the stabilizer of $c(j)$ in P for $j \in \mathbf{Z}$. Then P has the structure $P = \langle a \rangle \rtimes N$ as in Section 5.

Let now (\mathcal{H}, Q) be strongly nondegenerate of signature $(\aleph_0, 1)$ and let L be an isotropic line. The main objective of this section is to prove:

THEOREM 6.1. *For every continuous homomorphism $\chi : P \rightarrow \mathbf{R}^*$ with $|\chi(a)| > 1$, there is up to conjugation a unique continuous representation $\lambda : P \rightarrow \mathbf{O}_L(Q)$ such that*

- (i) P acts on L by multiplication by χ .
- (ii) There is no proper closed P -invariant subspace of \mathcal{H} which is nondegenerate of index one.

This result is based on Section 4, Section 5 and on the following generalization of [N]:

PROPOSITION 6.2. *Let P_x be the stabilizer in P of a vertex $x \in \mathcal{T}$. Then there is a unique (up to equivalence) irreducible orthogonal representation of P having a P_x -fixed vector and whose restriction to N is nontrivial.*

We precede the proof with some intermediate results. For $j \in \mathbf{Z}$, we denote by H_j the horosphere centered at ξ passing through $c(j)$; furthermore, for $\ell \geq 0$, let $H_j(2\ell)$ be the intersection of H_j with the sphere of radius 2ℓ centered at $c(j)$. Notice that for $r \leq j$, the group K_r acts on $H_j(2\ell)$ for every $\ell > 0$, and moreover:

LEMMA 6.3. *The group K_r acts transitively on $H_j(2\ell)$ for all $r \leq j$ and $\ell \geq 0$.*

Proof. Write $\xi_{\pm} := c(\pm\infty)$ and pick $a, b \in H_j(2\ell)$. Complete the geodesic segments $[c(j+\ell), a]$ and $[c(j+\ell), b]$ to infinite rays $[c(j+\ell), \alpha]$ and $[c(j+\ell), \beta]$ respectively, where $\alpha, \beta \in \partial\mathcal{T}$. We may assume $\ell \neq 0$, thus $\alpha, \beta \neq \xi_-$. By double transitivity, there is $g \in P$ with $g(\alpha) = \beta$ and $g(\xi_-) = \xi_-$. The center of the tripods (ξ_+, α, ξ_-) and (ξ_+, β, ξ_-) is $c(j+\ell)$, hence g fixes that point. But since g fixes ξ_{\pm} and hence preserves $c(\mathbf{Z})$, it follows that $g \in \bigcap_{n \in \mathbf{Z}} K_n$. \square

COROLLARY 6.4. *(N, K_j) is a Gelfand pair for all $j \in \mathbf{Z}$.* \square

Proof. As mentioned Section 6.1, it is enough to prove that $n^{-1} \in K_j n K_j$ for all $n \in N$ and $j \in \mathbf{Z}$. We may assume that $n \notin K_j$. Then $n \in K_{\ell} \setminus K_{\ell-1}$ for some $\ell \geq j+1$ and $n(c(j)), n^{-1}(c(j)) \in H_j(2(\ell-j))$. By Lemma 6.3, this implies the existence of $k \in K_j$ with $kn(c(j)) = n^{-1}(c(j))$ and hence $n^{-1} \in K_j n K_j$. \square

Let \widehat{N} be the set of (equivalence classes of) irreducible orthogonal representations of N ; the group $\langle a \rangle$ acts on \widehat{N} by $a_*\pi(g) = \pi(aga^{-1})$, thus preserving the subset \widehat{N}^1 of representations that have a K_j -fixed vector for some $j \in \mathbf{Z}$. In fact, if we set

$$\widehat{N}_j^1 := \{(\pi, \mathcal{H}) \in \widehat{N}^1 : \mathcal{H}^{K_j} \neq 0, \mathcal{H}^{K_{j+1}} = 0\},$$

then $\widehat{N}^1 = \bigsqcup_{j \in \mathbf{Z}} \widehat{N}_j^1$ and $a_*\widehat{N}_j^1 = \widehat{N}_{j-1}^1$, so that any \widehat{N}_j^1 is a fundamental domain for the action of $\langle a \rangle$ on \widehat{N}^1 .

LEMMA 6.5. $|\widehat{N}_j^1| = 1$ for all $j \in \mathbf{Z}$.

Proof. Since $K_\ell \leq K_j$ for $\ell \leq j$, for any K_j -spherical representation (π, \mathcal{H}) we have that $\mathcal{H}^{K_\ell} \supset \mathcal{H}^{K_j}$; since (N, K_j) is a Gelfand pair, these spaces are of dimension one and hence $\mathcal{H}^{K_\ell} = \mathcal{H}^{K_j}$. Thus, in order to show that $|\widehat{N}_j^1| = 1$, it is sufficient to show that there is a unique positive definite K_j -spherical function φ with

$$\int_{K_{j+1}} \varphi(kg) dk = 0, \quad \forall g \in N.$$

Since it suffices to show that $|\widehat{N}_0^1| = 1$, we start by showing that the space

$$S_0 := \left\{ \varphi \in (\mathbf{C}(N)^{\natural})^{K_0} : \int_{K_1} \varphi(kg) dk = 0 \quad \forall g \in N \right\}$$

is of dimension one. By identifying N/K_0 with the horosphere H_0 , we can identify S_0 with the space of K_0 -invariant functions on $H_0 = \bigsqcup_{\ell \geq 0} H_0(2\ell)$, and by applying Lemma 6.3 we deduce that any function $\varphi \in S_0$ can be written as

$$\varphi = \sum_{\ell=0}^{\infty} \kappa_\ell \mathbf{1}_{H_0(2\ell)},$$

where $\kappa_\ell \in \mathbf{R}$. The condition defining S_0 means that the sum of the values of φ over any K_1 -orbit in H_0 is zero; denoting by q the valence of \mathcal{T} , this implies that $\kappa_\ell = 0$ for all $\ell \geq 2$ and $\kappa_0 + (q-2)\kappa_1 = 0$, thus proving the claim.

Thus, let $\varphi_0 \in S_0$ be the unique function such that $\varphi_0(e) = 1$. To complete the proof we need to show that for all $f \in \mathbf{C}_c(G)^{\natural K}$ there exists a constant c_f such that $\varphi_0 * f = c_f \varphi_0$. To this end, observe that any $f \in \mathbf{C}_c(G)^{\natural K}$ is a linear combination of characteristic functions $\mathbf{1}_{K_0}$ and $\mathbf{1}_{K_n \setminus K_{n-1}} := \chi_n$ for $n > 0$. In this notation we have that

$$\varphi_0 = \mathbf{1}_{K_0} - \frac{1}{q-2} \mathbf{1}_{K_1 \setminus K_0}.$$

Moreover, a direct computation shows that

$$\chi_n * \chi_m = \begin{cases} \mu(K_n \setminus K_{n-1}) \chi_m & \text{if } n < m \\ \mu(K_n \setminus K_{n-1}) \mathbf{1}_{K_n} - \mu(K_{n-1}) \chi_n & \text{if } n = m, \end{cases}$$

wherein μ is the Haar measure implicit in the chosen convolution structure on $C_c(G)$. Now it follows that

$$\begin{aligned}\varphi_0 * \chi_m &= 0 && \text{if } m > 1 \\ \varphi_0 * \chi_1 &= -\mu(K_0)\varphi_0 \\ \varphi_0 * \mathbf{1}_{K_0} &= \mu(K_0)\varphi_0,\end{aligned}$$

and hence shows that φ_0 is spherical.

Finally, observe that since φ_0 is compactly supported, it follows from the identity

$$\varphi_0 * \check{\varphi}_0 = c_{\check{\varphi}_0}\varphi_0,$$

where as usual $\check{\varphi}_0(x) := \varphi_0(x^{-1})$, that φ_0 is positive definite. \square

Proof of Proposition 6.2. There is no loss of generality in assuming $x = c(0)$, so that $P_x = K_0$. Let π_j be a representative of the unique equivalence class in \widehat{N}_j^1 and observe that $a_*\pi_j \cong \pi_{j-1}$. Thus the N -representation $\pi := \bigoplus_{j \in \mathbf{Z}} \pi_j$ extends canonically to a P -representation. To verify that π is irreducible, observe that if σ is any sub- P -representation of π , then $\sigma|_N$ is a direct sum of sub- N -representations σ_j of π_j , because the π_j are irreducible and pairwise inequivalent. Therefore, each σ_j is either zero or irreducible. The a -invariance of σ shows that either σ is zero or it coincides with π , hence π is irreducible as a P -representation. The uniqueness of π follows from Lemma 6.5 and the existence of a K_0 -fixed vector by construction. \square

Proof of Theorem 6.1. Existence: Let $\pi : P \rightarrow \mathbf{O}(E)$ be the continuous orthogonal representation of P constructed in Proposition 6.2 with underlying Hilbert space E . Let $\varrho := |\chi| \otimes \pi \otimes \chi^{-1}$ and endow E with the P -action defined by $\chi \otimes \varrho$. Since we have by construction that $E^{K^{-1}} \ominus E^{K_0} \neq 0$, then Proposition 5.2(i) implies that $H_c^1(P, \chi \otimes \varrho) \neq 0$ and hence Proposition 4.4 (with $G = P$) gives us a representation $\lambda : P \rightarrow \mathbf{O}_L(Q) \subseteq \mathbf{O}(Q)$, so that $(\chi \otimes \varrho)|_N = \pi|_N$ acting on L by χ . By Proposition 4.3(iii), one can extract the corresponding ‘‘irreducible’’ part and hence the existence is proved.

Uniqueness: Let us set $L_+ = L$. Since $|\chi(a)| \neq 1$, $\lambda(a)$ is hyperbolic; let L_- be the isotropic line representing the repelling fixed point and $\varrho : P \rightarrow \mathbf{O}(E)$ be the orthogonal representation obtained as in Section 4.1 *via* the identification $E \rightarrow {}^+L_+/L_+$ and $\eta \in H_c^1(P, \chi \otimes \varrho)$ the cohomology class defined by the above action. The irreducibility hypothesis, Propositions 4.7 and 5.2 then imply that

$$E = \overline{\langle f_{\sigma(v)}(p) : p \in P \rangle},$$

where $f_{\sigma(v)}$ is the standard cocycle representing η and moreover E is the cyclic subspace generated by $v \in E^{K^{-1}} \ominus E^{K_0}$.

Consider the orthogonal representation $\psi := |\chi|^{-1} \otimes \chi \otimes \varrho$ on E and let π denote the orthogonal representation of P given by Proposition 6.2. Then

$$\psi = m\pi \oplus \pi_1 \oplus \pi_2,$$

where $\pi_1|_N = \mathbf{1}$ and π_2 does not have any nonzero K_n -invariant vectors. Since $v \in E^{K^{-1}} \ominus E^{K_0}$, the projections of v to the components of π_1 and π_2 is zero. Being a cyclic vector, this implies $\psi = m\pi$ and therefore also $m = 1$. It now follows that $\varrho = |\chi| \otimes \chi^{-1} \otimes \pi$; in the notation of Proposition 5.2, $\tau = |\chi| \otimes \pi$. By Proposition 5.2, we have $H_c^1(P, \tau) \cong E^{K^{-1}} \ominus E^{K_0}$. Let now (π_n, \mathcal{H}_n) be the unique K_n -spherical representation of N that is not K_{n+1} -spherical. Then it follows from Lemma 6.5 that $\tau|_N = \pi|_N = \bigoplus_{n \in \mathbf{Z}} \pi_n$. Hence,

$$E^{K^{-1}} = \bigoplus_{n \geq -1}^{\wedge} \mathcal{H}_n^{K^{-1}}, \quad E^{K_0} = \bigoplus_{n \geq 0}^{\wedge} \mathcal{H}_n^{K_0}.$$

Observing that $\mathcal{H}_n^{K^{-1}} = \mathcal{H}_n^{K_0}$ for all $n \geq 0$, we deduce that $E^{K^{-1}} \ominus E^{K_0}$ has dimension one. Hence $\dim H_c^1(P, \tau) = 1$, which implies now by Proposition 4.4 that, up to conjugation, $\lambda : P \rightarrow \mathbf{O}_L(Q)$ is completely determined by χ . \square

7. REPRESENTATIONS OF G INTO $\mathbf{O}(Q)$

7.1. In this section \mathcal{T} denotes a regular or biregular tree of finite bivalency (r, s) with $r, s \geq 3$. A subgroup $G < \text{Aut}(\mathcal{T})$ satisfies the property T_2^+ if for every $\xi_1 \neq \xi_2$ in $\partial\mathcal{T}$ and $\eta_1, \eta_2 \in \partial\mathcal{T} \setminus \{\xi_1, \xi_2\}$ such that the distance between the projections of η_1 and η_2 on the geodesic $[\xi_1, \xi_2]$ is even, there exists $h \in G$ fixing ξ_1, ξ_2 and $h(\eta_1) = \eta_2$.

This property is implied by triple transitivity of the G -action on $\partial\mathcal{T}$ and implies double transitivity. The main result of this section is:

THEOREM 7.1. *Let $G < \text{Aut}(\mathcal{T})$ be a closed subgroup satisfying property T_2^+ , $\xi \in \partial\mathcal{T}$ and P the stabilizer of ξ in G . Let (\mathcal{H}, Q) be a strongly nondegenerate quadratic space of index 1 and $\pi : G \rightarrow \mathbf{O}(Q)$ a continuous, nonelementary representation. Then $\pi|_P$ has an irreducible indefinite component $\mathcal{H}_{1,P}$ and the canonical orthogonal decomposition $\mathcal{H} = \mathcal{H}_{1,P} \oplus \mathcal{H}_{0,P}$ is G -invariant.*

Let $\pi_1, \pi_2 : G \rightarrow \mathbf{O}(Q)$ be nonelementary continuous representations such that $\pi_1|_P = \pi_2|_P$. Then the restriction of π_1 and π_2 to the indefinite irreducible components of P coincide.

7.2. Let G be a locally compact group boundedly generated by $\{s\} \cup P$, where $s \in G$ and $P < G$ is a closed subgroup with the structure considered in Section 5; assume further that:

- (1) $\langle s \rangle$ is relatively compact, and
- (2) $\{a^n s a^n s^{-1} : n \geq 1\}$ is relatively compact.

(We recall that a group G is said *boundedly generated* by a subset $X \subseteq G$ if there is $n \in \mathbf{N}$ with $X^n = G$.)

PROPOSITION 7.2. *Let X be a complete CAT(-1) space and $G \times X \rightarrow X$ a continuous, nonelementary isometric action. Then a acts as a hyperbolic element, s exchanges both fixed points of a and $(\partial X)^P$ is the attracting fixed point of a .*

Proof. We use Lemma 5.1 and we distinguish the three cases for a .

a is elliptic: Then $X^P \neq \emptyset$ and since $\langle s \rangle$ is relatively compact and G is boundedly generated by $\langle s \rangle$ and P , the G -orbits in X are bounded. Hence $X^G \neq \emptyset$, contradicting non-elementarity.

a is parabolic: Then $\partial X^P = \{\xi\}$. Let $\{n_i\}_{i \geq 1}$ be a sequence such that $a^{n_i}x \rightarrow \xi$; then $a^{-n_i}x \rightarrow \xi$, and since $\{a^n s a^n s^{-1} : n \geq 1\}$ is bounded, we have that $s a^{n_i} s^{-1} x \rightarrow \xi$ which, in view of the fact that $a^{n_i} s^{-1} x \rightarrow \xi$, implies that $s(\xi) = \xi$ and hence $G\xi = \xi$, contradiction.

Thus, *a is hyperbolic.* Let now ξ_-, ξ_+ be respectively the repelling and the attracting fixed points of a on ∂X . Since $a^n x \rightarrow \xi_+$, $a^{-n} x \rightarrow \xi_-$, and $\{a^n s a^n s^{-1} : n \geq 1\}$ is bounded, we deduce that $s a^n s^{-1} x \rightarrow \xi_-$ and hence $s(\xi_+) = \xi_-$ and $s(\xi_-) = \xi_+$.

Finally, we know $(\partial X)^P \ni \xi_+$ from Lemma 5.1; since $(\partial X)^{\langle a \rangle} = \{\xi_{\pm}\}$, it remains only to observe that ξ_- is not P -fixed, since otherwise the set $\{\xi_{\pm}\}$ would be preserved by G . \square

7.3. Let now $G < \text{Aut}(\mathcal{T})$ be any closed subgroup which acts doubly transitively on $\partial \mathcal{T}$. Then (see [BM2] Sec. 4.1 and 0.4):

- For every $\xi \in \partial \mathcal{T}$, the Busemann character $\chi_{\xi} : G_{\xi} \rightarrow \mathbf{Z}$ has image \mathbf{Z} or $2\mathbf{Z}$ depending on whether G is vertex transitive or not;
- For every geodesic $c : \mathbf{Z} \rightarrow \mathcal{T}$ there is $s \in G$ and $n_0 \in \mathbf{Z}$ with $s(c(\pm\infty)) = c(\mp\infty)$ and $s(c(n_0)) = c(n_0)$;
- For any $\xi \neq \eta$ in $\partial \mathcal{T}$ and $s \in G$ exchanging ξ and η , we have $G = G_{\xi} \cup G_{\xi} s G_{\xi}$.

LEMMA 7.3. *Let $G < \text{Aut}(\mathcal{T})$ be a closed subgroup satisfying T_2^+ , $c : \mathbf{Z} \rightarrow \mathcal{T}$ a geodesic, $\xi_{\pm} = c(\pm\infty)$, and $S = \{s \in G : s(\xi_{\pm}) = (\xi_{\mp}), s(c(0)) = c(0)\}$. Let also $K_j = G_{\xi} \cap G(c(j))$. Then, given $j \in \mathbf{Z}$, $n \in K_j \setminus K_{j-1}$ and $s \in S$, there exists $h \in G$ such that:*

- (i) For all $q \in \mathbf{Z}$, $hc(q) = c(q - 2j)$;
- (ii) For all $s', s'' \in S$, $s' n s h n s'' \in K_{-j} \setminus K_{-j-1}$.

Proof. Since $n \in K_j \setminus K_{j-1}$, then $\text{Pr}_{[\xi_+, \xi_-]}(n\xi_-) = c(j)$ and $\text{Pr}_{[\xi_+, \xi_-]}(s^{-1}n^{-1}\xi_-) = c(-j)$. Thus property T_2^+ implies that there exists $h \in G_{\xi_+} \cap G_{\xi_-}$ such that $hn\xi_- = s^{-1}n^{-1}\xi_-$ so that (i) follows. Let now $s', s'' \in S$ and set $g = s' n s h n s''$. Then we have:

$$(\circ) \quad g\xi_+ = s'nshns''\xi_+ = s'nshn\xi_- = s'nss^{-1}n^{-1}\xi_- = s'\xi_- = \xi_+.$$

(\circ) $gc(-j) = s'nshns''c(-j) = s'nshnc(j) = s'nshc(j) = s'nsc(-j) = s'nc(j) = s'c(j) = c(-j)$. Thus $g \in K_{-j}$.

(\circ) $gc(-j-1) = s'nshns''c(-j-1) = s'nshnc(j+1) = s'nshc(j+1) = s'nsc(-j+1) = s'nc(j-1)$. But since $n \in K_j$, we have that $nc(j-1) \neq c(j+1)$ and hence $s'nc(j-1) \neq c(-j-1)$. Thus $g \notin K_{-j-1}$. \square

7.4. We are finally ready to give the

Proof of Theorem 7.1. Let $\pi : G \rightarrow \mathbf{O}(Q) \rightarrow \text{Isom}(\mathbf{H})$ be a continuous nonelementary action. Being the stabilizer of $\xi \in \partial\mathcal{T}$, P has the structure of Section 5; we shall use the corresponding notation for a . In view of Section 7.3, we can choose s and a parametrisation $c : \mathbf{Z} \rightarrow \mathcal{T}$ of the axis of a such that $sc(0) = c(0)$ and $sc(\pm\infty) = c(\mp\infty)$. Observe further that these notations also put us in the setting of Section 7.2. By Proposition 7.2, $\pi(a)$ is hyperbolic and hence cannot fix a point in \mathbf{H} or preserve any horosphere. It follows from Proposition 4.3 applied to P that $\pi|_P$ has an irreducible indefinite component $\mathcal{H}_{1,P}$.

We need to show that $\mathcal{H}_{1,P}$ is G -invariant and that, on $\mathcal{H}_{1,P}$, the representation π is determined by $\pi|_P$. Let L_{\pm} be the attr./repell. fixed points of $\pi(a)$ and let $L = L_+ \oplus L_-$, $F = {}^{\perp}L$. We are in the setting of Proposition 4.3 for P (instead of G) and we adopt its notation. By Proposition 4.7, $\mathcal{H}_{1,P} = L \oplus j^{-1}(I_{\eta})$. By Proposition 7.2, $\pi(s)$ exchanges L_{\pm} . It is enough to show that $\pi(s)$ preserves $j^{-1}(I_{\eta})$ and that its restriction to $\mathcal{H}_{1,P}$ is determined by $\pi|_P$.

We adopt the notation of Appendix A with ℓ_{\pm} . Then

$$\pi(s) = \begin{pmatrix} 0 & \mu & 0 \\ \mu^{-1} & 0 & 0 \\ 0 & 0 & \pi_0(s) \end{pmatrix},$$

where $\pi_0(s)$ is orthogonal and $\mu \in \mathbf{R}^*$.

Fix $j \in \mathbf{Z}$, $n \in K_j \setminus K_{j-1}$ and h as in Lemma 7.3 such that $g := snshns \in K_{-j} \setminus K_{-j-1}$. We write

$$\pi(n) = \begin{pmatrix} \chi(n) & \alpha(n) & N_2^+ \\ 0 & \chi(n)^{-1} & 0 \\ 0 & N_3^- & \pi_0(n) \end{pmatrix},$$

and likewise

$$\pi(g) = \begin{pmatrix} \chi(g) & \alpha(g) & M_2^+ \\ 0 & \chi(g)^{-1} & 0 \\ 0 & M_3^- & \pi_0(g) \end{pmatrix}.$$

As to $\pi(h)$, since it fixes both L_{\pm} , it is of the form

$$\pi(h) = \begin{pmatrix} \chi(h) & 0 & 0 \\ 0 & \chi(h)^{-1} & 0 \\ 0 & 0 & \pi_0(h) \end{pmatrix}.$$

Computing $g = snshns$ we find

$$(7.1) \quad \chi(g) = \mu^{-1} \chi(n^{-1}h) \alpha(n),$$

and

$$(7.2) \quad M_3^- = \chi(hn) \pi_0(s) N_3^-.$$

Equation (7.1) shows that μ is determined in terms of $\pi|_P$. We are left to determine $\pi_0(s)$. Write f for the standard cocycle associated to η ; then (7.2) gives

$$\pi_0(s) f(n) = \chi(gh)^{-1} f(g).$$

We obtain such a formula for every $n \in N \setminus \bigcap_{j \in \mathbf{Z}} K_j$. Thus we are done since by Proposition 5.2(ii), I_η is spanned by these $f(n)$. \square

Proof of Theorem C. By Proposition 3.4, we are reduced to study homomorphisms from G into $\mathbf{O}_+(Q)$ and can thus apply the results obtained so far. The existence statement in Theorem C follows from Theorem A and Proposition 4.3(iii).

For the uniqueness part, let $\xi \in \partial \mathcal{T}_r$, P the stabilizer of ξ in G and $a \in P$ a hyperbolic element with attracting fixed point ξ and translation length 1. Let $\pi : G \rightarrow \mathbf{O}_+(Q)$ be a nonelementary continuous representation. Let $c : \mathbf{Z} \rightarrow \mathcal{T}_r$ be a parametrization of the axis of a , and let K_n be, as usual, the stabilizer in P of $c(n)$. Using that G is doubly transitive on $\partial \mathcal{T}_r$, we see that any other hyperbolic element b with translation length 1 is conjugate to an element of the form $a \cdot k$, where $k \in \bigcap_{n \in \mathbf{Z}} K_n$. Since $\pi(a)$ is hyperbolic (Proposition 7.2), and using that a normalises $\bigcap_{n \in \mathbf{Z}} K_n$, one sees that $\pi(k)$ fixes pointwise the axis of $\pi(a)$ and hence $\pi(a)$ and $\pi(b)$ have the same translation length, say ℓ_π . Let L_+ be the attractive fixed point of $\pi(a)$ and χ the character by which P acts on L_+ . Then, since π takes values in $\mathbf{O}_+(Q)$, we have that χ takes values in \mathbf{R}_+ which implies first that $\chi(a) = e^{\ell_\pi}$ and then that χ is trivial on N , since N is an increasing union of compact groups. This shows that χ is completely determined by ℓ_π .

Assume now that π is irreducible. Then Theorem 7.1 implies that $\pi|_P$ is irreducible, Theorem 6.1 that it is completely determined by χ (and hence by ℓ_π), and Theorem 7.1 again that π is completely determined by ℓ_π . \square

8. EXPLICIT CONSTRUCTIONS

8.1. Let \mathcal{T} be any simplicial tree. Let $\alpha + 1$ be the cardinal of the vertex set V of \mathcal{T} and let (\mathcal{H}, Q) be a strongly nondegenerate quadratic space of signature $(\alpha, 1)$; let \mathbf{H} be the corresponding hyperbolic space. Denote by G the (abstract)

group $G = \text{Aut}(\mathcal{T})$. We denote by d both the metric on \mathbf{H} and the metric on (the geometric realization of) \mathcal{T} that gives unit length edges.

THEOREM 8.1. *For every $\lambda > 1$ there is an embedding $\Psi : \mathcal{T} \rightarrow \mathbf{H}$ and a representation $\pi : G \rightarrow \mathbf{O}(Q) \rightarrow \text{Isom}(\mathbf{H})$ such that:*

- (i) *The map Ψ is G -equivariant for π .*
- (ii) *$\lambda^{d(x,y)} = \cosh d(\Psi x, \Psi y)$ for any two vertices x, y of \mathcal{T}*
- (iii) *Ψ extends to an equivariant boundary map $\partial\Psi : \partial\mathcal{T} \rightarrow \partial\mathbf{H}$ which is a homeomorphism onto its image.*
- (iv) *$\Psi(V)$ is cobounded in the convex hull $\mathcal{C} \subseteq \mathbf{H}$ of the image of $\partial\Psi$.*

REMARK 8.2. The formula in (ii) shows in particular that $d(\Psi x, \Psi y)$ is asymptotically proportional to $d(x, y)$. If we denote by $b^{\mathcal{T}}$ the Busemann cocycle for \mathcal{T} and b is the one for \mathbf{H} as in Section 3, we have

$$b_{\Psi\xi}(\Psi x, \Psi y) = b_{\xi}^{\mathcal{T}}(x, y) \ln \lambda \quad \forall \xi \in \partial\mathcal{T}, \forall x, y \in V.$$

We can give right away the construction of Ψ ; the remainder of the section will be devoted to proving the properties stated in Theorem 8.1.

Fix a vertex $w \in V$. By Proposition 2.7, we may identify \mathcal{H} with $\ell^2(V)$ in such a way that the bilinear form B associated to Q reads

$$B(f, g) = \sum_{v \in V, v \neq w} f(v)g(v) - f(w)g(w).$$

We define a map $V \rightarrow \mathcal{H}$, $v \mapsto f_v$ as follows. Denote for $u \in V$ by δ_u the unit function supported on u ; then

$$f_v := \lambda^{d(w,v)} \delta_w + \sqrt{\lambda^2 - 1} \sum_{k=1}^{d(w,v)} \lambda^{d(w,v)-k} \delta_{u_k},$$

where $w, u_1, u_2, \dots, u_{d(w,v)} = v$ is the geodesic from w to v (it is understood that the right hand side summation is zero when $v = w$). A computation gives $Q(f_v) = -1$ so that f_v is in the negative cone C_- ; now Ψ is the resulting map $V \rightarrow C_- \rightarrow \mathbf{H}$ extended to \mathcal{T} by sending each edge to a geodesic segment. Each element $\xi \in \partial\mathcal{T}$ can be realized by a unique geodesic ray of vertices $\{v_k\}_{k=0}^{\infty}$ with $v_0 = w$; we define $(\partial\Psi)(\xi)$ by considering the element f_{ξ} of the isotropic cone C_0 given by

$$f_{\xi} := \delta_w + \sqrt{\lambda^2 - 1} \sum_{k=1}^{\infty} \lambda^{-k} \delta_{v_k}.$$

Observe that one obtains a multiple of f_{x_k} , hence the same point Ψx_k , by truncating the above sum at k . It follows that the resulting map $\overline{\mathcal{T}} \rightarrow \overline{\mathbf{H}}$ is continuous; the claim (iii) now follows from Remark 8.2 and claim (i). The formula of claim (ii) can be verified by inspection; however, we shall see that it can be reduced to the obvious case $u = w$.

The strategy for the proof of Theorem 8.1 is first to construct π in the case where \mathcal{T} is *regular*, that is, G acts transitively on V . The general case will follow by the naturality of our construction with respect to the pointed tree (\mathcal{T}, w) .

8.2. Regular Case. We assume that $G = \text{Aut}(\mathcal{T})$ acts transitively on V . Fix a neighbour z of w and denote by $K < G$ the stabilizer of w , by L the subgroup of G preserving the set $\{w, z\}$ and by $E = K \cap L$ the pointwise stabilizer of $\{w, z\}$. The K -action on V preserves B and hence induces a representation $\pi : K \rightarrow \mathbf{O}(Q)$ by $\pi(g)\delta_u = \delta_{gu}$ ($g \in K$). We extend now π by defining $\pi : L = E \sqcup (L \setminus E) \rightarrow \mathbf{O}(Q)$ as follows: the map is already defined on $E = K \cap L$; for every $g \in L \setminus E$ and every $u \in V$, set

$$(8.1) \quad \pi(g)\delta_u := \begin{cases} \lambda\delta_w + \sqrt{\lambda^2 - 1}\delta_z & \text{if } u = w. \\ -\sqrt{\lambda^2 - 1}\delta_w - \lambda\delta_z & \text{if } u = z. \\ \delta_{gu} & \text{otherwise.} \end{cases}$$

It is a matter of computation to verify that $\pi(g)$ is in $\mathbf{O}(Q)$.

PROPOSITION 8.3. *The map $\pi : L \rightarrow \mathbf{O}(Q)$ extends uniquely to a homomorphism $\pi : G \rightarrow \mathbf{O}(Q)$.*

Proof. We start by showing that the map $\pi : L \rightarrow \mathbf{O}(Q)$ is a homomorphism; that is, we need to verify that $\pi(g)\pi(g') = \pi(gg')$ holds on L , which we do by discussing the cases according to where g, g' are in the coset decomposition $L = E \sqcup (L \setminus E)$. There is nothing to do if g, g' are both in E since π is a homomorphism on K . We shall write out the verification in the case $g, g' \in L \setminus E$; the two remaining cases are simpler and similar. Let thus $g, g' \in L \setminus E$. Then $\pi(gg')\delta_u = \delta_{gg'u}$ for all $u \in V$ since $gg' \in E$ (as E has index two in L). On the other hand, we have:

Case $u = w$:

$$\begin{aligned} \pi(g)\pi(g')\delta_w &= \pi(g)(\lambda\delta_w + \sqrt{\lambda^2 - 1}\delta_z) = \\ &= \lambda(\lambda\delta_w + \sqrt{\lambda^2 - 1}\delta_z) + \sqrt{\lambda^2 - 1}(-\sqrt{\lambda^2 - 1}\delta_w - \lambda\delta_z) = \delta_w, \end{aligned}$$

which is indeed $\delta_{gg'u}$ since E fixes w .

Case $u = z$:

$$\begin{aligned} \pi(g)\pi(g')\delta_z &= \pi(g)(-\sqrt{\lambda^2 - 1}\delta_w - \lambda\delta_z) = \\ &= -\sqrt{\lambda^2 - 1}(\lambda\delta_w + \sqrt{\lambda^2 - 1}\delta_z) - \lambda(-\sqrt{\lambda^2 - 1}\delta_w - \lambda\delta_z) = \delta_z, \end{aligned}$$

which is indeed $\delta_{gg'u}$ since E fixes z .

Case $u \neq w, z$: then we have also $g'u \neq w, z$ and hence

$$\pi(g)\pi(g')\delta_u = \pi(g)\delta_{g'u} = \delta_{gg'u} = \pi(gg')\delta_u.$$

To show that the map π defined on L extends uniquely to a homomorphism on G , observe the G -action on (the first barycentric subdivision of) \mathcal{T} determines a

splitting of G into an amalgamation $G = K *_E L$. Therefore the statement follows from the universal property of amalgamations. \square

REMARK 8.4. The definition of π on G is independent of the choice of z : Indeed, observe first that K acts transitively on the set of neighbours of w . If k is any element of K , we obtain another neighbour kz of w and another amalgamation $G = K *_E kLk^{-1}$. With π defined as before using L , one checks immediately that for $g \in kLk^{-1} \setminus kEk^{-1}$ the formula (8.1) for $\pi(g)$ remains valid upon replacing z with kz .

We turn to point (i) in Theorem 8.1. Pick a vertex $v \in V$ and let $n = d(v, w)$. There is a hyperbolic element a of translation length one admitting an axis $\{u_k\}_{k \in \mathbf{Z}}$ such that $u_0 = w$, $u_n = v$ and $au_k = u_{k+1}$ for all $k \in \mathbf{Z}$. Notice that K and a generate G ; since moreover Ψ is K -invariant by its construction, we need only verify that $\Psi(av) = \pi(a)\Psi(v)$. By Remark 8.4 there is no loss of generality in assuming that $z = u_1$. Let s be an element of $L \setminus E$ preserving $\{u_k\}$; that is, $su_k = u_{1-k}$ for all $k \in \mathbf{Z}$. Then $a = st$ for $t \in K$ such that $tu_k = u_{-k}$ for all $k \in \mathbf{Z}$, and thus an immediate computation using (8.1) for $\pi(s)$ shows that we have for all $k \in \mathbf{Z}$

$$\pi(a)\delta_{u_k} = \pi(s)\pi(t)\delta_{u_k} = \begin{cases} \lambda\delta_w + \sqrt{\lambda^2 - 1}\delta_{u_1} & \text{if } k = 0. \\ -\sqrt{\lambda^2 - 1}\delta_w - \lambda\delta_{u_1} & \text{if } k = -1. \\ \delta_{u_{k+1}} & \text{otherwise.} \end{cases}$$

Now we can compute

$$\begin{aligned} \pi(a)f_v &= \pi(a)\left(\lambda^n\delta_w + \sqrt{\lambda^2 - 1}\sum_{k=1}^n \lambda^{n-k}\delta_{u_k}\right) \\ &= \lambda^n(\lambda\delta_w + \sqrt{\lambda^2 - 1}\delta_{u_1}) + \sqrt{\lambda^2 - 1}\sum_{k=1}^n \lambda^{n-k}\delta_{u_{k+1}} \\ &= \lambda^{n+1}\delta_w + \sqrt{\lambda^2 - 1}\sum_{k=1}^{n+1} \lambda^{n+1-k}\delta_{u_k} = f_{u_{k+1}} = f_{av}, \end{aligned}$$

and claim (i) is proved. Now the transitivity of G reduces (ii) to the case where one of the vertices is w , which is an immediate computation. As (iii) was addressed before, we are left with proving (iv).

PROPOSITION 8.5. *Every point $x \in \mathcal{C}$ is at distance at most $\cosh^{-1}\sqrt{1 + \lambda}$ of some element of $\Psi(V)$.*

Proof. Every element of \mathbf{H} is represented by a unique function in $\ell^2(V)$ with value one on w . In fact, if D' denotes the unit ball in $\ell^2(V \setminus \{w\})$, the set of such functions is $D := \delta_w + D'$. This gives the *Klein model* in finite dimensions, and

thus it follows that geodesics in \mathbf{H} correspond to affine lines in D . It is therefore enough to show the claim for any finite convex combination $x = \sum_{\xi \in S} s_\xi f_\xi$ where $S \subseteq \partial \mathcal{T}$ is a finite set and $s : S \rightarrow (0, 1)$ is any function $\xi \mapsto s_\xi$ of sum one (observe that S must contain at least two points since $0 < s_\xi < 1$).

We write $\beta = b(\cdot, \delta_w)$ for Busemann function on \mathbf{H} normalised at δ_w and $\beta^\mathcal{T} = b^\mathcal{T}(\cdot, w)$ for the analogous Busemann function on \mathcal{T} . Consider the function $\psi_s : V \rightarrow \mathbf{R}_+^*$ defined by

$$\psi_s(v) := \sum_{\xi \in S} s_\xi \lambda^{\beta_\xi^\mathcal{T}(v)}.$$

This function admits a minimum $v_0 \in V$; indeed, outside a finite subset of V determined by the configuration of S , it increases monotonically with the distance to this subset. We shall see that $\Psi(v_0)$ is at distance at most $\cosh^{-1} \sqrt{1 + \lambda}$ of x .

Pick $g \in G$ such that $gv_0 = w$; then w is a minimum of the function $v \mapsto \psi_s(g^{-1}v)$, which in view of $\beta_\xi^\mathcal{T}(g^{-1}v) = \beta_{g\xi}(v) - \beta_{g\xi}(gw)$ reads

$$\psi_s(g^{-1}v) = \sum_{\eta \in gS} s_{g^{-1}\eta} \lambda^{\beta_\eta^\mathcal{T}(v) - \beta_\eta^\mathcal{T}(gw)}.$$

Thus, setting $(g_\star s)_\xi := \lambda^{-\beta_\xi^\mathcal{T}(gw)} s_{g^{-1}\xi}$, it follows that w is also a minimum of the function $\psi_{s'}$ for $s' := g_\star s / \sum_\xi (g_\star s)_\xi$. Setting

$$\sigma_v := \sum \{s'_\xi : v \text{ is in the ray } [w, \xi]\} \quad \forall v \in V$$

the minimality implies for every neighbour v of w

$$1 = \psi_{s'}(w) \leq \psi_{s'}(v) = \lambda^{-1} \sum \{s'_\xi : v \in [w, \xi]\} + \lambda \sum \{s'_\xi : v \notin [w, \xi]\} = \lambda^{-1} \sigma_v + \lambda(1 - \sigma_v),$$

and hence $\sigma_v \leq \lambda/(1 + \lambda)$. This in turn implies

$$\sigma_v \leq \frac{\lambda}{1 + \lambda} \quad \forall v \in V, v \neq w.$$

The formula (3.1) in Section 3 shows that for every $g \in G$ and $h \in C_0$ we have

$$\beta_{gh}(g\delta_w) = \ln \frac{B(gh, g\delta_w) \sqrt{-Q(\delta_w)}}{B(gh, \delta_w) \sqrt{-Q(g\delta_w)}} = \ln \frac{B(h, \delta_w)}{B(gh, \delta_w)} = \ln \frac{h(w)}{(gh)(w)}.$$

Thus we have for all $\xi \in \partial \mathcal{T}$

$$(gf_\xi)(w) f_\xi(w) e^{-\beta_{gf_\xi}(g\delta_w)} = f_\xi(w) \lambda^{-\beta_{g\xi}^\mathcal{T}(gw)}$$

and since gf_ξ is proportional to $f_{g\xi}$ we deduce $gf_\xi = \lambda^{-\beta_{g\xi}^\mathcal{T}(gw)} f_{g\xi}$. We conclude that gx is represented in \mathbf{H} by the element $y := \sum_{\xi \in gS} s'_\xi f_\xi$ of D . We proceed

now to compute

$$\cosh d(\delta_w, y) = -\frac{B(\delta_w, y)}{\sqrt{Q(\delta_w)Q(y)}} = \frac{1}{\sqrt{-Q(y)}}.$$

If S_k is the sphere of radius k around w , we deduce from the definition of f_ξ

$$y = \delta_w + \sqrt{\lambda^2 - 1} \sum_{k=1}^{\infty} \sum_{v \in S_k} \lambda^{-k} \sigma_v \delta_v$$

and therefore

$$Q(y) = -1 + (\lambda^2 - 1) \sum_{k=1}^{\infty} \lambda^{-2k} \sum_{v \in S_k} \sigma_v^2 \leq -1 + (\lambda^2 - 1) \frac{\lambda}{1 + \lambda} \sum_{k=1}^{\infty} \lambda^{-2k} \sum_{v \in S_k} \sigma_v.$$

Using $\sum_{v \in S_k} \sigma_v = 1$ one gets finally $Q(y) \leq -1/(1 + \lambda)$. It follows that

$$d(\Psi v_0, x) = d(\Psi w, gx) = d(\delta_w, y) \leq \cosh^{-1} \sqrt{1 + \lambda}.$$

□

This proposition completes the proof of Theorem 8.1 in the regular case. □

8.3. General Case. Suppose now that \mathcal{T} is a general tree with vertex set V of cardinal $\alpha + 1$ and $G = \text{Aut}(\mathcal{T})$. Complete \mathcal{T} to a regular tree \mathcal{T}' with vertex set $V' \supseteq V$ of cardinal $\alpha' + 1$. We keep the notation of Section 8.1 and define likewise $\mathcal{H}' = \ell^2(V')$, \mathbf{H}' , B' , $G' = \text{Aut}(\mathcal{T}')$, etc.; take $w' := w \in V \subseteq V'$ and observe that \mathbf{H} is a hyperbolic subspace of \mathbf{H}' . Let Ψ', π' be the maps associated to \mathcal{T}' by the proof for the regular case.

Denote by $L_0 < L < G'$ the pointwise stabilizer, respectively stabilizer, of \mathcal{T} ; since any automorphism of \mathcal{T} can be extended to some automorphism(s) of \mathcal{T}' , we have a natural identification $G = L/L_0$. It follows at once from the definition of Ψ, Ψ' and π, π' that the restriction of Ψ' gives Ψ , while π' descends to π . In fact, as all definitions for \mathcal{T}' vanish on $V' \setminus V$ when restricted to \mathcal{T} , the only part of Theorem 8.1 for \mathcal{T} that does not follow immediately from the case of \mathcal{T}' is point (iv). But the proof given above, when applied to \mathcal{T}' and to the corresponding convex hull \mathcal{C}' , shows in fact that whenever $x \in \mathcal{C} \subseteq \mathcal{C}'$ is a finite affine convex combination of elements in $\Psi(\partial\mathcal{T})$, then the vertex $v_0 \in V'$ is actually in V . This follows indeed from the definition of the function ψ_s and thus concludes the proof of Theorem 8.1. □

APPENDIX A. MATRIX REPRESENTATIONS

Let (\mathcal{H}, Q) a strongly nondegenerate quadratic space of signature $(\alpha, 1)$. If L_-, L_+ are two distinct isotropic lines, define $L := L_+ \oplus L_-$ and $F = {}^\perp L$, so that

$$\mathcal{H} = L \oplus F.$$

On L and F we consider the restrictions $B|_{L \times L}$ and $B|_{F \times F}$, and if A is a continuous linear operator between any of these spaces, A^* denotes the adjoint with respect to these strongly nondegenerate bilinear forms. For any continuous linear operator $T : \mathcal{H} \rightarrow \mathcal{H}$, define

$$\begin{aligned} A_1 &= p_L T|_L, & A_2 &= P_L T|_F, \\ A_3 &= P_F T|_L, & A_4 &= p_F T|_F. \end{aligned}$$

Then $T \in \mathbf{O}(Q)$ if and only if the following conditions are satisfied:

$$\begin{aligned} A_1^* A_1 + A_3^* A_3 &= Id_L; \\ A_2^* A_2 + A_4^* A_4 &= Id_F; \\ A_1^* A_2 + A_3^* A_4 &= 0; \\ A_2^* A_1 + A_4^* A_3 &= 0; \end{aligned}$$

Observe that by taking adjoints, the last two conditions are equivalent.

We shall look more closely at $\mathbf{O}_{L_+}(Q)$, the stabilizer in $\mathbf{O}(Q)$ of L_+ . For this, let $L_{\pm} = \mathbf{R}\ell_{\pm}$, with $B(\ell_+, \ell_-) = 1$. We represent A_1 by a two-by-two real matrix; $A_2 : F \rightarrow L$ will be represented by two linear forms A_2^+ and A_2^- given by $A_2(e) = A_2^+(e)\ell_+ + A_2^-(e)\ell_-$; $A_3 : L \rightarrow F$ will be represented by two vectors $A_3^+ = A_3(\ell_+)$, $A_3^- = A_3(\ell_-)$, and hence

$$T = \begin{pmatrix} A_1 & A_2^+ \\ A_3^+ & A_3^- & A_4 \end{pmatrix}.$$

Then $T \in \mathbf{O}_{L_+}(Q)$ if and only if it has the form

$$\begin{pmatrix} \lambda & \alpha & A_2^+ \\ 0 & \lambda^{-1} & 0 \\ 0 & A_3^- & A_4 \end{pmatrix}$$

with $\lambda \in \mathbf{R}^{\times}$, $A_3^- \in F$, $A_4 \in \mathbf{O}(F)$ and α, A_2^+ are determined by

$$\alpha = -\frac{\lambda}{2}Q(A_3^-),$$

and

$$A_2^+(v) = -\lambda B(A_4(v), A_3^-), \quad \text{for all } v \in F.$$

The inverse of

$$S = \begin{pmatrix} \mu & \beta & B_2^+ \\ 0 & \mu^{-1} & 0 \\ 0 & B_3^- & B_4 \end{pmatrix} \in \mathbf{O}_{L_+}(Q)$$

is given by

$$S^{-1} = \begin{pmatrix} \mu^{-1} & \beta & -\mu^{-1}B_2^+B_4^{-1} \\ 0 & \mu & 0 \\ 0 & -\mu B_4^{-1}(B_3^-) & B_4^{-1} \end{pmatrix}$$

and the conjugate STS^{-1} of T by S has the following entries

$$\begin{aligned} (STS^{-1})_{1,1} &= \lambda = (STS^{-1})_{2,2}^{-1} \\ (STS^{-1})_{2,1} &= (STS^{-1})_{2,3} = (STS^{-1})_{3,1} = 0 \\ (STS^{-1})_{1,2} &= \lambda\mu\beta + \mu^2\alpha - \mu^2 A_2^+ B_4^{-1} B_3^- + \lambda^{-1}\mu\beta - \mu B_2^+ A_3^- - \mu B_2^+ A_4 B_4^{-1} B_3^- \\ (STS^{-1})_{1,3} &= -\lambda B_2^+ B_4^{-1} + \mu A_2^+ B_4^{-1} + B_2^+ A_4 B_4^{-1} \\ (STS^{-1})_{3,2} &= \lambda^{-1}\mu B_3 + \mu B_4^{-1} A_3^- - \mu B_4 A_4 B_4^{-1} B_3^- \\ (STS^{-1})_{3,3} &= B_4 A_4 B_4^{-1} \end{aligned}$$

In particular, if $|\lambda| \neq 1$, by choosing $\mu = 1$, $B_4 = Id$, there exists B_3^- such that

$$A_3^- + (\lambda^{-1} - A_4)(B_3^-) = 0.$$

Then, using the relations (i) and (ii), one can see that T is conjugate to

$$\begin{pmatrix} \lambda & 0 & 0 \\ 0 & \lambda^{-1} & 0 \\ 0 & 0 & A_4 \end{pmatrix}.$$

and is hence hyperbolic; conversely, one can show that if $|\lambda| \neq 1$, T is hyperbolic.

REFERENCES

- [A] O. Amann. *Groups of tree automorphisms and their unitary representations*. Dissertation ETH No. 15292, 2003.
- [BH] M. R. Bridson and A. Haefliger. *Metric spaces of nonpositive curvature*. Springer-Verlag, Berlin, 1999.
- [BM1] M. Burger and S. Mozes. Groups acting on trees: from local to global structure. *Inst. Hautes Études Sci. Publ. Math.*, 92:113–150, 2000.
- [BM2] M. Burger and S. Mozes. Lattices in products of trees. *Inst. Hautes Études Sci. Publ. Math.*, 92:151–194, 2000.
- [FN] A. Figá-Talamanca and C. Nebbia. *Harmonic analysis and representations theory for groups acting on homogeneous trees*. London Mathematical Society Lecture Notes, 162, Cambridge University Press, 1991.
- [G] M. Gromov. Asymptotic invariants of infinite groups. In *Geometric group theory, Vol. 2 (Sussex, 1991)*, pages 1–295. Cambridge Univ. Press, Cambridge, 1993.
- [K] A. Karlsson. Nonexpanding maps and Busemann functions. *Ergodic Theory Dynam. Systems*, 21(5):1447–1457, 2001.
- [KN] A. Karlsson and G. Noskov. Some groups having only elementary actions on metric spaces with hyperbolic boundaries. *Geom. Dedicata*, 104:119–137, 2004.
- [N] C. Nebbia. Classification of all irreducible unitary representations of the stabilizer of the horocycles of a tree. *Isr. J. Math.*, 70:343–351, 1990.
- [O] A. Yu. Ol’sanskii. SQ-universality of hyperbolic groups. *Mat. Sb.*, 186:119–132, 1995.
- [S] J.-P. Serre. *Arbres, amalgames et SL_2* . Asterisque.
- [Si] M. Simonnet. *Measures and probabilities*. Universitext. Springer-Verlag, New York, 1996. With a foreword by Charles-Michel Marle.
- [T] J. Tits. Sur le groupe des automorphismes d’un arbre. In *Essays on Topology and Related Topics*, pages 188–211. Springer, 1970.

- [V] A. Valette. Cocycles d'arbres et représentations uniformément bornées. *C. R. Acad. Sci. Paris Sér. I Math.*, 310 (10):703–708, 1990.

M.B.: FIM, ETHZ, ZÜRICH, CH-8092, SWITZERLAND
E-mail address: burger@math.ethz.ch

A.I.: DEPARTMENT MATHEMATIK, ETHZ, ZÜRICH, CH-8092, SWITZERLAND
E-mail address: iozzi@math.ethz.ch

N.M.: UNIVERSITY OF CHICAGO, CHICAGO, IL 60637, U. S. A.
E-mail address: monod@math.uchicago.edu