

SELF-CONSISTENT SOLUTION FOR THE POLARIZED VACUUM IN A NO-PHOTON QED MODEL

CHRISTIAN HAINZL, MATHIEU LEWIN, AND ERIC SÉRÉ

ABSTRACT. We study the Bogoliubov-Dirac-Fock model introduced by Chaix and Iracane (*J. Phys. B.*, 22, 3791–3814, 1989) which is a mean-field theory deduced from no-photon QED. The associated functional is bounded from below. In the presence of an external field, a minimizer, if it exists, is interpreted as the polarized vacuum and it solves a self-consistent equation.

In a recent paper, we proved the convergence of the iterative fixed-point scheme naturally associated with this equation to a global minimizer of the BDF functional, under some restrictive conditions on the external potential, the ultraviolet cut-off Λ and the bare fine structure constant α . In the present work, we improve this result by showing the existence of the minimizer by a variational method, for any cut-off Λ and without any constraint on the external field.

We also study the behaviour of the minimizer as Λ goes to infinity and show that the theory is “nullified” in that limit, as predicted first by Landau: the vacuum totally cancels the external potential. Therefore the limit case of an infinite cut-off makes no sense both from a physical and mathematical point of view.

Finally, we perform a charge and density renormalization scheme applying simultaneously to all orders of the fine structure constant α , on a simplified model where the exchange term is neglected.

1. INTRODUCTION

Despite the incredible predictive power of Quantum Electrodynamics (QED) its description in terms of perturbation theory restricts its range of applicability. In fact a mathematical consistent formulation is still unknown. We want to make a tiny step in that direction.

Following ideas of Chaix and Iracane [7], we study in this paper a model for the polarized vacuum in a Hartree-Fock type approximation. This so-called Bogoliubov-Dirac-Fock (BDF) model has been derived from no-photon QED in [7] as a possible cure to the fundamental problems associated with standard relativistic quantum chemistry calculations.

The vacuum polarization (VP) is, quoting [16], “one of the most interesting of the phenomena predicted by contemporary quantum electrodynamics”. Although it plays a minor role in the calculation of the Lamb-shift for the ordinary hydrogen atom (comparing to other electrodynamic phenomena), it is important for High- Z atoms [37, 49] and even plays a crucial role for muonic atoms [16, 21]. It also explains the production of electron-positron pairs, observed experimentally in heavy ions collision [2, 42, 31, 47, 20].

In [7], Chaix and Iracane noticed that the vacuum polarization effects are “necessary for the internal consistency of the relativistic mean-field theory and should therefore be taken into account in proper self-consistent calculations, independently of the magnitude of the physical effects” [7, page 3813]. Taking into account these

The authors are thankful to Robert Seiringer and Vladimir M. Shabaev for valuable comments. They acknowledge support through the European Union’s IHP network Analysis & Quantum HPRN-CT-2002-00277. E.S. acknowledges support from the Institut Universitaire de France.

effects, they restricted the no-photon QED Hamiltonian (normal-ordered with respect to the free electrons and positrons) to Bogoliubov transformations of the free vacuum. This allowed them to obtain a bounded-below energy, a property which is a huge advantage compared to the usual Dirac-Fock theory [54]: the Dirac-Fock energy is unbounded from below, which is the cause of important computational [7, 9] and theoretical [13, 38, 14, 15] problems .

In this paper, we show the existence of a global minimizer for the Bogoliubov-Dirac-Fock functional of Chaix-Iracane in the presence of an external field, which is interpreted as the polarized vacuum. This vacuum is represented by a projector of infinite rank which solves a *self-consistent equation*: it is the projector on the negative eigenspace of an effective mean-field Dirac operator taking into account the vacuum polarization potentials. This equation naturally leads to an iterative fixed-point procedure for solving it. In a previous work [25], we proved the convergence of such an iterative scheme to a global minimizer of the BDF functional, but under some assumptions on the external field and the ultraviolet cut-off. Our goal here is to show the existence of a minimizer without any restriction, by means of a direct – non constructive – minimization argument.

In the case where no external field is present, the free vacuum is already known to be a minimizer of the BDF energy, as shown by Chaix-Iracane-Lions [8] and Bach-Barbaroux-Helffer-Siedentop [4]. In [4], an external field is also considered but vacuum polarization is neglected: the model studied there is thus very different from the one considered by Chaix-Iracane in [7] and in the present paper.

Of course the vacuum case is only a first step in the study of the Chaix-Iracane model. In order to consider atoms and molecules, one has to minimize the BDF energy in a fixed charge sector, a much more complicated problem from a mathematical point of view. A minimizer would then solve a self-consistent equation which takes the form of the usual *unprojected Dirac-Fock equations*, perturbed by the vacuum polarization potentials.

To deal with divergencies, we impose a ultraviolet momentum cutoff Λ . Our only restriction on Λ is its finiteness. Additionally we study the behaviour of our solution when $\Lambda \rightarrow \infty$ and show that the model becomes meaningless since the vacuum density totally cancels the external potential. In physics, this “nullification” of the theory as the cut-off diverges has been first predicted by Landau *et al.* [33, 32, 1, 34] and later thoroughly studied by Pomeranchuk *et al.* [40].

We also discuss a simplified model in more detail, neglecting the exchange energy. For the corresponding self-consistent solution we perform a fully – to any order in the coupling constant α – consistent charge renormalization scheme. This procedure has already been performed in perturbation theory by means of Feynman diagrams, see e.g. [41, page 194] and [23]. In particular, we recover the well-known fact [32, 33, 1] (see also, e.g., [28, Eq. (7.18)]) that the physical (renormalized) coupling constant α_r is related to the bare α by a relation of the form

$$(1) \quad \alpha_r = \frac{\alpha}{1 + \alpha B_\Lambda}.$$

where $B_\Lambda \sim_{\Lambda \rightarrow \infty} 2/(3\pi) \log \Lambda$. Therefore the limit case of an infinite cut-off appears as unphysical [33] since it would correspond to $\alpha_r = 0$, which means no more electrostatic interactions.

The paper is organized as follows. In the next section, we recall the BDF model. Our main existence result is stated in Section 3, together with the behaviour of the solution as $\Lambda \rightarrow \infty$. In Section 4, we study the reduced model and interpret the self-consistent equation thanks to a renormalization of the charge and the density. Finally, the last section is devoted to the proof of our main results.

2. THE BOGOLIUBOV-DIRAC-FOCK MODEL

For the sake of clarity, we first briefly recall the Bogoliubov-Dirac-Fock (BDF) model introduced by Chaix-Iracane in [7] and studied in [25]. Details can be found in [25].

We use relativistic units $\hbar = c = 1$, set the particle mass equal to one and $\alpha = e^2/(4\pi)$. We emphasize that in the first part e represents the *bare* charge of the electron. We assume the presence of an external field $\varphi = n * \frac{1}{|\cdot|}$ describing one or more extended nuclei with overall charge density $n(x)$. We do not assume in this work that n is a non-negative function, since our model allows to treat the vacuum interacting with both matter and antimatter. We denote by $D^0 = \boldsymbol{\alpha} \cdot \mathbf{p} + \beta$ the free Dirac operator and by $D^\varphi := D^0 - \alpha\varphi$ the Dirac operator with external potential. Throughout the paper we use the notation $\chi_{(-\infty,0)}(H)$ to denote the projector on the negative spectral subspace of H . In the physical literature $\chi_{(-\infty,0)}(H)$ is often denoted as $\Lambda^-(H)$.

When the external field is not too strong, a good approximation is to use the Furry picture [18] in the Lamb-shift calculations of atomic bound states (see, e.g., [37, 49]). This means that, in order to evaluate corrections due to Vacuum Polarization, the dressed vacuum is represented by the projector associated with the negative spectrum of the Dirac operator with external potential D^φ

$$P^\varphi = \chi_{(-\infty;0)}(D^\varphi).$$

In reality, the polarized vacuum modifies the electrostatic field, and the virtual electrons react to the corrected field. This remark naturally leads to a self-consistent equation for the dressed vacuum of the form

$$P_{\text{scf}} = \chi_{(-\infty;0)}(D^\varphi + V_{\text{eff}})$$

where V_{eff} is an effective potential already including the Vacuum Polarization potentials. The BDF model [7] allows to derive such an effective potential V_{eff} in a *self-consistent* way, P_{scf} being interpreted as a minimizer in the class of Bogoliubov transformations of the free vacuum $P^0 = \chi_{(-\infty;0)}(D^0)$.

In practice, V_{eff} can be computed by a fixed point iterative procedure studied in details in [25]. If one starts the procedure from P^0 , the first iteration gives P^φ , and this explains why the Furry picture is a good approximation. But corrections to the Furry picture are necessary for high accuracy computations of electronic levels near heavy nuclei. These corrections can be interpreted as the second iteration in a Banach fixed-point algorithm (see, e.g., [37, section 8.2]).

Self-consistent equations leading to a fixed-point iterative scheme are well-known and widely used in full QED. The solutions of the Schwinger-Dyson equations [46, 12] involving the different four-dimensional Feynman propagators are usually found by means of perturbation theory. Our approach for the special case of the Hartree-Fock theory without photon studied in this paper is mathematically rigorous, non-perturbative and works for any charge Z of the external potential.

The momentum cutoff Λ is implemented in the Hilbert space

$$(2) \quad \mathfrak{H}_\Lambda = \{f \in L^2(\mathbb{R}^3, \mathbb{C}^4) \mid \text{supp } \hat{f} \subset B(0, \Lambda)\},$$

that is the space of spin valued functions whose Fourier transform has support inside a ball with radius Λ . Such a sharp cut-off does not allow to keep gauge invariance when photons are present. Since we neglect photons, we shall however use (2) for simplicity.

The space \mathfrak{H}_Λ can be decomposed as a direct sum of the negative and positive subspaces of the free Dirac operator D^0 , i.e. $\mathfrak{H}_\Lambda = \mathfrak{H}_\Lambda^0 \oplus \mathfrak{H}_\Lambda^+$ where $\mathfrak{H}_\Lambda^0 = P^0 \mathfrak{H}_\Lambda$ and $\mathfrak{H}_\Lambda^+ = (1 - P^0) \mathfrak{H}_\Lambda$, $P^0 = \chi_{(-\infty;0)}(D^0)$. The Fock space \mathcal{F} is built upon this

splitting as usual [7, 55]:

$$\mathcal{F} := \bigoplus_{n,m=1}^{\infty} \mathcal{F}_+^{(n)} \otimes \mathcal{F}_-^{(m)},$$

where $\mathcal{F}_+^{(n)} := \bigwedge_{i=1}^n \mathfrak{H}_+^0$ is the n -electron state subspace, $\mathcal{F}_-^{(m)} := \bigwedge_{j=1}^m C\mathfrak{H}_-^0$ is the m -positron state subspace, and $\mathcal{F}_+^{(0)} = \mathcal{F}_-^{(0)} = \mathbb{C}$. Here C is the charge conjugation operator [55]. The bare annihilation operators for electrons $a_0(f)$ and positrons $b_0(f)$ are then defined in the usual way [7, 55], for any $f \in \mathfrak{H}_\Lambda$. The field operator reads

$$\Psi(f) = a_0(f) + b_0^*(f).$$

The free vacuum $\Omega_0 = 1 \in \mathbb{C} \subset \mathcal{F}$ is characterized up to a phase by the properties $a_0(f)\Omega_0 = b_0(f)\Omega_0 = 0$ for any $f \in \mathfrak{H}_\Lambda$, and $\|\Omega_0\|_{\mathcal{F}} = 1$.

Let us now define the BDF class in the Fock space. Given a new (dressed) projector P , we define the dressed annihilation operators by $a_P(f) = \Psi((1-P)f)$ and $b_P(f) = \Psi^*(Pf)$. The associated dressed Ω_P is a state in the Fock space such that $a_P(f)\Omega_P = b_P(f)\Omega_P = 0$ for any $f \in \mathfrak{H}_\Lambda$, and $\|\Omega_P\|_{\mathcal{F}} = 1$. By the Shale-Stinespring Theorem [50], such an Ω_P is known to exist and is unique up to a phase, if and only if $P - P^0 \in \mathfrak{S}_2(\mathfrak{H}_\Lambda)$, the space of Hilbert-Schmidt operators on \mathfrak{H}_Λ (see also [30]). The state Ω_P can be expressed as a rotation of the free vacuum, $\Omega_P = \mathbb{U}\Omega_0$, \mathbb{U} being called a Bogoliubov transformation. An explicit formula for Ω_P can be found in lots of papers [55, 30, 43, 44, 47, 20]. The BDF class is therefore the subset of \mathcal{F}

$$\mathcal{B} := \{\Omega_P \mid P \text{ orth. projector, } P - P^0 \in \mathfrak{S}_2(\mathfrak{H}_\Lambda)\}.$$

The charge of Ω_P can be easily computed

$$\begin{aligned} (3) \quad \langle \Omega_P | \mathcal{Q} | \Omega_P \rangle &= \text{tr}(P^0(P - P^0)P^0) + \text{tr}((1 - P^0)(P - P^0)(1 - P^0)) \\ &= \text{tr}(Q^{--}) + \text{tr}(Q^{++}) \end{aligned}$$

where $Q = P - P^0 \in \mathfrak{S}_2(\mathfrak{H}_\Lambda)$ and $Q^{--} = P^0QP^0$, $Q^{++} = (1 - P^0)Q(1 - P^0)$. In (3), \mathcal{Q} is the usual charge operator acting on the Fock space \mathcal{F} [55, Eq. (10.52)],

$$\mathcal{Q} = \sum_{i \geq 1} a_0^*(f_i^+) a_0(f_i^+) - \sum_{i \geq 1} b_0^*(f_i^-) b_0(f_i^-),$$

$(f_i^+)_{i \geq 1}$ and $(f_i^-)_{i \geq 1}$ being respectively orthonormal basis of \mathfrak{H}_+^0 and \mathfrak{H}_-^0 .

Due to (3), we have introduced in [25] the notion of P^0 -trace class operators. We say $A \in \mathfrak{S}_2(\mathfrak{H}_\Lambda)$ is P^0 -trace class if the operators $A^{++} := (1 - P^0)A(1 - P^0)$ and $A^{--} := P^0AP^0$ are trace-class ($\in \mathfrak{S}_1(\mathfrak{H}_\Lambda)$), and we define the P^0 -trace of A by

$$(4) \quad \text{tr}_{P^0} A = \text{tr} A^{++} + \text{tr} A^{--}.$$

Notice, if A is even trace-class then $\text{tr}_{P^0} A = \text{tr} A$. In the following, we denote by $\mathfrak{S}_1^{P^0}(\mathfrak{H}_\Lambda)$ the set of all P^0 -trace class operators. Remark that by definition $\mathfrak{S}_1^{P^0}(\mathfrak{H}_\Lambda) \subset \mathfrak{S}_2(\mathfrak{H}_\Lambda)$.

We have shown in [25, Lemma 2] that any difference of two projectors satisfying the Shale-Stinespring criterion, $Q = P - P^0 \in \mathfrak{S}_2(\mathfrak{H}_\Lambda)$, is automatically in $\mathfrak{S}_1^{P^0}(\mathfrak{H}_\Lambda)$. The charge $\langle \Omega_P | \mathcal{Q} | \Omega_P \rangle = \text{tr}_{P^0}(Q)$ is therefore a well-defined number which indeed is always an integer, as proved in [25, Lemma 2]. The P^0 -trace is an adequate tool for describing charge sectors, without using the explicit expression of Ω_P which can be found in the literature.

In this paper, we study the case of the vacuum: namely we want to show the existence of a BDF state $\Omega_P \in \mathcal{B}$ with *lowest energy*, which we call a *BDF-stable vacuum*. For a small external field, this vacuum will not be charged but if the external field is strong enough, we could end up with a charged vacuum,

$\langle \Omega_P | \mathcal{Q} | \Omega_P \rangle = \text{tr}_{P^0}(P - P^0) \neq 0$. In order to study atoms or molecules, one has to minimize the energy in different charge sectors

$$\mathcal{B}_N := \{\Omega_P \in \mathcal{B} \mid \langle \Omega_P | \mathcal{Q} | \Omega_P \rangle = N\} \subset \mathcal{B}.$$

In this case, as explained in [7, section 4.2] (see also [25, Remark 6]), the electronic orbitals will solve the unprojected Dirac-Fock equations, perturbed by the vacuum polarization potentials. It is our goal to study this constrained minimization problem in the near future.

According to Chaix and Iracane [7, Formula (4.1)], the energy of a state Ω_P is defined using the renormalized Hamiltonian, acting on the Fock space \mathcal{F} ,

$$(5) \quad \mathbb{H} = \int dx : \Psi^*(x) D^\varphi \Psi(x) :_{P^0} + \frac{\alpha}{2} \int dx \int dy \frac{:\Psi^*(x)\Psi(x)\Psi^*(y)\Psi(y):_{P^0}}{|x-y|}$$

where $\Psi(x) = \sum_{i \geq 1} \Psi(f_i) f_i(x)$, $(f_i)_{i \geq 1}$ being an orthonormal basis of \mathfrak{H}_Λ . The choice of the normal ordering with respect to P^0 corresponds to subtracting the energy of the free Dirac sea P^0 and the interaction potentials involving P^0 . We emphasize that by this choice we make the assumption that the free vacuum is unobservable, as done by Dirac [10, 11], Heisenberg [27] and Weisskopf [57] (see also [26]). In principle, other choices could be made [36].

Evaluating the expectation value of Ω_P , we obtain [25, Appendix]

$$(6) \quad \langle \Omega_P | \mathbb{H} | \Omega_P \rangle = \mathcal{E}(Q)$$

where $Q = P - P^0 \in \mathfrak{S}_1^{P^0}(\mathfrak{H}_\Lambda)$ and \mathcal{E} is the Bogoliubov-Dirac-Fock energy

$$(7) \quad \mathcal{E}(Q) = \text{tr}_{P^0}(D^0 Q) - \alpha D(\rho_Q, n) + \frac{\alpha}{2} D(\rho_Q, \rho_Q) - \frac{\alpha}{2} \iint \frac{|Q(x, y)|^2}{|x-y|} dx dy.$$

Here $\rho_Q(x) = \text{tr}_{\mathbb{C}^4} Q(x, x)$ and

$$D(f, g) = 4\pi \int_{\mathbb{R}^3} \frac{\widehat{f}(k) \widehat{g}(k)}{|k|^2} dk.$$

Notice that the density ρ_Q is well defined due to the ultraviolet cut-off [25, Eq. (9)], and that $D(f, g) = \iint_{\mathbb{R}^6} \frac{f(x)g(y)}{|x-y|} dx dy$ when f and g are smooth enough.

As this is seen from (6), the energy of Ω_P only depends on $Q = P - P^0$, which is interpreted as the *renormalized one-body density matrix* of Ω_P .

3. EXISTENCE OF A BDF-STABLE POLARIZED VACUUM

Following a usual method for Hartree-Fock type theories [35, 3, 5], we may define and study the functional \mathcal{E} on the extended convex set

$$(8) \quad \mathcal{S}_\Lambda = \{Q \mid 0 \leq Q + P^0 \leq 1, Q \in \mathfrak{S}_1^{P^0}(\mathfrak{H}_\Lambda), \rho_Q \in \mathcal{C}\},$$

where \mathcal{C} is the so-called Coulomb space consisting of functions with finite Coulomb norm

$$\|\rho\|_{\mathcal{C}}^2 := D(\rho, \rho) = 4\pi \int_{\mathbb{R}^3} \frac{|\widehat{\rho}(k)|^2}{|k|^2} dk.$$

More precisely, \mathcal{C} is the Fourier inverse of the L^2 space with weight $1/|k|^2$.

As our main result we obtain that, for any Λ , \mathcal{E} is bounded-below and has a minimizer on \mathcal{S}_Λ , therefore there exists a BDF-stable vacuum.

Theorem 1. *Let $0 \leq \alpha < 4/\pi$, $n \in \mathcal{C}$. Then \mathcal{E} satisfies, for any $Q \in \mathcal{S}_\Lambda$,*

$$(9) \quad \mathcal{E}(Q) + \frac{\alpha}{2} D(n, n) \geq 0$$

and it is therefore bounded from below on S_Λ . Moreover, there exists a minimizer \bar{Q} of \mathcal{E} on S_Λ such that $\bar{P} = \bar{Q} + P^0$ is a projector satisfying the self-consistent equation

$$(10) \quad \bar{P} = \chi_{(-\infty, 0)} \left(D^0 - \alpha\varphi + \alpha\rho_{\bar{Q}} * \frac{1}{|\cdot|} - \alpha \frac{\bar{Q}(x, y)}{|x - y|} \right).$$

Additionally, if α and n satisfy

$$(11) \quad 0 \leq \alpha \frac{\pi}{4} \left\{ 1 - \alpha \left(\frac{\pi}{2} \sqrt{\frac{\alpha/2}{1 - \alpha\pi/4}} + \pi^{1/6} 2^{11/6} \right) \|n\|_{\mathcal{C}} \right\}^{-1} \leq 1,$$

then this global minimizer \bar{Q} is unique and the associated polarized vacuum is neutral:

$$\langle \Omega_{\bar{P}} | \mathcal{Q} | \Omega_{\bar{P}} \rangle = \text{tr}_{P^0}(\bar{Q}) = 0.$$

The proof of this result is given in Section 5.

Equation (10) corresponds to Dirac's picture that the "correct" vacuum \bar{P} should be the projector on the negative spectrum of an effective one-body Hamiltonian. In the case without external potential, $n = 0$, the free projector P^0 solves (10) and is the unique BDF-stable vacuum [8, 4].

Numerically the self-consistent solution of (10) could be evaluated by a fixed point algorithm, starting with P^0 . In [25] we proved the convergence of this algorithm to a BDF-stable vacuum solving (10), under reasonable restrictions of the form $\alpha\sqrt{\|n\|_{L^2}^2 + \|n\|_{\mathcal{C}}^2} \leq C_1$ and $\alpha\sqrt{\log \Lambda} \leq C_2$, using the Banach fixed point theorem. This proof is much more constructive than the direct variational proof which is given in Section 5. However, the result of [25] is local in the sense that it is valid for weak external potentials $\varphi = n * 1/|\cdot|$ only.

The condition (11) means that if the overall charge of the nuclei is not too big and α is small enough, the BDF-stable vacuum is unique and stays neutral, cf. [22, 24]. In general, the solution found in Theorem 1 can correspond to a charged vacuum.

There is an interesting symmetry property of the solutions of (10) when n is replaced by $-n$. Namely, if P is a solution of (10) with external density n , then $P' = Q' + P^0$ is a solution of (10) with external density $-n$, where $Q' = -CQC^{-1}$, C being the charge conjugation operator [55, page 14]. The two dressed vacua P and P' have the same BDF energies and satisfy $\rho_{Q'} = -\rho_Q$, as suggested by the intuition. For this symmetry between matter and antimatter to be true, it is essential to have the Fermi level at 0 and not at -1 (see, e.g., the comments of [44, page 197] about this fact).

In Theorem 1, the cut-off Λ can be chosen arbitrarily large and it is therefore natural to describe the behaviour of our solution as $\Lambda \rightarrow \infty$.

Theorem 2. *Let be $n \in \mathcal{C} \cap L^2(\mathbb{R}^3)$ and $0 \leq \alpha < 4/\pi$. Then the solution $\bar{Q}_\Lambda = \bar{P}_\Lambda - P^0$ obtained in Theorem 1 satisfies*

$$\| |D^0|^{1/2} \bar{Q}_\Lambda \|_{\mathfrak{S}_2} \rightarrow 0, \quad \alpha \| \rho_{\bar{Q}_\Lambda} - n \|_{\mathcal{C}} \rightarrow 0$$

as $\Lambda \rightarrow \infty$, and therefore

$$(12) \quad \lim_{\Lambda \rightarrow \infty} \min_{S_\Lambda} \mathcal{E} = -\frac{\alpha}{2} D(n, n).$$

In words, when $\Lambda \rightarrow \infty$, the vacuum polarization density *totally cancels the external density* n , for $\rho_{\bar{Q}_\Lambda} \rightarrow n$ in \mathcal{C} . But since $\bar{Q}_\Lambda = \bar{P}_\Lambda - P^0 \rightarrow 0$, this means that in the limit $\Lambda \rightarrow \infty$, \bar{Q}_Λ and $\rho_{\bar{Q}_\Lambda}$ become independent. Therefore, the minimization without cut-off makes no sense both from a mathematical and physical point of view. Indeed (12) easily implies that when no cut-off is imposed and when $\varphi \neq 0$,

the infimum of the functional \mathcal{E} is *not attained*. In physics, this “nullification” of the theory as the cut-off Λ diverges has been first suggested by Landau *et al.* [33, 32, 1, 34] and later studied by Pomeranchuk *et al.* [40].

In the next section, we propose a renormalization procedure in which we show an inequality of the form $\frac{2}{3\pi}\alpha_r \log \Lambda \leq 1$ where α_r is the physical (renormalized) coupling constant, different from α . With the usual value $\alpha_r \simeq \frac{1}{137}$, this leads to the physical bound $\Lambda \leq 10^{280}$ (in units of mc^2).

The proof of Theorem 2 is given in Section 5.

Remark. If n is smooth enough, it can be shown that

$$\| |D^0|^{1/2} \bar{Q}_\Lambda \|_{\mathfrak{S}_2} \leq C_1 (\log \Lambda)^{-1}, \quad \alpha \| \rho_{\bar{Q}_\Lambda} - n \|_{\mathcal{C}} \leq C_2 (\log \Lambda)^{-1}$$

for some constants C_1 and C_2 .

4. REDUCED ENERGY FUNCTIONAL AND CHARGE RENORMALIZATION

Recall up to now the charge was kept to be the bare one. Next we want to derive a renormalization scheme consistent to any order of α for the solution of our minimization problem. Note that this procedure is well known in perturbation theory, see e.g. [41, page 194] and [23].

We first simplify our BDF energy by neglecting the exchange term,

$$(13) \quad \mathcal{E}_{\text{red}}(Q) = \text{tr}_{P^0}(D^0 Q) - \alpha \int \rho_Q \varphi + \frac{\alpha}{2} D(\rho_Q, \rho_Q).$$

From a physical point of view this is quite natural, since the exchange term is usually treated together with a term describing the interaction with the photon field to form the standard electron *self-energy* that is a subject of the mass renormalization.

Notice that since $\mathcal{E}_{\text{red}} \geq \mathcal{E}$, the energy functional \mathcal{E}_{red} is obviously bounded from below on \mathcal{S}_Λ , by Theorem 1. We now state our

Theorem 3. *Let $0 \leq \alpha < 4/\pi$, $n \in \mathcal{C}$. Then \mathcal{E}_{red} possesses a minimizer \bar{Q} on \mathcal{S}_Λ , which satisfies*

$$(14) \quad \bar{Q} = \chi_{(-\infty, 0)} \left(D^0 - \alpha \varphi + \alpha \rho_{\bar{Q}} * \frac{1}{|\cdot|} \right) - P^0 + \gamma_0,$$

where γ_0 is a finite rank operator of the form

$$\gamma_0 = \sum_{i=1}^K n_i |\varphi_i\rangle \langle \varphi_i|, \quad 0 \leq n_i \leq 1,$$

$(\varphi_i)_{i=1}^K$ being an orthonormal basis of $\ker(D^0 - \alpha \varphi + \alpha \rho_{\bar{Q}} * 1/|\cdot|)$.

Additionally, if α and n satisfy

$$(15) \quad \alpha \pi^{1/6} 2^{11/6} \|n\|_{\mathcal{C}} < 1,$$

then this global minimizer \bar{Q} is unique and

$$\ker(D^0 - \alpha \varphi + \alpha \rho_{\bar{Q}} * 1/|\cdot|) = \{0\}$$

which implies

$$(16) \quad \bar{Q} = \chi_{(-\infty, 0)} \left(D^0 - \alpha \varphi + \alpha \rho_{\bar{Q}} * \frac{1}{|\cdot|} \right) - P^0.$$

The proof is much simpler than the one of Theorem 1: \mathcal{E}_{red} is now a coercive and convex continuous functional which is therefore weakly lower semi-continuous on the closed convex set \mathcal{S}_Λ , and possesses a minimizer. The proof that it satisfies the self-consistent equation (14) is the same as the one of Theorem 1, except that due to the absence of the exchange term, one is not always able to prove that $\bar{Q} + P^0$ is a projector, as usual in reduced Hartree-Fock type theories [53].

In order to perform our renormalization scheme we expand (16) in powers of α . Assuming that (15) holds, 0 is not in the spectrum of the mean-field operator $D^\varphi + \alpha\rho_{\bar{Q}} * 1/|\cdot|$ and we can use the resolvent representation [29, Section VI, Lemma 5.6] to derive from (16) the self-consistent equation for the VP-density $\rho_Q(x) = \text{tr}_{\mathbb{C}^4} Q(x, x)$

$$(17) \quad \rho_Q(x) = -\frac{1}{2\pi} \int_{-\infty}^{\infty} d\eta \text{tr}_{\mathbb{C}^4} \left[\frac{1}{D^0 - \alpha\varphi + \alpha\rho_Q * \frac{1}{|\cdot|} + i\eta} - \frac{1}{D^0 + i\eta} \right] (x, x).$$

Applying the resolvent equation

$$\frac{1}{A - \alpha B} - \frac{1}{A} = \alpha \frac{1}{A} B \frac{1}{A} + \alpha^2 \frac{1}{A} B \frac{1}{A} B \frac{1}{A} + \alpha^3 \frac{1}{A} B \frac{1}{A} B \frac{1}{A} B \frac{1}{A - \alpha B}$$

and using Furry's Theorem [17], telling us that the corresponding α^2 -term with two potentials vanish, we obtain

$$(18) \quad \rho_Q = \alpha F_1[\rho_Q - n] + F_3[\alpha\rho_Q - \alpha n]$$

with

$$F_3[\rho](x) = \int_{-\infty}^{\infty} d\eta \text{tr}_{\mathbb{C}^4} \left[\frac{1}{D^0 + i\eta} \rho * \frac{1}{|\cdot|} \frac{1}{D^0 + i\eta} \rho * \frac{1}{|\cdot|} \frac{1}{D^0 + i\eta} \rho * \frac{1}{|\cdot|} \frac{1}{D^0 - \alpha\rho + i\eta} \right] (x, x).$$

As realized first by Dirac [10, 11] and Heisenberg [27], cf. also [19], the term $F_1[\rho]$ plays a particular role since it is logarithmically ultraviolet divergent. Following, e.g., Pauli-Rose [39], one evaluates in Fourier representation

$$\widehat{F}_1[\rho](k) = -\hat{\rho}(k) B_\Lambda(k),$$

with

$$(19) \quad B_\Lambda(k) = \frac{1}{\pi} \int_0^{\frac{\Lambda}{\sqrt{1+\Lambda^2}}} \frac{z^2 - z^4/3}{1 - z^2} \frac{dz}{1 + |k|^2(1 - z^2)/4},$$

which can be decomposed into [39, Equ. (5)-(9)] $B_\Lambda(k) = B_\Lambda - C_\Lambda(k)$, with

$$(20) \quad B_\Lambda = B_\Lambda(0) = \frac{1}{\pi} \int_0^{\frac{\Lambda}{\sqrt{1+\Lambda^2}}} \frac{z^2 - z^4/3}{1 - z^2} dz = \frac{2}{3\pi} \log(\Lambda) - \frac{5}{9\pi} + \frac{2}{3\pi} \log 2 + O(1/\Lambda^2).$$

and

$$(21) \quad \lim_{\Lambda \rightarrow \infty} C_\Lambda(k) = C(k) = -\frac{1}{2\pi} \int_0^1 dx (1 - x^2) \log[1 + k^2(1 - x^2)/4],$$

which was first calculated by Serber and Uehling [48, 56].

Denote $\rho = \rho_Q - n$ the total density, then (18) reads in terms of ρ

$$(22) \quad \hat{\rho} + \hat{n} = -\alpha B_\Lambda \hat{\rho} + \alpha C_\Lambda(k) \hat{\rho} + \widehat{F}_3[\alpha\rho],$$

or equivalently

$$(23) \quad \alpha \hat{\rho} = -\alpha \hat{n} - \alpha^2 B_\Lambda \hat{\rho} + \alpha^2 C_\Lambda(k) \hat{\rho} + \alpha \widehat{F}_3[\alpha\rho]$$

and

$$(24) \quad \alpha \hat{\rho} = -\frac{\alpha}{1 + \alpha B_\Lambda} \hat{n} + \frac{\alpha}{1 + \alpha B_\Lambda} C_\Lambda(k) \alpha \hat{\rho} + \frac{\alpha}{1 + \alpha B_\Lambda} \widehat{F}_3[\alpha\rho].$$

To perform our renormalization scheme we fix as physical (renormalized) objects $\alpha_r \rho_r = \alpha\rho$, with (cf. [28, Equ. (7-18)])

$$(25) \quad \alpha_r = \frac{\alpha}{1 + \alpha B_\Lambda}.$$

Therefore we can rewrite the self-consistent equation (22) as

$$(26) \quad \alpha_r \hat{\rho}_r = -\alpha_r \hat{n} + \alpha_r^2 C_\Lambda(k) \hat{\rho}_r + \alpha_r \widehat{F}_3[\alpha_r \rho_r],$$

independently of the bare α . This equation uniquely defines the VP density only depending on the physical observable α_r , which is what we understand under consistent to any order. The α_r represents the dressed coupling constant, which is observed in experiment and whose value is approximately $1/137$. Notice that from formula (25), it follows that necessarily $\alpha_r B_\Lambda < 1$ and $\alpha_r B_\Lambda \rightarrow 1$ as $\Lambda \rightarrow \infty$. We emphasize that although in the literature the expression of α_r is sometimes expanded to get $\alpha_r \simeq \alpha(1 - \alpha B_\Lambda)$ leading to the condition $\alpha B_\Lambda < 1$, the real constraint indeed applies to the physically observed α_r and not the bare one.

Notice that equation (26) satisfied by $\alpha_r \rho_r$ is exactly the same as equation (23) satisfied by $\alpha \rho$, but with the logarithmically divergent term $\alpha^2 B_\Lambda \hat{\rho}$ dropped. Therefore, as usual in QED [12], the charge renormalization allows to simply justify the dropping of the divergent terms in the self-consistent equation. In practice [37], one would solve (26) with $\alpha_r \simeq 1/137$ and with $C_\Lambda(k)$ replaced by its limit $C(k)$.

Returning to the *effective* Hamiltonian $D^0 - \alpha\varphi + \alpha\rho_Q * 1/|\cdot|$ and inserting (26), i.e. expressing in terms of the physical objects, we obtain

$$(27) \quad D^0 + \alpha_r \rho_r * \frac{1}{|\cdot|} = D^0 - \alpha_r n * \frac{1}{|\cdot|} + V_{\text{eff}},$$

with

$$V_{\text{eff}} = \frac{2}{\pi^3} \mathcal{F}^{-1} \left[\frac{\alpha_r^2 C_\Lambda(k) \hat{\rho}_r(k) + \alpha_r \hat{F}_3(\alpha_r \rho_r)}{k^2} \right] (x)$$

the effective self-consistent potential, where \mathcal{F}^{-1} denotes the inverse Fourier transform. Notice, this equation is valid for any strength of the external potential. However, expanding ρ_r in α_r , we obtain to lowest order in α_r

$$\begin{aligned} V_{\text{eff}} &\simeq \alpha_r^2 \frac{2}{\pi^3} \mathcal{F}^{-1} \left[\frac{C_\Lambda(k) \hat{n}(k)}{k^2} \right] (x) \\ &\simeq \frac{\alpha_r^2}{3\pi} \int_1^\infty dt (t^2 - 1)^{1/2} \left[\frac{2}{t^2} + \frac{1}{t^4} \right] \int dx' e^{-2|x-x'|t} \frac{n(x')}{|x-x'|}, \end{aligned}$$

the Uehling potential [6]. Concerning a point like particle this potential was first written down in a closed form by Schwinger [45]. The next term in V_{eff} is of order $\alpha_r(\alpha_r Z)^3$. In principle all higher order corrections can be evaluated explicitly, which is not the task of our paper.

Finally we note that the convergence of the term in the right hand side of (22), in the case of the VP-density in the Furry picture, i.e. $\alpha \hat{F}_3(\alpha n)$, was shown in various papers. The most clarifying proof with respect to spurious third order contributions can probably be found in [52] (for earlier proofs, in particular corresponding to muonic atoms, we refer to the references in [52]). However the fact that this term, $\alpha \hat{F}_3(\alpha n)$, additionally gives rise to a well defined self-adjoint operator was recently proved in [26].

5. PROOF OF THEOREMS 1 AND 2

In this last Section, we give the proof of our main Theorems.

5.1. Proof of Theorem 1. The proof that \mathcal{E} is well-defined on \mathcal{S}_Λ can be found in details in [25, Theorem 1]. For simplicity, we extend \mathcal{E} to the closed convex set

$$\mathcal{S}'_\Lambda = \{Q \in \mathfrak{S}_2(\mathfrak{H}_\Lambda), 0 \leq Q + P^0 \leq 1, \rho_Q \in \mathcal{C}\}$$

of the Hilbert space $\mathcal{H} := \{Q \in \mathfrak{S}_2(\mathfrak{H}_\Lambda), \rho_Q \in \mathcal{C}\}$, by simply letting

$$\mathcal{E}(Q) = F(Q) - \alpha D(\rho_Q, n) + \frac{\alpha}{2} D(\rho_Q, \rho_Q),$$

$$(28) \quad F(Q) := \begin{cases} \operatorname{tr}_{P^0}(D^0 Q) - \frac{\alpha}{2} \iint \frac{|Q(x,y)|^2}{|x-y|} dx dy & \text{if } Q \in \mathfrak{S}'_1(P^0)(\mathfrak{H}_\Lambda) \\ +\infty & \text{if } Q \notin \mathfrak{S}'_1(P^0)(\mathfrak{H}_\Lambda) \end{cases}.$$

Let us recall the inequality established in [4]

$$F(Q) \geq (1 - \alpha\pi/4) \operatorname{tr}_{P^0}(D^0 Q) = (1 - \alpha\pi/4)(\operatorname{tr}(|D^0|Q^{++}) - \operatorname{tr}(|D^0|Q^{--}))$$

(notice that $Q^{++} \geq 0$ and $Q^{--} \leq 0$ when $Q \in \mathfrak{S}'_\Lambda$), which easily implies the bound (9) since

$$(29) \quad \mathcal{E}(Q) \geq (1 - \alpha\pi/4) \operatorname{tr}_{P^0}(D^0 Q) + \frac{\alpha}{2} \|\rho_Q - n\|_{\mathcal{C}}^2 - \frac{\alpha}{2} \|n\|_{\mathcal{C}}^2 \geq -\frac{\alpha}{2} \|n\|_{\mathcal{C}}^2.$$

This also easily shows that both F and \mathcal{E} are strongly lower semi-continuous and coercive on \mathfrak{S}'_Λ . We now prove that \mathcal{E} is indeed weakly lower semi-continuous (wlsc) on \mathfrak{S}'_Λ in \mathcal{H} , which will show the existence of a minimizer since \mathfrak{S}'_Λ is closed and convex, and therefore weakly closed.

Step 1: \mathcal{E} is wlsc on \mathfrak{S}'_Λ . Since the functional

$$Q \mapsto -\alpha D(\rho_Q, n) + \frac{\alpha}{2} D(\rho_Q, \rho_Q) = \frac{\alpha}{2} \|\rho_Q - n\|_{\mathcal{C}}^2 - \frac{\alpha}{2} \|n\|_{\mathcal{C}}^2$$

is easily seen to be wlsc on the convex set \mathfrak{S}'_Λ , it only remains to prove that F (defined in (28)) is wlsc on \mathfrak{S}'_Λ . To this end, we consider a weakly converging sequence $Q_n \rightarrow Q$ in \mathcal{H} , such that $Q_n \in \mathfrak{S}'_\Lambda$ for each n . If $\liminf_n F(Q_n) = \infty$, there is nothing to show and we can therefore assume that $(Q_n^{++})_{n \geq 1}$ and $(Q_n^{--})_{n \geq 1}$ are bounded in $\mathfrak{S}_1(\mathfrak{H}_\Lambda)$. Due to the cut-off Λ in Fourier space, $(Q_n(x, y))_{n \geq 1}$ is bounded in the Sobolev space $H^1(\mathbb{R}^6, \mathbb{C}^4 \otimes \mathbb{C}^4)$, $(\rho_{|D^0|Q_n^{++}})_{n \geq 1}$ and $(\rho_{|D^0|Q_n^{--}})_{n \geq 1}$ are bounded for instance in $H^1(\mathbb{R}^3, \mathbb{R})$. We may thus assume, up to a subsequence, that $Q_n(x, y) \rightarrow Q(x, y)$ in $L^2_{\text{loc}}(\mathbb{R}^6, \mathbb{C}^4 \otimes \mathbb{C}^4)$, that $\rho_{|D^0|Q_n^{++}} \rightarrow \rho_{|D^0|Q^{++}}$ and $\rho_{|D^0|Q_n^{--}} \rightarrow \rho_{|D^0|Q^{--}}$ in $L^1_{\text{loc}}(\mathbb{R}^3, \mathbb{R})$.

Let us now consider two real functions $\eta, \xi \in C^\infty([0; \infty); [0; 1])$ such that $\eta(t) = 1$ if $t \in [0; 1]$, $\eta(t) = 0$ if $t \geq 2$, $0 \leq \eta(t) \leq 1$ if $t \in [1; 2]$, and $\eta^2 + \xi^2 = 1$. We now define $\eta_R(x) := \eta(|x|/R)$ and $\xi_R(x) := \xi(|x|/R)$ for $x \in \mathbb{R}^3$. In the following, we also denote by η_R and ξ_R the multiplication operators by the functions η_R and ξ_R , acting on \mathfrak{H}_Λ .

Lemma 1. *We have $\|[\xi_R, |D^0|]\|_{\mathfrak{S}_\infty(\mathfrak{H}_\Lambda)} = O(1/R)$.*

Proof. We compute $\langle \psi | [\xi_R, |D^0|] | \chi \rangle$ in Fourier space, for some $\psi, \chi \in \mathfrak{H}_\Lambda$ (we use the notation $E(p) = \sqrt{1 + p^2}$):

$$\begin{aligned} \langle \psi | [\xi_R, |D^0|] | \chi \rangle &= \iint_{\mathbb{R}^6} \widehat{\xi}_R(p - q) \overline{\widehat{\psi}(p)} \widehat{\chi}(q) (E(q) - E(p)) dp dq \\ &= \iint_{\mathbb{R}^6} \widehat{\xi}_R(r) \overline{\widehat{\psi}\left(s + \frac{r}{2}\right)} \widehat{\chi}\left(s - \frac{r}{2}\right) \left(E\left(s - \frac{r}{2}\right) - E\left(s + \frac{r}{2}\right)\right) ds dr \end{aligned}$$

and therefore, using the inequality $|E(x) - E(x - y)| \leq |y|$, we obtain

$$|\langle \psi | [\xi_R, |D^0|] | \chi \rangle| \leq \left(\int_{\mathbb{R}^3} |r \widehat{\xi}_R(r)| dr \right) \|\widehat{\psi}\|_{L^2} \|\widehat{\chi}\|_{L^2}$$

and

$$\|[\xi_R, |D^0|]\|_{\mathfrak{S}_\infty(\mathfrak{H}_\Lambda)} \leq \int_{\mathbb{R}^3} |r \widehat{\xi}_R(r)| dr = \frac{C}{R} \int_{\mathbb{R}^3} |r \widehat{\xi}_1(r)| dr.$$

□

Using this Lemma, we may now write

$$\begin{aligned} \operatorname{tr}(|D^0|Q_n^{++}) &= \operatorname{tr}(\eta_R^2|D^0|Q_n^{++}) + \operatorname{tr}(\xi_R^2|D^0|Q_n^{++}) \\ &= \operatorname{tr}(\eta_R|D^0|Q_n^{++}\eta_R) + \operatorname{tr}(|D^0|\xi_R Q_n^{++}\xi_R) + \operatorname{tr}([\xi_R, |D^0|]Q_n^{++}\xi_R) \\ &= \operatorname{tr}(\eta_R|D^0|Q_n^{++}\eta_R) + \operatorname{tr}(|D^0|\xi_R Q_n^{++}\xi_R) + O(1/R) \end{aligned}$$

since

$$|\operatorname{tr}([\xi_R, |D^0|]Q_n^{++}\xi_R)| \leq \|[\xi_R, |D^0|]\|_{\mathfrak{S}_\infty(\mathfrak{H}_\Lambda)} \|Q_n^{++}\|_{\mathfrak{S}_1(\mathfrak{H}_\Lambda)} = O(1/R)$$

by Lemma 1 and since by assumption $(Q_n^{++})_{n \geq 1}$ is bounded in $\mathfrak{S}_1(\mathfrak{H}_\Lambda)$. With the same argument for Q_n^{--} , we obtain

$$\begin{aligned} \operatorname{tr}_{P^0}(D^0 Q_n) &= \operatorname{tr}(\eta_R|D^0|Q_n^{++}\eta_R) - \operatorname{tr}(\eta_R|D^0|Q_n^{--}\eta_R) \\ &\quad + \operatorname{tr}(|D^0|\xi_R Q_n^{++}\xi_R) - \operatorname{tr}(|D^0|\xi_R Q_n^{--}\xi_R) + O(1/R). \end{aligned}$$

On the other hand, we have

$$\begin{aligned} \iint \frac{|Q_n(x, y)|^2}{|x - y|} dx dy &= \iint \frac{\eta_R(x)^2 \eta_{3R}(y)^2 |Q_n(x, y)|^2}{|x - y|} dx dy \\ &\quad + \iint \frac{\xi_R(x)^2 |Q_n(x, y)|^2}{|x - y|} dx dy + O(1/R) \end{aligned}$$

since

$$\iint \frac{\eta_R(x)^2 \xi_{3R}(y)^2 |Q_n(x, y)|^2}{|x - y|} dx dy \leq \frac{\|Q_n\|_{\mathfrak{S}_2(\mathfrak{H}_\Lambda)}^2}{R}.$$

We therefore obtain

$$\begin{aligned} F(Q_n) &= \operatorname{tr}(\eta_R|D^0|Q_n^{++}\eta_R) - \operatorname{tr}(\eta_R|D^0|Q_n^{--}\eta_R) \\ &\quad - \frac{\alpha}{2} \iint \frac{\eta_R(x)^2 \eta_{3R}(y)^2 |Q_n(x, y)|^2}{|x - y|} dx dy + \operatorname{tr}(|D^0|\xi_R Q_n^{++}\xi_R) - \operatorname{tr}(|D^0|\xi_R Q_n^{--}\xi_R) \\ &\quad - \frac{\alpha}{2} \iint \frac{\xi_R(x)^2 |Q_n(x, y)|^2}{|x - y|} dx dy + O(1/R). \end{aligned}$$

Notice now that $0 \leq Q_n + P^0 \leq 1$ implies $|Q_n|^2 \leq Q_n^{++} - Q_n^{--}$ (see [4]). We now localize this inequality to obtain $\xi_R|Q_n|^2\xi_R \leq \xi_R Q_n^{++}\xi_R - \xi_R Q_n^{--}\xi_R$. By Kato's inequality [4], we now have

$$\begin{aligned} \iint \frac{\xi_R(x)^2 |Q_n(x, y)|^2}{|x - y|} dx dy &\leq \frac{\pi}{2} \operatorname{tr}(|D^0|\xi_R Q_n^2 \xi_R) \\ &\leq \frac{\pi}{2} (\operatorname{tr}(|D^0|\xi_R Q_n^{++}\xi_R) - \operatorname{tr}(|D^0|\xi_R Q_n^{--}\xi_R)) \end{aligned}$$

and therefore, since $(1 - \alpha\pi/4) \geq 0$,

$$\begin{aligned} F(Q_n) &\geq \operatorname{tr}(\eta_R|D^0|Q_n^{++}\eta_R) - \operatorname{tr}(\eta_R|D^0|Q_n^{--}\eta_R) \\ &\quad - \frac{\alpha}{2} \iint \frac{\eta_R(x)^2 \eta_{3R}(y)^2 |Q_n(x, y)|^2}{|x - y|} dx dy + O(1/R). \end{aligned}$$

Passing now to the limit as $n \rightarrow \infty$ and using the local compactness of $Q_n(x, y)$ in $L_{\text{loc}}^2(\mathbb{R}^6)$ and $\rho_{|D^0|Q_n^{++}}, \rho_{|D^0|Q_n^{--}}$ in $L_{\text{loc}}^1(\mathbb{R}^3)$, we obtain

$$\begin{aligned} \liminf_{n \rightarrow \infty} F(Q_n) &\geq \operatorname{tr}(\eta_R|D^0|Q^{++}\eta_R) - \operatorname{tr}(\eta_R|D^0|Q^{--}\eta_R) \\ &\quad - \frac{\alpha}{2} \iint \frac{\eta_R(x)^2 \eta_{3R}(y)^2 |Q(x, y)|^2}{|x - y|} dx dy + O(1/R). \end{aligned}$$

If we now let $R \rightarrow \infty$, we obtain $\liminf_{n \rightarrow \infty} F(Q_n) \geq F(Q)$ and therefore F is wsc on \mathcal{S}'_Λ .

Step 2: at least one of the minimizers satisfies (10). In the previous step, we have shown the existence of a minimizer. It now remains to show that one of them indeed satisfies (10).

Lemma 2. *Let \bar{Q} be a minimizer of \mathcal{E} in S'_Λ . Then either $\bar{Q} + P^0$ is a projector, or*

$$(30) \quad \bar{Q} + P^0 = \bar{P} + \mu|f\rangle\langle f|,$$

where \bar{P} is a projector, $\mu \in (0; 1)$ and $f \in \ker(D_{\bar{Q}})$, with

$$D_{\bar{Q}} := D^0 - \alpha\varphi + \alpha\rho_{\bar{Q}} * \frac{1}{|\cdot|} - \alpha \frac{\bar{Q}(x, y)}{|x - y|}.$$

Proof. Our proof is inspired by classical arguments already used in the Hartree-Fock theory [35, 3].

Notice that since \bar{Q} is compact, $\bar{Q} + P^0$ is a compact perturbation of P^0 and therefore its essential spectrum is $\sigma_{\text{ess}}(\bar{Q} + P^0) = \{0, 1\}$, meaning that $\sigma(\bar{Q} + P^0) \cap (0; 1)$ only contains eigenvalues of finite multiplicity accumulating at $\{0, 1\}$. Let us assume that $\bar{Q} + P^0$ possesses two different eigenvectors φ_1, φ_2 :

$$\bar{Q} + P^0 = \lambda_1|\varphi_1\rangle\langle\varphi_1| + \lambda_2|\varphi_2\rangle\langle\varphi_2| + G$$

where $\lambda_1, \lambda_2 \in (0; 1)$ and $G\varphi_1 = G\varphi_2 = 0$. We now introduce $\bar{Q}_\epsilon := \bar{Q} + \epsilon|\varphi_1\rangle\langle\varphi_1| - \epsilon|\varphi_2\rangle\langle\varphi_2|$ which belongs to S_Λ for ϵ small enough and compute

$$\mathcal{E}(\bar{Q}_\epsilon) = \mathcal{E}(\bar{Q}) + \epsilon(\langle\varphi_1|D_{\bar{Q}}|\varphi_1\rangle - \langle\varphi_2|D_{\bar{Q}}|\varphi_2\rangle) - \epsilon^2 \frac{\alpha}{2} \iint \frac{|\varphi_1 \wedge \varphi_2(x, y)|^2}{|x - y|} dx dy.$$

Therefore, using either the first order term in ϵ if it does not vanish, or the second order term, we can always decrease the energy. This is a contradiction which implies that $\sigma(\bar{Q} + P^0) \cap (0; 1)$ contains at most one eigenvalue of multiplicity 1 and thus

$$\bar{Q} + P^0 = \bar{P} + \mu|f\rangle\langle f|$$

where \bar{P} is a projector and $\mu \in [0; 1)$. If $\mu \neq 0$, using the same type of variation $\bar{Q}_\epsilon := \bar{Q} + \epsilon|f\rangle\langle f|$, we easily show that indeed $f \in \ker(D_{\bar{Q}})$. \square

If \bar{Q} is a minimizer of the form (30), we now see that

$$\begin{aligned} \mathcal{E}(\bar{Q}) &= \mathcal{E}(\bar{P} - P^0) + \mu\langle D_{\bar{P}-P^0} f, f \rangle \\ &= \mathcal{E}(\bar{P} - P^0) + \mu\langle D_{\bar{Q}} f, f \rangle \\ &= \mathcal{E}(\bar{P} - P^0) \end{aligned}$$

and therefore $\bar{P} - P^0$ is also a minimizer of \mathcal{E} (i.e. \bar{P} is BDF-stable vacuum). In [25, proof of Theorem 2], we have already shown that a minimizer of \mathcal{E} on S_Λ taking the form $\bar{P} - P^0$ where \bar{P} is an orthogonal projector, is indeed a solution of the self-consistent equation (10).

Step 3: uniqueness of the global minimizer of \mathcal{E} under the condition (11). Due to [25, Theorem 2], we know that the global minimizer \bar{Q} of \mathcal{E} is unique if $D_{\bar{Q}}$ satisfies

$$d|D_{\bar{Q}}| \geq |D^0|$$

for some d such that $\alpha d\pi/4 \leq 1$.

We know that $\mathcal{E}(\bar{Q}) \leq \mathcal{E}(0) = 0$ and therefore, by an argument similar to (29),

$$(31) \quad \left(\frac{2}{\pi} - \frac{\alpha}{2}\right) \iint_{\mathbb{R}^6} \frac{|\bar{Q}(x, y)|^2}{|x - y|} dx dy + \frac{\alpha}{2} \|\rho_{\bar{Q}} - n\|_{\mathcal{C}}^2 \leq \frac{\alpha}{2} \|n\|_{\mathcal{C}}^2$$

and thus

$$(32) \quad \iint_{\mathbb{R}^6} \frac{|\bar{Q}(x, y)|^2}{|x - y|} dx dy \leq \frac{\alpha\pi/4}{1 - \alpha\pi/4} \|n\|_{\mathcal{C}}^2$$

$$(33) \quad \|\rho_{\bar{Q}} - n\|_c \leq \|n\|_c.$$

Recall that $D_{\bar{Q}} = D^0 + \alpha\varphi'_{\bar{Q}} - \alpha R_{\bar{Q}}$ where $\varphi'_{\bar{Q}} = (\rho_{\bar{Q}} - n) * 1/|\cdot|$ and $R_{\bar{Q}}$ is the operator with kernel $\bar{Q}(x, y)/|x - y|$. Now, we have

$$\left\| \varphi'_{\bar{Q}} \frac{1}{|D^0|} \right\|_{\mathfrak{S}_\infty(\mathfrak{H}_\Lambda)} \leq \left\| \varphi'_{\bar{Q}} \frac{1}{|D^0|} \right\|_{\mathfrak{S}_6(\mathfrak{H}_\Lambda)} \leq (2\pi)^{-1/2} \|\varphi'_{\bar{Q}}\|_{L^6} \|E(\cdot)^{-1}\|_{L^6}$$

where we recall that $E(p) = \sqrt{1 + p^2}$, and by [51, Theorem 4.1]. Therefore

$$\left\| \varphi'_{\bar{Q}} \frac{1}{|D^0|} \right\|_{\mathfrak{S}_\infty(\mathfrak{H}_\Lambda)} \leq S_6 \|\varphi'_{\bar{Q}}\|_{L^6} \leq S_6 C_6 \|\nabla \varphi'_{\bar{Q}}\|_{L^2} = (4\pi) S_6 C_6 \|\rho_{\bar{Q}} - n\|_c$$

with $S_6 = 2^{-5/6} 3^{1/6} \pi^{-1/6}$ and where $C_6 = 3^{-1/6} 2^{2/3} \pi^{-2/3}$ is the Sobolev constant for the inequality $\|f\|_{L^6(\mathbb{R}^3)} \leq C_6 \|\nabla f\|_{L^2(\mathbb{R}^3)}$. Due to (33), this shows that

$$|\varphi'_{\bar{Q}}| \leq \kappa \|n\|_c |D^0|,$$

where $\kappa = (4\pi) S_6 C_6 = \pi^{1/6} 2^{11/6}$. On the other hand, we know from [25, Proof of Lemma 4] that

$$|R_{\bar{Q}}| \leq \sqrt{\frac{\pi}{2} \iint_{\mathbb{R}^6} \frac{|\bar{Q}(x, y)|^2}{|x - y|} dx dy} |D^0|$$

and therefore, using (32),

$$|R_{\bar{Q}}| \leq \frac{\pi}{2} \sqrt{\frac{\alpha/2}{1 - \alpha\pi/4}} \|n\|_c |D^0|.$$

As a conclusion, when

$$\alpha \left(\frac{\pi}{2} \sqrt{\frac{\alpha/2}{1 - \alpha\pi/4}} + \kappa \right) \|n\|_c < 1,$$

$D_{\bar{Q}}$ fulfills $d|D_{\bar{Q}}| \geq |D^0|$ with

$$d = \left\{ 1 - \alpha \left(\frac{\pi}{2} \sqrt{\frac{\alpha/2}{1 - \alpha\pi/4}} + \kappa \right) \|n\|_c \right\}^{-1}.$$

Applying now [25, Theorem 2], we obtain that the minimizer \bar{Q} is unique when $\alpha d\pi/4 \leq 1$, i.e. under the condition (11).

Assuming now that (11) holds, let us show that the unique BDF-stable vacuum \bar{P} is not charged. To this end, we define, for $t \in [0; 1]$,

$$\bar{Q}(t) = \chi_{(-\infty; 0)} \left(D^0 + \alpha t (\rho_{\bar{Q}} - n) * \frac{1}{|\cdot|} - \alpha t \frac{\bar{Q}(x, y)}{|x - y|} \right) - P^0.$$

$t \mapsto \bar{Q}(t)$ is a continuous function for the $\mathfrak{S}_2(\mathfrak{H}_\Lambda)$ topology, since by the previous estimates $D_{\bar{Q}}(t) = D^0 + \alpha t (\rho_{\bar{Q}} - n) * \frac{1}{|\cdot|} - \alpha t \frac{\bar{Q}(x, y)}{|x - y|}$ possesses a gap around 0, uniformly in $t \in [0; 1]$. This implies that

$$q : t \mapsto \text{tr}_{P^0}(Q(t)) = \text{tr}(Q(t)^3)$$

is continuous on $[0; 1]$, by [25, Lemma 2]. Since $q(0) = 0$ and $q(t)$ is an integer for any $t \in [0; 1]$, we therefore deduce that

$$q(1) = \langle \Omega_{\bar{P}} | \mathcal{Q} | \Omega_{\bar{P}} \rangle = \text{tr}_{P^0}(\bar{Q}) = 0.$$

This ends the proof of Theorem 1. \square

5.2. Proof of Theorem 2. We first prove (12) which will easily imply that \bar{Q}_Λ obtained by Theorem 1 behaves at stated as $\Lambda \rightarrow \infty$, due to (29). To this end, we introduce

$$Q_\Lambda := \chi_{(-\infty;0)} \left(D^0 - \alpha n_\Lambda * \frac{1}{|\cdot|} \right) - P^0 \in \mathcal{S}_\Lambda,$$

$$\widehat{n}_\Lambda(k) := \frac{\widehat{n}(k)}{1 + \alpha B_\Lambda(k)}$$

where we recall that $B_\Lambda(k) = B_\Lambda - C_\Lambda(k)$ is defined in (19). We now show that

$$\lim_{\Lambda \rightarrow \infty} \mathcal{E}(Q_\Lambda) = -\frac{\alpha}{2} D(n, n),$$

which will imply (12), by (9).

Let us now compute $\rho_\Lambda := \rho_{Q_\Lambda}$. By (22), ρ_Λ satisfies

$$\widehat{\rho}_\Lambda(k) = \frac{\alpha}{1 + \alpha B_\Lambda(k)} \widehat{n}(k) B_\Lambda(k) + \widehat{F}_3[\alpha n_\Lambda](k),$$

and therefore

$$\widehat{\rho}_\Lambda(k) - \widehat{n}(k) = -\widehat{n}_\Lambda(k) + \widehat{F}_3[\alpha n_\Lambda](k).$$

When $\alpha > 0$, since $(1 + \alpha B_\Lambda(k))^{-1} \rightarrow 0$ a.e., we obtain by Lebesgue's dominated convergence Theorem that $\|n_\Lambda\|_{\mathcal{C} \cap L^2} \rightarrow 0$ as $\Lambda \rightarrow \infty$. By the fixed-point estimates of [25] in the case of the reduced model (they are then independent on the cut-off Λ as this can be seen from the proof of [25, Theorem 3]), it is known that F_3 is continuous at 0 for the $\mathcal{C} \cap L^2$ topology. We therefore obtain

$$\lim_{\Lambda \rightarrow \infty} \|\rho_\Lambda - n\|_{\mathcal{C} \cap L^2} = 0.$$

On the other hand, we also know from the bounds proved in [25], that

$$\mathrm{tr}_{P^0}(D^0 Q_\Lambda)^{1/2} = \mathrm{tr}(|D^0| Q_\Lambda^2)^{1/2} \leq C\alpha \|\rho_\Lambda - n\|_{\mathcal{C} \cap L^2}$$

for some constant C independent of Λ . Therefore

$$\lim_{\Lambda \rightarrow \infty} \mathcal{E}(Q_\Lambda) = -\frac{\alpha}{2} D(n, n)$$

which ends the proof of Theorem 2. \square

REFERENCES

1. A.A. Abrikosov, I. Halatnikov and L.D. Landau. On the quantum theory of fields. *Nuovo Cim. Suppl.*, 3: 80–104, 1956. Reprinted in *Collected papers of L.D. Landau*, edited by D. Ter Haar, Pergamon Press, 1965.
2. A. Aste, G. Baur, K. Hencken, D. Trautmann and G. Scharf. Electron-positron pair production in the external electromagnetic field of colliding relativistic heavy ions. *Eur. Phys. J. C*, 23(3): 545–550, 2002.
3. V. Bach. Error bound for the Hartree-Fock energy of atoms and molecules. *Comm. Math. Phys.*, 147: 527–548, 1992.
4. V. Bach, J.-M. Barbaroux, B. Helffer and H. Siedentop. On the Stability of the Relativistic Electron-Positron Field. *Commun. Math. Phys.*, 201: 445–460, 1999.
5. V. Bach, E.H. Lieb and J.P. Solovej. Generalized Hartree-Fock theory and the Hubbard model. *J. Statist. Phys.*, 76 (1-2): 3–89, 1994.
6. E. Borie and G.A. Rinker. *Rev. Mod. Phys.*, 54: 67 (1982)
7. P. Chaix and D. Iracane. From quantum electrodynamics to mean field theory: I. The Bogoliubov-Dirac-Fock formalism. *J. Phys. B.*, 22(23): 3791–3814, 1989.
8. P. Chaix, D. Iracane and P.L. Lions. From quantum electrodynamics to mean field theory: II. Variational stability of the vacuum of quantum electrodynamics in the mean-field approximation. *J. Phys. B.*, 22(23): 3815–3828, 1989.
9. P. Chaix. *Une Méthode de Champ Moyen Relativiste et Application à l'Etude du Vide de l'Electrodynamique Quantique*. PhD Thesis, University Paris VI, 1990.
10. P.-A.-M. Dirac. Théorie du positron. Solvay report, pages 203–212. Paris: Gauthier-Villars. XXV, 353 S., 1934 (reprinted in *Selected papers on Quantum Electrodynamics*, edited by J. Schwinger, Dover, 1958).
11. P.A.M. Dirac. Discussion of the infinite distribution of electrons in the theory of the positron. *Proc. Camb. Philos. Soc.*, 30: 150–163, 1934.

12. F. J. Dyson. The S Matrix in Quantum Electrodynamics. *Phys. Rev.*, 75 (11): 1736–1755, 1949.
13. M.J. Esteban, E. Séré. Solutions of the Dirac-Fock Equations for Atoms and Molecules. *Comm. Math. Phys.*, 203: 499–530, 1999.
14. M.J. Esteban, E. Séré. Nonrelativistic limit of the Dirac-Fock equations. *Ann. Henri Poincaré*, 2(5): 941–961, 2001.
15. M.J. Esteban, E. Séré. A max-min principle for the ground state of the Dirac-Fock functional. *Contemp. Math.*, 307: 135–139, 2002.
16. L.L. Foldy and E. Eriksen. Some Physical Consequences of Vacuum Polarization. *Phys. Rev.*, 95(4): 1048–1051, 1954.
17. W.H. Furry. A symmetry theorem in the positron theory. *Phys. Rev.*, 51: 125, 1937.
18. W.H. Furry. On Bound States and Scattering in Positron Theory. *Phys. Rev.*, 81(1): 115–124, 1951.
19. W.H. Furry and J.R. Oppenheimer. On the Theory of the Electron and Positive. *Phys. Rev., II. Ser.*, 45: 245–262, 1934.
20. H. Fierz and G. Scharf. Particle interpretation for external field problems in QED. *Helv. Phys. Acta*, 52: 437–453, 1979.
21. R. Glauber, W. Rarita, and P. Schwed. Vacuum polarization effects on energy levels in μ -mesonic atoms. *Physical Review*, 120(2): 609–613, 1960.
22. W. Greiner, B. Müller, and J. Rafelski. *Quantum Electrodynamics of Strong Fields*. Texts and Monographs in Physics. Springer-Verlag, 1st edition, 1985.
23. A. Hamm and D. Schütte. How to remove divergences from the QED-Hartree approximation. *J. Phys. A: Math. Gen.*, 23: 3969–3982, 1990.
24. C. Hainzl. On the Vacuum Polarization Density caused by an External Field. *Ann. Henri Poincaré*, 5(6): 1137–1157, 2004.
25. C. Hainzl, M. Lewin and E. Séré. Existence of a stable polarized vacuum in the Bogoliubov-Dirac-Fock approximation, to appear in *Commun. Math. Phys.*
26. C. Hainzl and H. Siedentop. Non-Perturbative Mass and Charge Renormalization in Relativistic no-photon Quantum Electrodynamics. *Comm. Math. Phys.*, 243: 241–260, 2003.
27. W. Heisenberg. Bemerkungen zur Diracschen Theorie des Positrons. *Z. Phys.*, 90: 209–231, 1934.
28. C. Itzykson and J.-B. Zuber. *Quantum Field Theory*. McGraw-Hill, New York, 1980.
29. T. Kato. Perturbation Theory for Linear Operators, volume 132 of *Grundlehren der mathematischen Wissenschaften*. Springer-Verlag, Berlin, 1 edition, 1966.
30. M. Klaus and G. Scharf. The regular external field problem in quantum electrodynamics. *Helv. Phys. Acta*, 50: 779–802, 1977.
31. M. Klaus and G. Scharf. Vacuum Polarization in Fock Space. *Helv. Phys. Acta*, 50: 803–814, 1977.
32. L.D. Landau. On the quantum theory of fields. *Bohr Volume*, Pergamon Press, Oxford, 1955. Reprinted in *Collected papers of L.D. Landau*, edited by D. Ter Haar, Pergamon Press, 1965.
33. L.D. Landau and I. Pomeranchuk. On point interactions in quantum electro dynamics. *Dokl. Akad. Nauk SSSR*, 102: 489–492, 1955. Reprinted in *Collected papers of L.D. Landau*, edited by D. Ter Haar, Pergamon Press, 1965.
34. L.D. Landau. Fundamental problems. *Pauli Memorial Volume*, Interscience, p 245, 1960. Reprinted in *Collected papers of L.D. Landau*, edited by D. Ter Haar, Pergamon Press, 1965.
35. E.H. Lieb. Variational Principle for Many-Fermion Systems. *Phys. Rev. Lett.*, 46: 457–459, 1981.
36. E.H. Lieb and H. Siedentop. Renormalization of the regularized relativistic electron-positron field. *Comm. Math. Phys.*, 213 (3): 673–683, 2000.
37. P. J. Mohr, G. Plunien and G. Soff. QED corrections in Heavy Atoms. *Phys. Rep.*, 293: 227–369, 1998.
38. E. Paturel. Solutions of the Dirac-Fock equations without projector. *Ann. Henri Poincaré*, 1(6): 1123–1157, 2000.
39. W. Pauli and M.E. Rose. Remarks on the Polarization Effects in the Positron Theory. *Phys. Rev II*, 49: 462–465, 1936.
40. I.Ya Pomeranchuk, V.V. Sudakov and K.A. Ter-Martirosyan. Vanishing of renormalized charges in field theories with point interaction. *Phys. Rev.*, 103 (3): 784–802, 1956.
41. P.-G. Reinhard, W. Greiner and H. Arenhvel. Electrons in Strong External Fields. *Nucl. Phys. A*, 166: 173–197, 1971.
42. J. Reinhardt, B. Müller and W. Greiner. Theory of positron production in heavy-ion collision. *Phys. Rev. A*, 24(1): 103–128, 1981.
43. S.N.M. Ruijsenaars. On Bogoliubov transformations for systems of relativistic charged particles. *J. Math. Phys.*, 18(3): 517–526, 1977.

44. G. Scharf and H.P. Seipp. Charged Vacuum, Spontaneous Positron Production and All That. *Phys. Lett.*, 108B(3): 196–198, 1982.
45. J. Schwinger. Quantum Electrodynamics II. Vacuum Polarization and Self-Energy. *Phys. Rev., II. Ser.*, 75: 651–679, 1949.
46. J. Schwinger. On the Green’s Function of Quantized Fields. II. *Proc. Nat. Acad. Sci.*, 37(7):455–459, 1951.
47. H.P. Seipp. On the S -operator for the external field problem of QED. *Helv. Phys. Acta*, 55: 1–28, 1982.
48. R. Serber. Linear modifications in the Maxwell field equations. *Phys. Rev., II. Ser.*, 48: 49–54, 1935.
49. V.M. Shabaev. Two-time Green’s function method in quantum electrodynamics of high- Z few-electron atoms. *Phys. Rep.*, 356, 119–228 (2002)
50. D. Shale and W.F. Stinespring. Spinor representations of infinite orthogonal groups. *J. Math. and Mech.*, 14: 315–322, 1965.
51. B. Simon. *Trace Ideals and their Applications*. Vol 35 of *London Mathematical Society Lecture Notes Series*. Cambridge University Press, 1979.
52. G. Soff and P. Mohr. *Phys. Rev. A*, 38, 5066 (1988)
53. J.P. Solovej. Proof of the ionization conjecture in a reduced Hartree-Fock model. *Invent. Math.*, 104(2): 291–311, 1994.
54. B. Swirles. The relativistic self-consistent field. *Proc. Roy. Soc. A*, 152: 625–649, 1935.
55. B. Thaller. *The Dirac Equation*. Springer Verlag, 1992.
56. E.A. Uehling. Polarization effects in the positron theory. *Phys. Rev., II. Ser.*, 48: 55–63, 1935.
57. V. Weisskopf. Über die Elektrodynamik des Vakuums auf Grund der Quantentheorie des Elektrons. *Math.-Fys. Medd., Danske Vid. Selsk.*, 16(6):1–39, 1936 (reprinted in *Selected papers on Quantum Electrodynamics*, edited by J. Schwinger, Dover, 1958).

CEREMADE, UNIVERSITÉ PARIS-DAUPHINE, PLACE DU MARÉCHAL DE LATTRE DE TASSIGNY,
F-75775 PARIS CEDEX 16, FRANCE & LABORATOIRE DE MATHÉMATIQUES PARIS-SUD-BAT 425,
F-91405 ORSAY CEDEX

E-mail address: `hainzl@ceremade.dauphine.fr`

CEREMADE, UNIVERSITÉ PARIS-DAUPHINE, PLACE DU MARÉCHAL DE LATTRE DE TASSIGNY,
F-75775 PARIS CEDEX 16, FRANCE.

E-mail address: `lewin@ceremade.dauphine.fr`

CEREMADE, UNIVERSITÉ PARIS-DAUPHINE, PLACE DU MARÉCHAL DE LATTRE DE TASSIGNY,
F-75775 PARIS CEDEX 16, FRANCE.

E-mail address: `sere@ceremade.dauphine.fr`